

ON A CLASS NUMBER FORMULA FOR  
REAL QUADRATIC NUMBER FIELDS

DAVID M. BRADLEY, ALI E. ÖZLÜK AND C. SNYDER

For an even Dirichlet character  $\psi$ , we obtain a formula for  $L(1, \psi)$  in terms of a sum of Dirichlet  $L$ -series evaluated at  $s = 2$  and  $s = 3$  and a rapidly convergent numerical series involving the central binomial coefficients. We then derive a class number formula for real quadratic number fields by taking  $L(s, \psi)$  to be the quadratic  $L$ -series associated with these fields.

1. INTRODUCTION

In [1], acceleration formulæ are derived for Catalan's constant  $L(2, \chi_4)$ . (Here  $\chi_4$  is the non-principal Dirichlet character of modulus 4.) In some of these formulæ  $L(2, \chi_4)$  is given as the sum of two terms: one involving a rapidly convergent series and the other involving the natural logarithm of a unit in the ring of integers of a finite Abelian field extension of the rational number field  $\mathbb{Q}$ . The existence of the logarithmic terms suggested to the authors that these terms should somehow be related to the values of Dirichlet  $L$ -series at the argument  $s = 1$ . This leads to the general question of whether or not there exist relations between the value of  $L$ -series at  $s = 1$  and values of  $L$ -series at integer arguments larger than 1.

The purpose of this note is to exhibit such a relation between values of  $L$ -series. For an even Dirichlet character  $\psi$ , we obtain a formula for  $L(1, \psi)$  in terms of a sum of Dirichlet series evaluated at  $s = 2$  and  $s = 3$  and a convergent numerical series involving powers of twice special values of the sine function divided by  $\binom{2n}{n} n^3$ . See Theorem 1 below for a precise statement. (It is perhaps interesting to notice that not much is known about number theoretic properties of the values of the  $L$ -series on the right-hand side of the formula given in this theorem.) We then deduce a class number formula for real quadratic number fields by letting  $\psi$  be the quadratic character associated with a real quadratic number field; see Corollary 1. This class number formula seems new to us and is perhaps an interesting curiosity.

---

Received 28th August, 2001

---

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9727/02 \$A2.00+0.00.

To derive our results, we employ a formula of Zucker [5] that expresses

$$(1) \quad \sum_{n=1}^{\infty} \frac{x^{2n}}{\binom{2n}{n} n^3}, \quad |x| \leq 2,$$

in terms of periodic zeta functions. Proposition 1 below shows how periodic zeta functions may be expressed in terms of Dirichlet  $L$ -series. Thus, we can rewrite (1) in terms of  $L$ -series values, thereby obtaining our result.

### 2. PRELIMINARIES

Let  $m$  be a positive integer. We denote the group of Dirichlet characters of modulus  $m$  by  $\widehat{U}_m$ . The Dirichlet  $L$ -series associated with  $\chi \in \widehat{U}_m$  is

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}, \quad \text{Re}(s) > 1.$$

Similarly, for real  $\beta$  we define the *periodic zeta function* (a special case of the Lerch transcendent) by

$$\Phi(s, \beta) = \sum_{n=1}^{\infty} \frac{e^{2\pi i \beta n}}{n^s}, \quad \text{Re}(s) > 1.$$

Let  $\zeta_m = e^{2\pi i/m}$ . Throughout, the sum over a complete set of residues modulo  $m$  is denoted by  $\sum_{a \bmod m}$  and the sum over the positive integer divisors of  $m$  is denoted by  $\sum_{d|m}$ .

Thus, Ramanujan's sum is

$$c_m(k) = \sum_{\substack{\nu \bmod m \\ (\nu, m)=1}} \zeta_m^{\nu k},$$

and likewise the Gaussian sum attached to  $\chi$  is

$$\tau(\chi) = \sum_{\nu \bmod m} \chi(\nu) \zeta_m^{\nu}.$$

Also,  $\bar{\chi}$  denotes the inverse—or equivalently, the complex conjugate—of the character  $\chi$ . Finally, as customary,  $\mu()$ ,  $\varphi()$ , and  $\zeta()$  denote the Möbius, Euler totient, and Riemann zeta functions, respectively.

Our immediate goal is to represent periodic zeta functions in terms of  $L$ -series. It turns out to be easier to do the reverse first. The following result is well known, so we omit the proof.

**LEMMA 1.** *Let  $m$  be a positive integer, let  $\chi$  be a Dirichlet character of modulus  $m$ , and let  $L(s, \chi)$  be the associated Dirichlet  $L$ -series. Then*

$$L(s, \chi) = \frac{1}{m} \sum_{a \bmod m} \chi(a) \sum_{b \bmod m} \zeta_m^{-ab} \Phi(s, b/m), \quad \text{Re}(s) > 1.$$

**LEMMA 2.** *Let  $a$  and  $m$  be positive integers. Then*

$$\frac{1}{\varphi(m)} \sum_{\chi \in \hat{U}_m} \chi(a)\tau(\bar{\chi})L(s, \chi) = \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) c_m(a - b), \quad \text{Re}(s) > 1.$$

**PROOF:** First recall that

$$\sum_{\chi \in \hat{U}_m} \bar{\chi}(c)\chi(a) = \begin{cases} \varphi(m) & \text{if } (ac, m) = 1 \text{ and } a \equiv c \pmod m, \\ 0 & \text{otherwise.} \end{cases}$$

We claim that if  $(c, m) = 1$ , then

$$(2) \quad \frac{\varphi(m)}{m} \sum_{b \pmod m} \zeta_m^{-bc} \Phi(s, b/m) = \sum_{\chi \in \hat{U}_m} \chi(c)L(s, \chi).$$

By Lemma 1,

$$\begin{aligned} \sum_{\chi \in \hat{U}_m} \bar{\chi}(c) L(s, \chi) &= \frac{1}{m} \sum_{a \pmod m} \sum_{b \pmod m} \zeta_m^{-ab} \Phi(s, b/m) \sum_{\chi \in \hat{U}_m} \bar{\chi}(c)\chi(a) \\ &= \frac{\varphi(m)}{m} \sum_{b \pmod m} \zeta_m^{-bc} \Phi(s, b/m). \end{aligned}$$

On the other hand, if  $(c, m) > 1$ , then clearly

$$\sum_{\chi \in \hat{U}_m} \bar{\chi}(c) L(s, \chi) = 0.$$

We now multiply equation (2) by  $\zeta_m^{ac}$  with  $(a, m) = 1$ , and then sum over all  $c$  modulo  $m$ , obtaining

$$\begin{aligned} \frac{\varphi(m)}{m} \sum_{b \pmod m} \sum_{\substack{c \pmod m \\ (c,m)=1}} \zeta_m^{(a-b)c} \Phi(s, b/m) &= \sum_{c \pmod m} \zeta_m^{ac} \sum_{\chi \in \hat{U}_m} \bar{\chi}(c)L(s, \chi) \\ &= \sum_{c \pmod m} \zeta_m^{ac} \sum_{\chi \in \hat{U}_m} \bar{\chi}(c)L(s, \chi) \\ &= \sum_{\chi \in \hat{U}_m} \sum_{c \pmod m} \bar{\chi}(c)\zeta_m^{ac} L(s, \chi) \\ &= \sum_{\chi \in \hat{U}_m} \chi(a)\tau(\bar{\chi})L(s, \chi). \end{aligned}$$

Rewriting this latter equation in terms of Ramanujan sums completes the proof. □

We now state the main proposition of this section.

**PROPOSITION 1.** *Let  $a$  and  $m$  be coprime positive integers. Then*

$$m^s \Phi(s, a/m) = \sum_{d|m} \frac{d^s}{\varphi(d)} \sum_{\chi \in \hat{U}_d} \chi(a)\tau(\bar{\chi}) L(s, \chi), \quad \text{Re}(s) > 1.$$

Before proving Proposition 1, we state and prove two lemmata which are used in the proof of Proposition 1.

**LEMMA 3.** *Let  $f : \mathbb{Z} \rightarrow \mathbb{C}$  be multiplicative and such that for all positive integers  $m$ ,*

$$F(m) := \sum_{d|m} \mu^2(d)f(d)$$

*is non-zero. Furthermore, let*

$$g(m) := \sum_{d|m} \frac{\mu(d)}{F(d)}.$$

*Then for all positive integers  $k$  and  $m$  such that  $k$  divides  $m$ ,*

$$\sum_{\substack{d|m \\ k|d}} \mu^2(d)f(d) = F(m)\mu^2(k)g(k).$$

*In particular,*

$$\sum_{\substack{d|m \\ k|d}} \frac{\mu^2(d)}{\varphi(d)} = \frac{m}{\varphi(m)} \frac{\mu^2(k)}{k}.$$

**PROOF:** First, let us define  $F(x) = f(x) = 0$  if  $x$  is not an integer. Next, observe that  $F(p^a) = F(p)$  for all positive primes  $p$  and positive integers  $a$ . We may write  $m$  as  $\prod_p p^{a_p}$  and  $k$  as  $\prod_p p^{b_p}$  where  $p$  ranges over all positive primes and  $a_p$  and  $b_p$  are non-negative integers with  $b_p \leq a_p$ . Since  $F$  is multiplicative, we have

$$\begin{aligned} \sum_{\substack{d|m \\ k|d}} \mu^2(d)f(d) &= \prod_{p|m} \left( \sum_{\nu_p=b_p}^{a_p} \mu^2(p^{\nu_p})f(p^{\nu_p}) \right) \\ &= \prod_{p|m} (F(p^{a_p}) - F(p^{b_p-1})) \\ &= \prod_{p|m} F(p^{a_p}) \left( 1 - \frac{F(p^{b_p-1})}{F(p^{a_p})} \right) \\ (3) \qquad &= F(m) \prod_{p|m} \left( 1 - \frac{F(p^{b_p-1})}{F(p^{a_p})} \right). \end{aligned}$$

Notice that the final product in (3) vanishes if any  $b_p \geq 2$ , for then  $F(p^{b_p-1}) = F(p) = F(p^{a_p})$ . Hence if  $k$  is not square-free, then the lemma is trivially true as both sides are equal to 0. Therefore, we may assume henceforth that  $k$  is square-free. Now if  $b_p = 0$ , then  $1 - F(p^{b_p-1})/F(p^{a_p}) = 1$ , and thus (under the assumption that  $k$  is square-free), we

may restrict the final product in (3) to primes  $p$  for which  $b_p = 1$ . This yields

$$\begin{aligned} \sum_{\substack{d|m \\ k|d}} \mu^2(d)f(d) &= F(m) \prod_{p|k} \left(1 - \frac{1}{F(p^{a_p})}\right) = F(m) \prod_{p|k} \left(1 - \frac{1}{F(p)}\right) \\ &= F(m) \sum_{d|k} \frac{\mu(d)}{F(d)} \\ &= F(m)g(k). \end{aligned}$$

Thus, in general, we have

$$\sum_{\substack{d|m \\ k|d}} \mu^2(d)f(d) = F(m)\mu^2(k)g(k).$$

The special case is obtained by taking  $F(m) = m/\varphi(m)$ , so that if  $k$  is square-free, then

$$g(k) = \sum_{d|k} \frac{\mu(d)\varphi(d)}{d} = \prod_{p|k} \left(1 - \frac{\varphi(p)}{p}\right) = \prod_{p|k} \frac{1}{p} = \frac{1}{k}.$$

This completes the proof of Lemma 3. □

**LEMMA 4.** *Let  $m$  be a positive integer and let  $\beta$  be any real number. Then*

$$\sum_{\substack{n=1 \\ (n,m)=1}}^{\infty} \frac{e^{2\pi i\beta n}}{n^s} = \sum_{d|m} \frac{\mu(d)}{d^s} \Phi(s, \beta d).$$

**PROOF:** Let  $x$  be any complex number with  $|x| \leq 1$ . Then

$$\begin{aligned} \sum_{\substack{n=1 \\ (n,m)=1}}^{\infty} \frac{x^n}{n^s} &= \sum_{n=1}^{\infty} \frac{x^n}{n^s} \sum_{d|(n,m)} \mu(d) = \sum_{n=1}^{\infty} \frac{x^n}{n^s} \sum_{\substack{d|n \\ d|m}} \mu(d) = \sum_{d|m} \mu(d) \sum_{k=1}^{\infty} \frac{x^{kd}}{(kd)^s} \\ &= \sum_{d|m} \frac{\mu(d)}{d^s} \sum_{k=1}^{\infty} \frac{x^{kd}}{k^s}. \end{aligned}$$

Replacing  $x$  by  $e^{2\pi i\beta}$  completes the proof. □

**PROOF OF PROPOSITION 1:** First, recall (see for example [4, p. 238]) that Ramanujan’s sum has the explicit representation

$$c_m(k) = \varphi(m) \frac{\mu(m/(m, k))}{\varphi(m/(m, k))}.$$

Hence, we have

$$\begin{aligned}
 \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) c_m(a - b) &= \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) \varphi(m) \frac{\mu(m/(m, a - b))}{\varphi(m/(m, a - b))} \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \sum_{\substack{b \pmod m \\ (a-b, m) = m/d}} \Phi(s, b/m) \frac{\mu(d)}{\varphi(d)} \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{\substack{\nu \pmod d \\ (\nu, d) = 1}} \Phi\left(s, \frac{a + m\nu/d}{m}\right) \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{n=1}^{\infty} n^{-s} \sum_{\substack{\nu \pmod d \\ (\nu, d) = 1}} \zeta_m^{(a+m\nu/d)n} \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{n=1}^{\infty} n^{-s} \zeta_m^{an} c_d(n) \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{n=1}^{\infty} n^{-s} \zeta_m^{an} \varphi(d) \frac{\mu(d/(n, d))}{\varphi(d/(n, d))} \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \mu(d) \sum_{f|d} \frac{\mu(f)}{\varphi(f)} \sum_{\substack{n=1 \\ (n, f) = 1}}^{\infty} (nd/f)^{-s} \zeta_{fm/d}^{an} \\
 (4) \qquad &= \frac{\varphi(m)}{m} \sum_{d|m} \mu(d) \sum_{f|d} \frac{\mu(f)}{\varphi(f)} \left(\frac{d}{f}\right)^{-s} \sum_{\substack{n=1 \\ (n, f) = 1}}^{\infty} n^{-s} \zeta_{fm/d}^{an}.
 \end{aligned}$$

By Lemma 4 the final expression in (4) can be rewritten as

$$(5) \qquad \frac{\varphi(m)}{m} \sum_{d|m} \mu(d) \sum_{f|d} \frac{\mu(f)}{\varphi(f)} \left(\frac{d}{f}\right)^{-s} \sum_{\delta|f} \delta^{-s} \mu(\delta) \Phi\left(s, \frac{ad\delta}{fm}\right).$$

Now transform (5) by changing the variable  $f$  to  $d/f$ , then letting  $k = f\delta$  (noticing that the only non-zero terms occur when  $d$  is square-free), then observing that  $\sum_{f|k} \varphi(f) = k$ , and finally replacing  $d$  by  $kd$ . Thus, from (4) and (5),

$$\begin{aligned}
 \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) c_m(a - b) &= \frac{\varphi(m)}{m} \sum_{d|m} \mu(d) \sum_{f|d} \frac{\mu(d/f)}{\varphi(d/f)} f^{-s} \sum_{\delta|(d/f)} \delta^{-s} \mu(\delta) \Phi(s, af\delta/m) \\
 &= \frac{\varphi(m)}{m} \sum_{d|m} \sum_{k|d} \frac{\mu^2(d)}{\varphi(d)} k^{-s} \mu(k) \Phi(s, ak/m) \sum_{f|k} \varphi(f) \\
 (6) \qquad &= \frac{\varphi(m)}{m} \sum_{k|m} k^{1-s} \mu(k) \Phi(s, ak/m) \sum_{\substack{d|m \\ k|d}} \frac{\mu^2(d)}{\varphi(d)}.
 \end{aligned}$$

By applying Lemma 3 to (6) and then replacing  $m/k$  by  $d$ , we find that

$$\begin{aligned}
 \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) c_m(a - b) &= \sum_{k|m} k^{-s} \mu(k) \Phi(s, ak/m) \\
 (7) \qquad \qquad \qquad &= \frac{1}{m^s} \sum_{d|m} d^s \mu(m/d) \Phi(s, a/d).
 \end{aligned}$$

Hence by (7) and Lemma 2, we see that

$$\begin{aligned}
 \frac{1}{m^s} \sum_{d|m} d^s \mu(m/d) \Phi(s, a/d) &= \frac{1}{m} \sum_{b \pmod m} \Phi(s, b/m) c_m(a - b) \\
 &= \frac{1}{\varphi(m)} \sum_{\chi \in \bar{U}_m} \chi(a) \tau(\bar{\chi}) L(s, \chi).
 \end{aligned}$$

An application of Möbius inversion now completes the proof. □

### 3. MAIN RESULTS

We are now in a position to derive our class number formula. To this end, for  $|x| \leq 2$  and  $2 \leq k \in \mathbb{Z}$ , put

$$s(k, x) := \sum_{n=1}^{\infty} \frac{x^{2n}}{\binom{2n}{n} n^k}.$$

Let  $0 < \theta < \pi$  and  $x = 2 \sin \theta/2$ . Then [3, p. 61 (2)]  $2s(2, x) = \theta^2$  and by formula (2.7) of [5],

$$\begin{aligned}
 \theta^2 \log(2 \sin \theta/2) &= 2\zeta(3) + \sum_{n=1}^{\infty} \frac{(2 \sin \theta/2)^{2n}}{\binom{2n}{n} n^3} \\
 (8) \qquad \qquad \qquad &\qquad \qquad -2\theta \operatorname{Im} \Phi(2, \theta/2\pi) - 2\operatorname{Re} \Phi(3, \theta/2\pi),
 \end{aligned}$$

where  $\operatorname{Re}$  and  $\operatorname{Im}$  denote the real and imaginary parts of a complex number, respectively. Now substitute  $\theta = 2\pi a/m$  with  $(a, m) = 1$  and  $0 < a < m/2$  in (8) to obtain

$$\begin{aligned}
 \log \left( 2 \sin \frac{\pi a}{m} \right) &= \frac{m^2}{2\pi^2 a^2} \zeta(3) + \frac{m^2}{4\pi^2 a^2} \sum_{n=1}^{\infty} \frac{(2 \sin \pi a/m)^{2n}}{\binom{2n}{n} n^3} \\
 (9) \qquad \qquad \qquad &\qquad \qquad - \frac{m}{\pi a} \operatorname{Im} \Phi \left( 2, \frac{a}{m} \right) - \frac{m^2}{2\pi^2 a^2} \operatorname{Re} \Phi \left( 3, \frac{a}{m} \right).
 \end{aligned}$$

In our main result, character sums of consecutive integer powers arise, and it is convenient to fix some notation for these.

DEFINITION 1. Let  $m$  be a positive integer. If  $\chi$  is a Dirichlet character of modulus  $m$  and  $j$  is any integer, put

$$(10) \quad \mathcal{B}_j(\chi) := \sum_{0 < a < m/2} a^j \chi(a).$$

We now state and prove our main result.

THEOREM 1. Let  $m$  be a positive integer, let  $\psi$  be an even primitive character of modulus  $m$ , and let  $\mathcal{B}_j$  be as in (10). Then

$$\begin{aligned} L(1, \psi) &= \frac{2\tau(\psi)}{\pi i m^2} \sum_{d|m} \frac{d^2}{\varphi(d)} \sum_{\substack{\chi \in \widehat{U}_d \\ \chi \text{ odd}}} \mathcal{B}_{-1}(\chi\bar{\psi})\tau(\bar{\chi})L(2, \chi) \\ &\quad + \frac{\tau(\psi)}{\pi^2 m^2} \sum_{d|m} \frac{d^3}{\varphi(d)} \sum_{\substack{\chi \in \widehat{U}_d \\ \chi \text{ even}}} \mathcal{B}_{-2}(\chi\bar{\psi})\tau(\bar{\chi})L(3, \chi) - \frac{m\tau(\psi)}{\pi^2} \mathcal{B}_{-2}(\bar{\psi})\zeta(3) \\ &\quad - \frac{m\tau(\psi)}{2\pi^2} \sum_{n=1}^{\infty} \frac{1}{\binom{2n}{n}n^3} \sum_{0 < a < m/2} \frac{\bar{\psi}(a)}{a^2} \left(2 \sin \frac{\pi a}{m}\right)^{2n}. \end{aligned}$$

PROOF: We start with (9) and write  $\text{Im } \Phi(2, a/m)$  and  $\text{Re } \Phi(3, a/m)$  in terms of  $L$ -series via Proposition 1. First observe that

$$\begin{aligned} \text{Im} \left( \sum_{\chi \bmod d} \chi(a)\tau(\bar{\chi})L(2, \chi) \right) &= \frac{1}{2i} \sum_{\chi \in \widehat{U}_d} \left( \chi(a)\tau(\bar{\chi})L(2, \chi) - \overline{\chi(a)\tau(\bar{\chi})L(2, \chi)} \right) \\ &= \frac{1}{2i} \sum_{\chi \in \widehat{U}_d} \left( \chi(a)\tau(\bar{\chi})L(2, \chi) - \chi(-1)\bar{\chi}(a)\tau(\chi)L(2, \bar{\chi}) \right), \end{aligned}$$

since  $\overline{\tau(\bar{\chi})} = \chi(-1)\tau(\bar{\chi})$ . Now split the sum over the two terms and in the second sum replace  $\chi$  by  $\bar{\chi}$ . The even characters cancel and we obtain

$$\text{Im} \left( \sum_{\chi \in \widehat{U}_d} \chi(a)\tau(\bar{\chi})L(2, \chi) \right) = \frac{1}{i} \sum_{\substack{\chi \in \widehat{U}_d \\ \chi(-1)=-1}} \chi(a)\tau(\bar{\chi})L(2, \chi).$$

Similarly, we see that

$$\text{Re} \left( \sum_{\chi \in \widehat{U}_d} \chi(a)\tau(\bar{\chi})L(3, \chi) \right) = \sum_{\substack{\chi \in \widehat{U}_d \\ \chi(-1)=1}} \chi(a)\tau(\bar{\chi})L(3, \chi).$$

Thus by (9) and Proposition 1,

$$\begin{aligned}
 \log \left( 2 \sin \frac{\pi a}{m} \right) &= \frac{m^2}{2\pi^2 a^2} \zeta(3) + \frac{m^2}{4\pi^2 a^2} \sum_{n=1}^{\infty} \frac{(2 \sin \pi a/m)^{2n}}{\binom{2n}{n} n^3} \\
 &\quad - \frac{1}{\pi i m a} \sum_{d|m} \frac{d^2}{\varphi(d)} \sum_{\substack{\chi \in \hat{U}_d \\ \chi \text{ odd}}} \chi(a) \tau(\bar{\chi}) L(2, \chi) \\
 (11) \quad &\quad - \frac{1}{2\pi^2 m a^2} \sum_{d|m} \frac{d^3}{\varphi(d)} \sum_{\substack{\chi \in \hat{U}_d \\ \chi \text{ even}}} \chi(a) \tau(\bar{\chi}) L(3, \chi).
 \end{aligned}$$

Next, recall (see for example [2, p. 336]) that if  $\psi$  is an even primitive character of modulus  $m$ , then

$$\begin{aligned}
 L(1, \psi) &= -\frac{\tau(\psi)}{m} \sum_{a=1}^{m-1} \bar{\psi}(a) \log \left( 2 \sin \frac{\pi a}{m} \right) \\
 (12) \quad &= -\frac{2\tau(\psi)}{m} \sum_{0 < a < m/2} \bar{\psi}(a) \log \left( 2 \sin \frac{\pi a}{m} \right).
 \end{aligned}$$

Substituting (12) into (11) completes the proof. □

Let  $D$  be a (positive fundamental) discriminant of a real quadratic number field. Let  $h(D)$  denote its class number,  $\varepsilon = \varepsilon_D$  its fundamental unit  $> 1$ , and  $\chi_D = (D/\cdot)$ , the Kronecker symbol, that is, the Dirichlet character associated with the quadratic field of discriminant  $D$ . Then by Dirichlet (see for example [2, p. 343]), we know that

$$2h(D) \log \varepsilon_D = \sqrt{D} L(1, \chi_D).$$

Hence by Theorem 1, using the fact that  $\tau(\chi_D) = \sqrt{D}$ , we obtain the following class number formula.

**COROLLARY 1.** Class Number Formula

$$\begin{aligned}
 h(D) \log \varepsilon_D &= \frac{1}{\pi D i} \sum_{d|D} \frac{d^2}{\varphi(d)} \sum_{\substack{\chi \in \hat{U}_d \\ \chi \text{ odd}}} \mathcal{B}_{-1}(\chi \chi_D) \tau(\bar{\chi}) L(2, \chi) \\
 &\quad + \frac{1}{2\pi^2 D} \sum_{d|D} \frac{d^3}{\varphi(d)} \sum_{\substack{\chi \in \hat{U}_d \\ \chi \text{ even}}} \mathcal{B}_{-2}(\chi \chi_D) \tau(\bar{\chi}) L(3, \chi) \\
 &\quad - \frac{D^2}{2\pi^2} \mathcal{B}_{-2}(\chi_D) \zeta(3) \\
 &\quad - \frac{D^2}{4\pi^2} \sum_{n=1}^{\infty} \frac{1}{\binom{2n}{n} n^3} \sum_{0 < a < D/2} \frac{\chi_D(a)}{a^2} \left( 2 \sin \frac{\pi a}{D} \right)^{2n}.
 \end{aligned}$$

3.1. A COMPUTATION. As an amusing conclusion, we now show how to use our class number formula to compute  $h(5)$ , the class number of the quadratic field  $\mathbb{Q}(\sqrt{5})$ . Since the discriminant  $D = 5$ , the only relevant moduli of characters are  $m = 1$  and  $m = 5$ . For  $m = 1$ , the unique character is the even constant character 1. For  $m = 5$ , we have four characters determined by the homomorphisms from  $(\mathbb{Z}/5\mathbb{Z})^\times$  into  $\mathbb{C}^\times$ , namely  $\chi_\nu$  for  $\nu = 0, 1, 2, 3$  determined by  $\chi_\nu(2) = i^\nu$ . Notice that  $\overline{\chi_1} = \chi_3$  and that  $\chi_2 = (5/\cdot) = \chi_5$ , the Kronecker character modulo 5.

By Corollary 1, we have  $h(5) = (A + B + C + S)/\log \varepsilon_5$ , where

$$\begin{aligned}
 A &= \frac{5}{4\pi i} (\mathcal{B}_{-1}(\chi_3)\tau(\chi_3)L(2, \chi_1) + \mathcal{B}_{-1}(\chi_1)\tau(\chi_1)L(2, \chi_3)), \\
 B &= \frac{1}{10\pi^2} \left( \mathcal{B}_{-2}(\chi_5)\tau(1)\zeta(3) + \frac{125}{4} (\mathcal{B}_{-2}(\chi_5)\tau(\chi_0)L(3, \chi_0) + \mathcal{B}_{-2}(\chi_0)\tau(\chi_5)L(3, \chi_5)) \right) \\
 C &= -\frac{25}{2\pi^2} \mathcal{B}_{-2}(\chi_5)\zeta(3) \\
 S &= -\frac{25}{4\pi^2} \sum_{n=1}^{\infty} \frac{1}{\binom{2n}{n} n^3} \left( \chi_5(1)(2 \sin \pi/5)^{2n} + \frac{1}{4} \chi_5(2)(2 \sin 2\pi/5)^{2n} \right),
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{B}_{-1}(\chi_1) &= \chi_1(1) + \frac{1}{2}\chi_1(2) = 1 + \frac{1}{2}i \\
 \mathcal{B}_{-1}(\chi_3) &= 1 - \frac{1}{2}i \\
 \mathcal{B}_{-2}(\chi_0) &= 1 + \frac{1}{4} = \frac{5}{4} \\
 \mathcal{B}_{-2}(\chi_5) &= 1 - \frac{1}{4} = \frac{3}{4} \\
 \tau(1) &= 1 \\
 \tau(\chi_0) &= -1 \\
 \tau(\chi_5) &= \sqrt{5} \\
 \tau(\chi_1) &= \zeta_5 + i\zeta_5^2 - i\zeta_5^3 - \zeta_5^4 = \left( i + \frac{1 - \sqrt{5}}{2} \right) \sqrt{\frac{5 + \sqrt{5}}{2}} \\
 \tau(\chi_3) &= \zeta_5 - i\zeta_5^2 + i\zeta_5^3 - \zeta_5^4 = \left( i + \frac{\sqrt{5} - 1}{2} \right) \sqrt{\frac{5 + \sqrt{5}}{2}} \\
 L(3, \chi_0) &= (1 - 5^{-3}) \zeta(3) = \frac{124}{125} \zeta(3).
 \end{aligned}$$

In order to evaluate  $L(s, \chi_\nu)$  for  $\nu = 0, 1, 2, 3$  and  $s = 2, 3$  we write

$$L(s, \chi_\nu) = 5^{-s} \sum_{r=1}^4 \chi_\nu(r)\zeta(s, r/5),$$

where  $\zeta(s, a) = \sum_{n=0}^{\infty} (n+a)^{-s}$  is the Hurwitz zeta function. Hence to evaluate these  $L$ -series, it suffices to evaluate the Hurwitz zeta functions. The following table gives the appropriate approximations.

$r$	$\zeta(2, r/5)$	$\zeta(3, r/5)$
1	26.26737720...	125.73901805...
2	7.27535659...	16.1195643...
3	3.63620967...	4.98141576...
4	2.29947413...	2.21505785...

Thus, we find that

$$L(2, \chi_1) = 0.95871612\dots + (0.14556587\dots)i$$

$$L(2, \chi_3) = 0.95871612\dots - (0.14556587\dots)i$$

$$L(3, \chi_5) = 0.85482476\dots$$

Furthermore,  $\zeta(3) = 1.20205690\dots$ , so that

$$A = 1.24907310\dots$$

$$B = 0.48248793\dots$$

$$C = -1.14181713\dots$$

$$S = -0.10853146\dots$$

Finally, the fundamental unit

$$\varepsilon_5 = \frac{1 + \sqrt{5}}{2},$$

(which generally can be computed efficiently by continued fractions).

From all of this we obtain  $h(5) = 1.0000000\dots + (0.0000000\dots)i$ , whence  $h(5) = 1$ .

#### REFERENCES

- [1] D. Bradley, 'A class of series acceleration formulae for Catalan's constant', *Ramanujan J.* **3** (1999), 159–173.
- [2] A. Borevich and I. Shafarevich, *Number theory* (Academic Press, New York and London, 1966).
- [3] I.S. Gradshteyn and I.M. Ryzhik, *Table of integrals, series, and products*, (5th edition) (Academic Press, Boston, 1994).
- [4] G.H. Hardy and E.M. Wright, *An introduction to the theory of numbers*, (5th edition) (Clarendon Press, Oxford, 1979).
- [5] I.J. Zucker, 'On the series  $\sum_{k=1}^{\infty} \binom{2k}{k}^{-1} k^{-n}$  and related sums', *J. Number Theory* **20** (1985), 92–102.

Department of Mathematics and Statistics  
University of Maine  
Orono, Maine 04469  
United States of America  
e-mail: [bradley@gauss.umemat.maine.edu](mailto:bradley@gauss.umemat.maine.edu)  
[ozluk@gauss.umemat.maine.edu](mailto:ozluk@gauss.umemat.maine.edu)  
[snyder@gauss.umemat.maine.edu](mailto:snyder@gauss.umemat.maine.edu)