Longitudinal assessment of food insecurity status on the gut microbiome and metabolome of first year college students

A.E. Mohr^{1,2}, P. Jasbi³, I. van Woerden⁴, D.A. Bowes⁵, J. Chi^{1,6}, H. Gu^{1,6}, M. Bruening⁷*, C.M. Whisner^{1,2}*

¹College of Health Solutions, Arizona State University, Phoenix, AZ, USA

²Center for Health Through Microbiomes, Biodesign Institute, Arizona State University, Tempe, AZ, USA

³Systems Precision Engineering and Advanced Research (SPEAR), Theriome Inc., Phoenix, AZ, USA

⁴Department of Community and Public Health, Idaho State University, Pocatello, ID, USA

⁵Department of Environmental Health Sciences, Arnold School of Public Health, University of South Carolina, 921 Assembly Street, Columbia, SC 29208, USA.

⁶Center for Translational Science, Florida International University, Port St. Lucie, FL, USA

⁷Department of Nutritional Sciences, College of Health and Human Development, Pennsylvania State University, University Park, PA, USA

*Correspondence: Meg Bruening, PhD, MPH, RD, mmb203@psu.edu; Corrie M. Whisner,

PhD, cwhisner@asu.edu

Shortened Title: Food insecurity affects gut microbiome

Keywords: Food security; adolescence; gut microbiota; metabolome; multi-omics; systems biology; stability



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114525103668

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abbreviations:

- GM: Gut microbiome
- GI: Gastrointestinal
- FI: Food insecure
- FS: Food secure
- VAR: Variable food secure
- SPARC: Social impact of Physical Activity and nutRition in College
- ASU: Arizona State University
- BMI: Body mass index
- QIIME2: Quantitative Insights Into Microbial Ecology 2
- ASV: Amplicon sequence variants
- GTDB: Genome Taxonomy Database
- PICRUSt2: Phylogenetic Investigation of Communities by Reconstruction of Unobserved

States 2

- ACN: Acetonitrile
- MeOH: Methanol
- LC-MS: Liquid chromatography-mass spectrometry
- QC: Quality control
- HILIC: Hydrophilic interaction chromatography
- ESI: Electrospray ionization
- **RT**: Retention time
- MS: Exact mass
- HMDB: Human Microbiome Database
- ANOVA: Analysis of variance
- PERMDISP: Permutation test for homogeneity in multivariate dispersion
- LME: Linear-mixed effect
- PERMANOVA: Permutational multivariate analysis of variance
- GLM: General linear model
- PLS-DA: Partial least squares-discriminant analysis
- GFLASSO: Graph-guided fused least absolute shrinkage and selection operator
- RMSE: Root mean squared error

ER: Enrichment ratio

PA: Polyamine

Abstract

Food insecurity affects the health of college-aged individuals, but its impact on the gut microbiome (GM) over time is poorly understood. This study explored the association between food insecurity and the GM in 85 college students, identifying microbial taxa, metabolites, and pathways linked to food security status and examining GM stability and microbe-metabolite interactions. Longitudinal GM and metabolomic data were collected from first-year students over an academic year, encompassing periods of variable food security status. Participants were categorized into three groups: food insecure (FI, n=13), food secure (FS, n=44), and variable (VAR, n=28) status. GM composition varied significantly between FS classifications (Bray-Curtis dissimilarity, $P \le 0.005$). Stability analysis revealed correlations between stability scores and microbial features, pathways, and metabolites. Specific microbes (e.g., Bifidobacterium species, Faecalibacterium prausnitizii D, and Lachnospiraceae), pathways (energy and microbial turnover), and metabolites (cadaverine, N-acetylcadaverine, putrescine, testosterone sulfate, and creatine) associated with FI status were identified. Multi-omic integration revealed metabolic pathways influenced by differentially abundant microbial species and co-occurring fecal metabolites in food-insecure participants related to the microbial production of polyamines, detoxification, and energy metabolism. The transition from FS to FI showed no significant differences at specific taxonomic, functional, or metabolite levels. This study uncovers complex interactions between food security, GM composition, and metabolism. Significant differences were found in microbial community variability and metabolic pathways associated with food security status, but the transition from food security to insecurity disrupted the GM without clear taxonomic or functional distinctions, emphasizing the need for further research into these mechanisms.

Introduction

Food insecurity is an increasing public health problem that disproportionally affects vulnerable populations across the lifespan. Despite targeted efforts, food insecurity continues to be prevalent in the United States, occurring in 12.8% of households as of 2021 ⁽¹⁾. Defined as a socio-cultural construct tied to the lack of consistent access to healthy food ⁽²⁾, food insecurity is particularly problematic amongst college-aged individuals with prevalence levels several times higher than the national average ⁽³⁾. Food insecurity in this population is correlated with poor physical and mental health, dietary, and academic outcomes ^(3–7). During the formative period of emerging adulthood (the general age of college students), food insecurity has the potential to significantly disrupt an individual's health trajectory, leading to suboptimal health that may persist into later life ⁽⁸⁾.

Food insecurity shares multiple characteristics with dietary behaviors involving nutrient or food deprivation, such as caloric restriction, intermittent fasting, and disordered eating (9-11). However, it is important to recognize that the psychological mechanisms surrounding food insecurity, including anxiety and depression related to food access, are distinct from voluntary dietary practices (e.g., caloric restriction)⁽¹²⁾. Moreover, direct comparison of the aforementioned dietary behaviors with food insecurity for factors such as energy intake, nutrient profile, and other related health considerations may not be appropriate. Therefore, more research in this area is greatly warranted, particularly longitudinal assessments in tandem with systems biology approaches to unravel the complex impact of food insecurity on individual health. In this context, the gut microbiome (GM), comprising the diverse community of microbes inhabiting the gastrointestinal (GI) tract, has emerged as a valuable and elucidating factor, responsive to dietary pressures akin to food security status. For example, evidence supports that caloric restriction, intermittent fasting, and disordered eating influence the ecological landscape of the GM, modulating microbial competitiveness favoring particular taxa and metabolic pathways⁽¹³⁻¹⁶⁾. The metabolites synthesized or modified within this environment have also been reported to play a crucial role in corresponding health implications for the host ⁽¹⁷⁾. For instance, microbialgenerated metabolites have been suggested as a mediator between an altered GM and behavior associated with anorexia nervosa⁽¹⁵⁾. Consequently, the GM and metabolome represent essential complementary omic layers, poised to offer predictive biomarkers and co-occurrent signatures for capturing pathological variation over time.

Previously we established distinct microbial and metabolic signatures in food insecure individuals compared to food secure counterparts in a college-based population ⁽¹⁸⁾. Our findings suggested that compared to food secure students, the GM of food insecure students exhibited greater within-sample differences and estimated metabolic pathway activity, particularly related to hydrolysis reactions, energy substrate biosynthesis, and macronutrient metabolism. Additionally, we observed elevated metabolites (i.e., picolinic acid, phosphocreatine, 2pyrrolidinone) associated with energy transfer and gut-brain-axis communication with food insecurity ⁽¹⁸⁾. While this work was cross-sectional, these results indicate that the architecture of a food insecure GM may be primed to handle periods of deprivation and encode for the production/transformation of metabolic products relevant to gut-brain communication. However, the ecological stability of microbial communities under dietary pressure remains unclear, as within-subject longitudinal variation depends on factors like host lifestyle or sudden changes in nutritional status ⁽¹⁹⁾. Furthermore, it is important to understand how functional attributes and metabolic end-products of the GM may shift over time depending on variable or fixed food security status.

As a continuation of our research to uncover the influence of food insecurity status on the GM and its functional output ⁽¹⁸⁾, we assembled longitudinal GM and metabolomic data from first-year college students, capturing a timeframe that accurately aligns with the academic year and is known to induce variability in food security status ⁽³⁾. We categorized participants into three groups: those with a fixed status of either food insecure (FI) or food secure (FS), and those with a variable (VAR) status throughout the study period. Our objectives were to identify specific taxa, metabolites, and functional pathways in the GI tract associated with food security status, while also shedding light on GM stability and meaningful microbe-metabolite co-occurrence signatures that may influence host health. This investigation represents an important, early step in deepening our understanding of the underlying mechanisms linking food security status to GM dynamics and metabolic profiles, which may ultimately provide valuable insight to improve health outcomes in this vulnerable population.

Methods

Participants and Study design

The devilWASTE study was a sub-study of the Social impact of Physical Activity and nutRition in College (SPARC) study, which sought to analyze relationships between lifestyle factors,

weight outcomes, and the social networks of first-year college students ⁽²⁰⁾. DevilWASTE participants were recruited from the SPARC cohort from six dorms across three different Arizona State University (ASU) campuses. Participants in devilSPARC were recruited from one dorm on each of the three campuses. Given participants lived on campus, all participants were required to have a meal plan. All participants had pre-paid for at least 8 meals per week. The exclusion criteria for devilWASTE included age less than 18 years, certain GI conditions such as malabsorptive diseases, history of eating disorders, antibiotic use 2-3 months prior to study visits, and conditions that affect the microbiome including HIV infection, diabetes, and high blood pressure. Inclusion criteria included living in a residence hall at ASU, English speaking, and participation in the SPARC study. Eligible participants provided written informed consent before enrollment. The devilWASTE study and the parent SPARC study were approved by the ASU Institutional Review Board.

Data Collection

Recruitment for devilWASTE took place during the academic year starting in August 2015 and data collection continued through May 2016. Stool samples were collected at up to three of the four SPARC study time points (beginning and end of fall and spring semesters). In the present analysis, only individuals who provided a stool sample at the start of the Fall semester (T1), either at the end of the Fall or start of the Spring semester (T2), and at the end of the Spring semester (T3) were retained to support this longitudinal analysis. At each time point, anthropometrics, physical activity, and diet information were collected. At baseline sociodemographic characteristics including age, sex, race, and ethnicity were collected. Participant Pell grant status for the Fall 2015 and Spring 2016 semesters was acquired from university records.

Anthropometrics were obtained by trained research staff using Seca 869 scales (Seca, USA) for weight (kg), Seca 217 stadiometers (Seca, USA) for height (cm), and flexible, tension spring-loaded Gulick measuring tapes for waist (cm) and hip (cm) circumference (Creative Health Products, USA). These measurements were taken at each of the four devilSPARC study visits and completed up to three times at each visit to ensure accuracy. The two measures that were within 0.5 kg and 0.5 cm for weight and height, respectively, were averaged for final measurements. If none of the first three measurements were within 0.5 units of each other, another set of up to three measurements were taken. Similarly, the two waist and hip measures

within 0.5 cm were averaged for final measures, and if none were within 0.5 cm another set of up to three measurements was obtained. Participant body mass indexes (BMI) were calculated and reported in kg/m^2 while waist-to-hip ratios were calculated as waist divided by hip.

Questionnaires were used to evaluate lifestyle behaviors. Physical activity was assessed with the Godin-Shephard Leisure-Time Physical Activity Questionnaire which categorizes and quantifies activity into vigorous, moderate, and light physical activity ⁽²¹⁾. Sedentary behavior was assessed with an additional question: "Yesterday, how much time did you spend in front of a screen (excluding time in class and being physically active)?" ⁽²²⁾. Participants selected a response from a range of zero to six hours. Self-reported dietary intake was reported using the National Cancer Institute Dietary Screener Questionnaire that assesses consumption frequency of key food items and food groups ⁽²³⁾. This tool does not estimate caloric intake but rather tracks consumption of food group categories such as fruits and vegetables, high-fat and processed foods, added sugar, dietary fiber, and whole grain intake. Weekly alcohol intake was assessed by the number of drinks consumed weekly by asking: "For each day of the week in the calendar below, indicate the number of alcoholic drinks typically consumed on that day (Only if yes to alcohol is selected)." The dropdown ranged from 1-15 drinks for all days of the week. For capturing depression ⁽²⁴⁾ and perceived stress ⁽²⁵⁾, students completed previously validated questionnaires with robust test-rest correlations (0.89 and 0.74, respectively) in a similar collegeaged population $^{(6)}$.

Food insecurity was measured using the United States Department of Agriculture 6-item food insecurity screener ⁽²⁶⁾. The time frame in the validated question was adapted and the framing of the question changed from "we" to "I", as has been done by others ⁽²⁷⁾. Of the 6 items administered, participants that provided an affirmative answer to two or more questions were categorized as food insecure in the past month. Participants were then distributed into three groups: those with a fixed status of FI or FS, and those with a VAR status throughout the study period.

Fecal sample collection and DNA extraction

Research staff delivered fecal sample collections kits to the residence halls of eligible participants. As previously noted, fecal samples were collected at up to three timepoints for each participant. Participants were asked to report any medication and supplement use within the last 3 months; if participants had taken any antibiotics, antifungals, or probiotics within the previous

3 months, a fecal sample was not obtained. Research staff picked up the fecal samples within 30 min of a participant's reported bowel movement and transported them to the laboratory where they were frozen at -80 °C until further processing. Frozen samples were thawed at 4 °C, and wet weight was recorded to the nearest 0.01 g after subtracting the weight of fecal collection materials. DNA was extracted from approximately 300 mg of feces, collected from the center of the sample, using a modified version of the manufacturer protocol (MoBio Power Soil DNA Isolation Kit #12888-100, MoBio, Carlsbad, CA). Per manufacturer recommendations, a heating step of 65 °C for 10 min was added to the protocol to reduce the influence of inhibitors commonly found in feces and increase DNA yield. DNA concentration and quality were quantified using the QIAxpert System (Qiagen, Germantown, MD) according to manufacturer instructions.

Fecal microbiome sequencing

High-throughput genomic sequencing of the 16S rRNA gene was performed at the Biodesign Institute at ASU in Tempe, Arizona using Illumina miSeq technology after ligating 515F and 806R primers and Illumina adapters via polymerase chain reaction. Negative controls were included and run with the study samples. A detailed report of methods to prepare and sequence DNA has been published ^(18,28).

Due to the complex nature of the devilWASTE study design, large number of participants, and low-quality sequencing for a few samples in initial runs, three sequencing runs were conducted. In cases where samples were sequenced multiple times, files were merged after performing quality control. This method has been supported by expert census from the Quantitative Insights Into Microbial Ecology 2 (QIIME2) software development team ⁽²⁹⁾, the bioinformatic pipeline used in the present analysis. Overall, the 16S rRNA sequencing produced a total of 22,628,375 reads across the three runs with a median of 46,238 (run 1), 33,395 (run 2), and 77,808 (run 3) reads per sample. Paired-end, demultiplexed data were imported and analyzed using QIIME 2 software version 2021.8. Upon examination of sequence quality plots, base pairs were trimmed at position 20 and truncated at position 240 and were run through DADA2 (each run separately to prevent error modeling) to remove low quality regions and construct a feature table using amplicon sequence variants (ASVs). Next feature tables and representative sequences from the three runs were merged using the functions 'feature-table merge' and 'feature-table merge-seqs', respectively. Parameters were set to 'sum' for '--p-overlap-method' to pool all

features from duplicated samples in one and summarize all identical features in that sample. Taxonomy was assigned by first constructing a classifier against the Genome Taxonomy Database (GTDB; r.202)⁽³⁰⁾ using a naive Bayes approach via the q2-feature-classifier. Refined taxonomic classification was produced by assembling bespoke taxonomic weights with q2clawback to achieve benchmarked species-level resolution ⁽³¹⁾. A phylogenic tree was then constructed using the fragment-insertion plugin. A rarefaction threshold was assessed at 8,000 and used to impute high-quality reads and normalize for uneven sequencing depth between samples ⁽³²⁾. A *phyloseq* (v1.38.0) object was created, and downstream analyses and visualizations were performed in R (v.4.1.2). Sequences were removed including mitochondrial and plant DNA. In addition, any remaining singleton ASVs across samples were removed. Alpha-diversity was calculated using observed ASVs and phylogenetic diversity metrics. Betadiversity was assessed using the Bray-Curtis dissimilarity metric. Estimated functional potential of the overall bacterial community was surveyed via the Phylogenetic Investigation of Communities by Reconstruction of Unobserved States 2 (PICRUSt2) algorithm (v2.4.2)⁽³³⁾. Pathway abundances were inferred based on structured pathway mappings of Enzyme Commission gene families to the MetaCyc database ⁽³⁴⁾.

Fecal Metabolomics

Acetonitrile (ACN), methanol (MeOH), ammonium acetate, and acetic acid, all liquid chromatography-mass spectrometry (LC-MS) grade, were purchased from Fisher Scientific (Pittsburgh, PA). Ammonium hydroxide was bought from Sigma-Aldrich (Saint Louis, MO). DI water was provided in-house by a Water Purification System from EMD Millipore (Billerica, MA). PBS was bought from GE Healthcare Life Sciences (Logan, UT). The standard metabolite compounds were purchased from Sigma-Aldrich (Saint Louis, MO) and Fisher Scientific (Pittsburgh, PA).

Briefly, each fecal sample (~20 mg) was homogenized in 200 μ L MeOH:PBS (4:1, v:v, containing 1,810.5 μ M ¹³C₃-lactate and 142 μ M ¹³C₅-glutamic Acid) in an Eppendorf tube using a Bullet Blender homogenizer (Next Advance, Averill Park, NY). Then 800 μ L MeOH:PBS (4:1, v:v, containing 1,810.5 μ M ¹³C₃-lactate and 142 μ M ¹³C₅-glutamic Acid) was added, and after vortexing for 10 s, the samples were stored at -20 °C for 30 min. The samples were then sonicated in an ice bath for 30 min. The samples were centrifuged at 14,000 RPM for 10 min (4 °C), and 800 μ L supernatant was transferred to a new Eppendorf tube. The samples were then

dried under vacuum using a CentriVap Concentrator (Labconco, Fort Scott, KS). Prior to MS analysis, the obtained residue was reconstituted in 150 µL 40% PBS/60% ACN. A quality control (QC) sample was pooled from all the study samples.

The untargeted LC-MS metabolomics method used here was modeled after that developed and used in a growing number of studies ^(28,35). Briefly, all LC-MS experiments were performed on a Thermo Vanquish UPLC-Exploris 240 Orbitrap MS instrument (Waltham, MA). Each sample was injected twice, 10 μ L for analysis using negative ionization mode and 4 μ L for analysis using positive ionization mode. Both chromatographic separations were performed in hydrophilic interaction chromatography (HILIC) mode on a Waters XBridge BEH Amide column (150 x 2.1 mm, 2.5 μ m particle size, Waters Corporation, Milford, MA). The flow rate was 0.3 mL/min, auto-sampler temperature was kept at 4 °C, and the column compartment was set at 40 °C. The mobile phase was composed of Solvents A (10 mM ammonium acetate, 10 mM ammonium hydroxide in 95% ACN/5% H₂O). After the initial 1 min isocratic elution of 90% B, the percentage of Solvent B decreased to 40% at t=11 min. The composition of Solvent B maintained at 40% for 4 min (t=15 min), and then the percentage of B gradually went back to 90%, to prepare for the next injection. Using mass spectrometer equipped with an electrospray ionization (ESI) source, we will collect untargeted data from 70 to 1050 m/z.

To identify peaks from the MS spectra, we made extensive use of the in-house chemical standards (~600 aqueous metabolites), and in addition, we searched the resulting MS spectra against the Human Metabolome Database (HMDB) library, Lipidmap database, METLIN database, as well as commercial databases including mzCloud, Metabolika, and ChemSpider. The absolute intensity threshold for the MS data extraction was 1,000, and the mass accuracy limit was set to 5 ppm. Identifications and annotations used available data for retention time (RT), exact mass (MS), MS/MS fragmentation pattern, and isotopic pattern. We used the Thermo Compound Discoverer 3.3 software for aqueous metabolomics data processing. The untargeted data were processed by the software for peak picking, alignment, and normalization. To improve rigor, only the signals/peaks with CV < 20% across QC pools, and the signals showing up in >80% of all the samples were included for further analysis. Chemical taxonomy was obtained by querying metabolome features against the most recent version of the HMDB (v.5.0) ⁽³⁶⁾. A

Canberra distance matrix was computed on the metabolomics feature table, as previously described ⁽³⁷⁾.

Statistical Analysis

Anthropometric, behavioral, and dietary data from participant's was first assessed for normality using QQ-plots and Shapiro-Wilk's test. Differences in baseline values between the three FS statuses were then analyzed by analysis of variance (ANOVA) or Kruskal-Wallis tests, depending on normality. Potential differences in anthropometric, behavioral, and dietary data at each of the three time points between the three FS classifications were assessed by ANOVA or Kruskal-Wallis tests with post-hoc comparisons, where appropriate.

To ensure that the study had sufficient power to detect differences in gut microbiome composition across the food security groups, we conducted a power analysis using the Xmcupo test from the *HMP* package (v2.0.2.). The Xmcupo test applies a Dirichlet-Multinomial model, which is particularly suited for over-dispersed microbiome data. The Xmcupo test demonstrated sufficient statistical power to detect significant differences in microbial community composition between the groups. The analysis returned an Xmcupo statistic of 4207.375 (P < 0.001), indicating a highly significant difference in microbiome composition across food security statuses. This result confirms that, despite the unequal group sizes, the study design provides adequate power to detect meaningful differences in gut microbial diversity and composition between the food security groups.

For analysis of the GM, features and estimated functional pathways that were not present in at least 10% of samples were removed. First, a permutation test for homogeneity in multivariate dispersion (PERMDISP) was conducted using the 'betadisper' function in the *vegan* package (v2.6.2.) to compare dispersion across FS classification. Next, calculated intraindividual differences for the microbiome, estimated functional pathways, and fecal metabolome and alpha diversity metrics were tested for the effect of time, group, and their interaction using linear-mixed effect (LME) modeling using the *nLME* package (v3.1.153.). To assess feature stability, we computed Bray-Curtis dissimilarity (for species and estimated pathways) and Canberra distance (for the fecal metabolome) using the *ecodist* package (v2.0.9.) for each participant across the three sample-collection time points. We then calculated a stability score for each feature as 1 minus its Bray–Curtis (or) Canberra value (i.e., "1 – dissimilarity"). Because a smaller dissimilarity value indicates higher similarity across time points, this approach yields a

stability measure that ranges from 0 (least stable) to 1 (most stable). This metric has been previously described ^(19,38) but here serves to highlight how consistently a given feature's abundance remains within an individual over multiple time points. We then pooled all featurelevel observations (i.e., each species, pathway, or metabolite in each participant) to perform a single Spearman's rank correlation (ρ) comparing the distribution of stability scores to the distribution of log-transformed abundances within each data type (microbial species, pathways, or metabolites). Next, to determine whether the slope (relationship) between stability and abundance differed by FS status, we fitted separate linear regression models for each data type (estimates reflect the average across all features within that data type). In these models, individual feature observations were nested within each FS classification, and FS status was treated as an interaction term with the stability measure. Where a significant interaction was observed, we conducted post-hoc pairwise comparisons between FS categories to test for differences in the slope of stability vs. abundance. If significant interactions were detected, pairwise contrasts were used to detect significant differences in beta-coefficients between the FS classifications. Permutational multivariate analysis of variance (PERMANOVA) models were constructed for Bray-Curtis dissimilarity metrics (microbiome and estimated functional pathways) in the vegan package testing the effects of the individual (nested factor), FS status, time, and the interaction between these factors, controlling for sex and baseline BMI (permutation n = 999). Differential abundance analyses for microbial features at the phylum, family, and species level and estimated functional pathways were performed using the MaAsLin2 package with default parameters (v1.12.0)⁽³⁹⁾. To detect differences in features between the three FS statuses, we built linear mixed models with FS as the reference group that included time, and the covariates of sex and baseline BMI and the participant as a random factor.

To assess differences in metabolite abundances between the three FS statuses, fecal metabolome samples from all time points were pooled and a covariate adjusted general linear model (GLM), accounting for time, sex, and baseline BMI, was assembled with FS status used as the reference group. Significant features were then selected and used to construct a supervised partial least squares-discriminant analysis (PLS-DA) as a model predictive of FI and FS status. Pathway and enzyme topology and enrichment models were analyzed between groups using MetaboAnalyst software ⁽⁴⁰⁾. For pathway analysis, impact was calculated using a hypergeometric test, while significance was determined using a test of relative betweenness

centrality. All metabolomics data were log10-transformed, and Pareto scaled to approximate normality prior to analysis.

For multi-omic analysis, we integrated significant microbial taxa from the linear mixed models and the reliably detected and annotated fecal metabolomic data with graph-guided fused least absolute shrinkage and selection operator (GFLASSO) regression (R package: *GFLASSO*, v0.0.0.9000). Using this correlation-based network approach, significant microbial features were entered as the predictor variable and fecal metabolites as the response variable. Solution parsimony was determined by an unweighted (i.e., presence or absence of association by imposing a correlation threshold) network structure. For our purposes, we imposed a Spearman's rank correlation and fusion parameters were determined from the smallest root mean squared error (RMSE) estimate via cross-validation, accounting for interdependencies among microbial features. The tested parameters encompassed all combinations between λ and γ with values ranging from 0 to 1 (inclusive) in step increments of 0.1. GFLASSO coefficient matrices were constructed using a threshold coefficient of >0.02 to discern the strongest associative signals. The highest beta coefficients detected from GFLASSO models were further assessed by performing Spearman correlations of select microbial features with the response variables.

Where appropriate, statistical models accounted for baseline BMI and sex as covariates with participant as a random effect. A *P*-value of 0.05 was used to denote statistical significance and *P*-value adjustments (*Q*) were performed where appropriate using the Benjamini-Hochberg (BH) procedure. A *Q*-value < 0.10 was considered statistically significant. All statistical analyses were performed in the R environment (v4.1.2.).

Results

Participant Characteristics

In total, 85 participants from the previously described parent study ⁽²⁰⁾ provided a full stool sample at the requisite time points and were included in this analysis (**Fig. 1**). No significant differences between FS classification at baseline (beginning of the Fall semester) were detected for age, body weight, waist circumference, waist/hip ratio, or BMI (ANOVA/Kruskal-Wallis tests, $P \ge 0.085$); nor for self-reported lifestyle behaviors, including moderate to vigorous physical activity (MVPA), screen time, stress, sleep, and dietary factors (ANOVA/Kruskal-Wallis tests, $P \ge 0.079$; **Table 1**). There was a significant difference in depression and total alcohol

consumption (Kruskal-Wallis tests, $P \le 0.011$), with FI individuals displaying greater initial levels compared to FS individuals (pairwise comparison, $P \le 0.05$). As a proxy for economic status, we did not detect a significant difference between classifications for Pell grant status (Fisher's exact test, P = 0.981). No significant time or interaction effects were detected for any anthropometric or behavioral variable ($P \ge 0.085$; **Supplementary Table S1**).

Community structure and stability of gut microbes, function, and metabolome display variability by food security status

Covariate controlled, nested adonis analysis revealed significant individual variance over time between the three food security classifications (PERMANOVA, $R^2 = 0.194$, P = 0.001), with the individual effects of group ($R^2 = 0.017$, P = 0.001) and time ($R^2 = 0.005$, P = 0.001) explaining less overall variance. This was not the case for estimated functional pathways, which was not significant for any factor as above (P = 0.180) or the individual variance over time for the fecal metabolome (P = 0.304). However, the fecal metabolome was significant for the individual effects of group ($R^2 = 0.015$, P = 0.001) and time ($R^2 = 0.005$, P = 0.001).

There were no significant time or interaction effects for the surveyed alpha diversity metrics, observed ASVs and phylogenic diversity ($P \ge 0.122$; **Supplementary Fig. S1A-B**). Comparison of distributions of Bray-Curtis dissimilarity with a total of 3,570 ASVs across 255 samples revealed significant within-group differences by food security status (ANOVA, P =0.004), with the GM of FS participants showing greater overall variability compared to FI and VAR participants (pairwise comparison, $P \le 0.008$; **Fig. 2A**). Assessing Bray-Curtis intraindividual dissimilarity, time had a significant effect (P = 1.0e-04), with the overall GM community appearing to display greater dissimilarity between the first two timepoints and then becoming more similar to baseline by the third time point (**Fig. 2B**). In contrast, the pooled 334 estimated microbial functional pathways were not significant by any factor using Bray-Curtis dissimilarity (LME, $P \ge 0.468$; **Fig. 2C**), suggestive of the established conserved functional pathways compared to taxonomic membership in the GM ⁽⁴¹⁾. For the fecal metabolome, the reliably detected 554 metabolites (i.e., QC CV < 20% and relative abundance > 1,000 in 80% of samples) displayed no significant effects ($P \ge 0.173$; **Fig. 2D**) via assessment of the Canberra distance.

We next extended these results to explore the stability of individual microbial features and estimated functional pathways and fecal metabolites. Pooling feature-level values across

individuals within each data type, we noted moderate, large, and small correlations between stability scores and log(abundance) for microbial species (Spearman's $\rho = 0.422$, P = 2.2e-16; Fig. 2E), estimated pathways (Spearman's $\rho = 0.882$, P < 2.2e-16; Fig. 2F), and fecal metabolites (Spearman's $\rho = 0.211$, P < 2.2e-16; Fig. 2G), respectively. To determine if these correlations differed by FS status, we fit linear regression models in which each feature (e.g., each species) contributed a stability-versus-abundance data point, and FS status served as an interaction term. We found a significant interaction for microbial species (P = 0.034) such that the slope between stability and abundance differed significantly for FI and VAR over FS status $(Q \le 0.021)$. In other words, for these groups, an increase in species abundance was associated with either a steeper or shallower increase in stability than what we observed in the FS group. This suggests that food-insecure and variably food-secure students might rely on certain microbial species to maintain a more consistent community (higher stability) under changing dietary conditions – particularly for those species that are more abundant. A similar pattern emerged for estimated functional pathways (P = 0.004), where FI and FS showed a steeper stability-abundance relationship than VAR ($Q \le 0.041$). These findings indicate that the more abundant functional pathways remained more stable across time for FI and FS, whereas for VAR, abundance did not translate into stability to the same extent. In contrast, there was no significant difference for fecal metabolome stability by FS classification (P = 0.189). Because these analyses are done at the feature level, each reported slope represents an average effect across all species, pathways, or metabolites.

Differential feature shift and abundance in gut microbes, functional pathways, and metabolites by food security status

To better understand the significant group × time interaction we examined the differences between the three FS classifications by multivariate regression models, controlling for time, sex, and baseline BMI with FS as the reference, we found that the FI group had significantly greater species abundance of *Bifidobacterium samirii*, *Collinsella sp900760325*, *Bifidobacterium callitrichidarum*, *Turicibacter sp001543345*, *UBA9502 sp003669055* (unclassified *Lachnospiraceae*), *Bifidobacterium bifidum*, *Faecalibacterium prausnitzii D*, *Clostridium Q sp003024715*, *Phascolarctobacterium A sp900544885*, *UBA7182 sp003480725* (unclassified *Lachnospiraceae*), and *Gemella haemolysans* (log fold-change (logFC): 0.55-2.69, $Q \le 0.090$; **Fig. 3A**). Conversely, there was a significantly lower abundance in the family *Butyricicoccaceae*

(logFC: -1.163, Q = 0.097) and species *Paramuribaculum sp900551515*, *Akkermansia sp001580195*, *Bacteroides sp002491635*, and *Bacteroides sp902362375* in the FI group (logFC: -2.25-1.12, $Q \le 0.008$). For VAR participants, we observed greater abundance in *Dialisteraceae* at the family level (logFC: 1.23, Q = 0.097) and *GCA 900066495 sp900545985* (unclassified *Peptostreptococcaceae*), *Collinsella sp900760325*, *Dialister invisus*, *Anaerostipes hadrus*, *Phocaeicola vulgatus*, and *Faecalibacterium prausnitzii D* at the species level (logFC: 1.01-1.64, $Q \le 0.096$; **Fig. 3B**). There was a lower abundance of *Akkermansia sp001580195*, *ER4 sp000765235* (unclassified *Oscillospiraceae*), CAG-110 sp900544945 (unclassified *Oscillospiraceae*), CAG-83 sp900545585 (unclassified *Oscillospiraceae*), *Alistipes A indistinctus*, and *Phascolarctobacterium sp900544795* in the VAR group (logFC: -0.85-1.50, $Q \le$ 0.090).

At the functional pathway level, examining the differences between FS classification with FS as the reference, we found that the FI group had significantly greater abundance of L-lysine biosynthesis II, L-methionine biosynthesis I, superpathway of S-adenosyl-L-methionine biosynthesis, superpathway of L-lysine, L-threonine and L-methionine biosynthesis I, and superpathway of L-methionine biosynthesis (transsulfuration) (logFC: 0.30-0.91, $Q \le 0.043$; **Fig. 3C**). Conversely, there was a significantly lower abundance in Chondroitin sulfate degradation I (bacterial), Pyridoxal 5'-phosphate biosynthesis I, Phosphate biosynthesis and salvage, and superpathway of L-methionine biosynthesis (by sulfhydrylation) (logFC: -1.41-0.93, $Q \le 0.038$). For VAR participants, we observed greater pathway abundance in octane oxidation, myo-inositol degradation I, hexitol fermentation to lactate, formate, ethanol and acetate, and superpathway of hexitol degradation (bacteria) (logFC: 0.44-0.97, $Q \le 0.086$; **Fig. 3D**).

To assess potential individual feature shifts in the fecal metabolome, we pooled samples from all time points and assembled a covariate adjusted GLM, accounting for time, sex, and baseline BMI. With FS status as the reference group, we observed significant differences across classifications on 26 metabolites ($Q \le 0.089$). FI samples displayed increases in edetic acid, parathion, and N-acetylvaline, and decreases in n-arbamoylputrescine, L-carnitine, allysine, N6,N6,N6-trimethyl-L-lysine, thymine, carnosine, cysteinyl-aspartate, 4-aceotamido-2-amino-6nitroluene, testosterone sulfate, creatinine, L-palmitoylcarnitine, creatine, putrescine, nacetylcadaverine, cadaverine, and 5-hydroxylysine (**Fig. 3E**). It should be noted that these metabolites did change over time, though many were depleted (e.g., edetic acid, D-leucic acid,

thymine, and cysteinyl-aspartate) or enriched in abundance (e.g., cadaverine, Nacetylcadaverine, 5-hydroxylysine, and putrescine) as compared to FS participants. Of note, VAR shared trends in abundance change with the other classifications, though unlike either FI or FS metabolite shifts, there was a pattern for increased N-carbamoylputrescine, carnosine, 4acetamido-2-amino-6-nitrotoluene, creatinine, L-palmitoylcarnitine, creatine, and cadaverine (logFC > 0.5).

Next, a PLS-DA model was constructed using levels of significant between-group metabolites solely for FS and FI classifications because of the complexities involved in the status change for VAR. The model accounted for 37.9% of between-group variance (**Supplementary Fig. S2A**). Notably, PLS-DA regression coefficients showed that cadaverine, Nacetylcadaverine, putrescine, testosterone sulfate, and creatine accounted for the greatest influence of between group differences (model coefficients ≥ 60.84) and were all weighted towards FI status. Although the PLS-DA model displayed considerable accuracy ($R^2X = 0.789$), it exhibited low explanatory capacity ($R^2Y = 0.308$) and poor predictive capacity ($R^2Q = 0.172$). To assess the robustness of the PLS-DA model, 1000 permutations were performed to confirm the model was not overfit (**Supplementary Fig. S2B**; observed P < 0.001).

Comparing FI to FS, pathway enrichment analysis detected 16 significant pathways ($Q \le 0.091$; **Fig. 3F**). The most significant pathways were glutathione metabolism, lysine degradation, and fatty acid degradation ($Q \le 0.01$), whereas those with the greatest impact included histidine metabolism (I = 0.770), taurine and hypotaurine metabolism (I = 0.714), and arginine and proline metabolism (I = 0.538). At the main chemical class level, enrichment analysis queried against a reference library of 464 metabolite sets revealed enrichment of fatty amines, carboxylic acids, amines, steroid conjugates, arylsulfates, organic carbonic acids, organooxygen compounds, sulfonic acids, pyrimidines, fatty esters, and fatty acids and conjugates (FI/FS: $Q \le 0.098$; **Fig. 3G**). Compared to FS, the fecal metabolome of VAR participants revealed significant enrichment of organic thiophosphoric acids, steroid conjugates, purines, and butyrophenones ($Q \le 0.094$; data not shown). This was not the case for metabolic pathway enrichment analysis, where no differences were detected ($Q \ge 0.329$; data not shown).

Multi-omic integration suggests enriched FI microbes and co-occurrent fecal metabolites influence important metabolic processes in the gut

Seeking to uncover potential multi-omic signatures within the FI microbiome environment, we selected differentially abundant FI species (vs. FS classification) and the full set of metabolome features channeling these data sets in a regularized high-dimensional regression framework via GFLASSO. This model was selected due to its ability to handle large numbers of predictors and outcomes in comparison to the number of samples. To determine metabolites predictive of co-occurrence with driving features of the FI microbiome, we selected those microbes with a total regression coefficient greater than 0.2 to determine the most prominent signals, as previously described ⁽⁴²⁾. This rendered 11 species sets of 63 metabolites each, containing a total of approximately 11% of the entire reliably detected and annotated fecal metabolome (**Fig. 4**). Metabolites with positive beta-coefficients for significantly greater microbial species abundance with FI status were then selected and re-entered into a pathway enrichment analysis to determine how these multi-omic signatures might be influencing metabolism in the GM. Significantly enriched pathways included glutathione metabolism (ER: 4.05) for FI relative to FS ($Q \le 0.091$).

Focusing on the most pronounced co-occurrent patterns, we observed notable correlations between specific species and fecal metabolites through the use of Spearman correlations as a complementary approach, bolstering confidence in GFLASSO findings. Noteworthy correlations included: 1) *Bifidobacterium bifidum* with homoanserine (Spearman's $\rho = 0.17$, Q = 0.027), desaminotyrosine (Spearman's $\rho = 0.23$, Q = 1.36e-04), cadaverine (Spearman's $\rho = 0.32$, Q = 7.73e-06), triethanolamine (Spearman's $\rho = 0.35$, Q = 9.25e-07), gamma-L-Glutamyl-L-cysteine (Spearman's $\rho = -0.26$, Q = 2.21e-04), as well as putrescine (Spearman's $\rho = 0.38$, Q = 1.55e-07), and n-acetylcadaverine (Spearman's $\rho = 0.35$, Q = 9.62e-07); 2) *Gmella haemolysans* with glucose 1-phosphate (Spearman's $\rho = 0.33$, Q = 3.20e-06), cadaverine (Spearman's $\rho = 0.26$, Q = 2.72e-04), n-acetylcadaverine (Spearman's $\rho = 0.27$, Q = 2.05e-04), putrescine (Spearman's $\rho = 0.27$, Q = 1.86e-04), desaminotyrosine (Spearman's $\rho = 0.27$, Q = 1.57e-04), and glycyl lysine (Spearman's $\rho = -0.18$, Q = 0.001); 3) *Phascolarctobacterium A sp900544885* with n-acetylcadaverine (Spearman's $\rho = 0.31$, Q = 1.59e-05); and 4) *Faecalibacterium prausnitzii D* with nicotinic acid (Spearman's $\rho = 0.22$, Q = 0.003).

Exploration of the gut microbiome during transition from FS to FI status

Exploration of the GM during the transition from FS to FI status was undertaken to gain insights into the dynamic changes that occur under this shift. Leveraging the variability within the VAR group, consisting of 28 individuals, we focused on those who transitioned from initial FS to FI status during the study period, totaling 19 individuals (Fig. 5A) and a pre/post sample set of 38 observations for analysis. Assessing intra-individual dissimilarity using Bray-Curtis for the GM, estimated functional pathways, and Canberra distance for the fecal metabolome during the transition from FS to FI classification, we observed significant differences across all three omic layers, with the microbial community exhibiting the most pronounced disruption ($Q \le 0.001$; Fig. 5B). While no statistical significance was observed between different FI states for observed ASVs and phylogenetic diversity, we noted a trend toward significance for elevated richness and phylogenetic diversity with FI ($P \ge 0.369$; Fig. 5C-D). To discern significant features during the transition from FS to FI status, we conducted further analyses at the phylum, family, and species levels, as well as for estimated functional pathways and fecal metabolites. However, we did not detect any statistically significant differences ($Q \ge 0.144$). Notably, the observed variability in individual responses and the relatively small sample size within the transition subgroup may have contributed to these results (Fig. 5E).

Discussion

In the present study, we longitudinally characterized fecal GM composition, estimated function, and associated fecal metabolites of 85 dormitory housed college students with different food security status over the academic year. This approach afforded novel application to an understudied population and deeper phenotyping of the FI microbiome with the combined synergy of microbial sequencing and the fecal metabolome. Overall, our findings indicated a greater overall stability of the GM in college students experiencing persistent or intermittent FI when compared to their FS counterparts. Despite changes in GM taxonomy and fecal metabolite abundance over time, the estimated functional pathways within the GM remained relatively conserved across all groups. Moreover, we observed moderate to large correlations between microbial and estimated functional pathway stability scores and the abundance of individual microbial species. Notably, in a subset of students who transitioned from FS to FI status, we observed significantly greater community-level dissimilarity (Bray–Curtis or Canberra distance) across the GM, predicted functional pathways, and fecal metabolome ($Q \le 0.001$); however, no specific phylum-, family-, or species-level features or pathways reached statistical significance,

and alpha diversity showed only a nonsignificant trend toward higher richness and phylogenetic diversity in FI.

This study delved into the mechanisms through which FI may induce perturbations in GM composition and function. The observed GM stability in individuals experiencing FI may suggest either the conservation of GM function or that the stimuli (e.g., FI assessment within the past 30 days relative to fecal collection) were not potent enough to significantly disrupt GM activity. Previous studies have highlighted the functional redundancy of the human GM, emphasizing the highly conserved gene composition across a diverse taxonomic landscape ^(43,44). Regardless, taxonomic community analysis revealed significant individual variance over time between the three food security classifications at the species level, indicating that food security status influences the composition of gut microbes and serves as a key factor driving changes in the GM. These shifts in microbial composition were confirmed in multivariate regression models, which showed significant differences in species abundance between food security classifications. For instance, Bifidobacterium samirii, Collinsella sp900760325, Bifidobacterium callitrichidarum, and others were more abundant in the food-insecure group, while Akkermansia sp001580195 and other species were less abundant. Understanding these differences is not straightforward as microbes positively associated with FI status like Phocaeicola vulgatus are context dependent and may have beneficial (improvement of lipid metabolism ⁽⁴⁵⁾) or detrimental (involved in human infections ⁽⁴⁶⁾) influence on the host.

In contrast to microbial composition, estimated functional pathways did not exhibit significant differences between food security classifications in the PERMANOVA analysis. This suggests that, at a functional level, the GM may maintain stability despite changes in microbial species. Nonetheless, examining specific functional pathway trends revealed varying profiles across the food security classifications. For example, food-insecure individuals displayed increased pathways related to energy and microbial turnover, while food-secure individuals showed a broad increase in fatty acid anabolism. These trends are consistent with the presumed health state of the individuals in these different FS classifications, in which FI students may experience erratic or reduced nutrient intake – potentially driving up energy-harvesting processes – while FS students have consistent access to meals, supporting anabolic pathways (e.g., lipid synthesis). However, to pinpoint the exact onset and duration of these functional shifts in the GM, more frequent longitudinal sampling would be necessary. Denser sampling could capture

day-to-day or week-to-week changes in diet and microbial function, offering greater insight into how quickly the microbiome responds to fluctuations in food availability.

In assessment of the fecal metabolome, metabolites that accounted for the most significant differences between food security groups include cadaverine, N-acetylcadaverine, putrescine, testosterone sulfate, and creatine, establishing a prominent role in distinguishing food-insecure individuals. Pathway enrichment analysis identified specific metabolic pathways that were significantly enriched in food-insecure individuals compared to food-secure individuals. These pathways included glutathione metabolism, lysine degradation, and fatty acid degradation, suggesting alterations in key metabolic processes associated with food insecurity. Because fecal metabolites can originate from both microbial and host processes, these findings likely reflect microbially mediated metabolic shifts and potential host responses to inconsistent or nutrient-poor food intake. For instance, increases in glutathione-related pathways may indicate heightened oxidative stress handling in the gut environment due to poor nutritional quality and the consumption of energy-dense, nutrient-poor foods ⁽⁴⁷⁾. This stress could be due to the cyclical nature of food availability, leading to periods of overeating followed by scarcity. In relation, greater lysine and fatty acid degradation could reflect adaptations in microbial and/or host metabolism when carbohydrate or protein availability is limited. This could also be a compensatory mechanism to maintain energy homeostasis when dietary intake is insufficient. Finally, the degradation of fatty acids, or beta-oxidation, is a metabolic process that provides energy when carbohydrate availability is low. In FI individuals, increased fatty acid degradation may suggest reliance on fat stores for energy due to inconsistent access to food, particularly carbohydrates. Although we discuss possible physiological implications for humans experiencing FI (e.g., cyclical intake of nutrient-dense yet low-quality foods), our data do not definitively isolate whether these catabolic processes are predominantly driven by host versus microbial mechanisms. Future studies employing direct host biomarkers (e.g., blood metabolites) or shotgun metagenomic/metatranscriptomic approaches would help clarify the specific contributions of the gut microbiome relative to the host under conditions of food insecurity.

To uncover potential multi-omic signatures within the food-insecure microbiome, a highdimensional regression framework was applied, identifying specific metabolites associated with food-insecure microbial species. Correlation analyses confirmed co-occurring patterns between certain microbial species and fecal metabolites. Notable positive correlations included those

between Bifidobacterium bifidum and cadaverine, triethanolamine, putrescine, and nacetylcadaverine. Similarly, *Gmella haemolysans* was positively associated with cadaverine, putrescine, and n-acetylcadaverine, as well as Phascolarctobacterium A sp900544885 with nacetylcadaverine and Faecalibacterium prausnitzii D with nicotinic acid. The current understanding of these specific metabolites is largely underdeveloped with respect to their overall presence, physiological role, functional pathways, and effects in the human body. Despite the paucity of studies acknowledging exogenous (i.e., food sources) and endogenous (i.e., cell synthesis) sources of these polyamines (PA), putrescine or cadaverine are thought to be more greatly produced from microbial sources ⁽⁴⁸⁾; a connection that is still relatively uncharacterized. Very few studies to date have accomplished reporting linkages between metabolite production and GM composition as demonstrated in the present study ⁽⁴⁹⁾. For example, Kitada et al., discovered novel linkages between three commensal bacterial populations, including *Bifidobacterium spp.* and the enhanced production of putrescine ⁽⁵⁰⁾, a PA commonly found in the intestinal lumen within the human gut as well as in a variety of foods and food products, with higher concentrations typically found in fermented dairy products and animal-based processed foods ⁽⁵¹⁾. Interestingly, soluble and fermentable dietary fiber consumption has been shown to enhance the gut microbial production of PAs putrescine, cadaverine, and spermidine in rodents with gut microbes playing a key role (52-54). Previous research suggests the importance of PAs from dietary constituents for meeting the needs of the human body ⁽⁽⁵⁵⁾ which support intestinal DNA, RNA and protein synthesis, as well as cellular proliferation and differentiation ^(56,57). Pathway enrichment analyses further support this idea in that gut microbes are capable of decarboxylating lysine to cadaverine, and lysine degradation was identified as a key pathway in FI participants. Overall, these data suggest that microbially-derived PAs may serve as energy substrate ⁽⁵⁸⁾ in the gut during periods of FI or intermittent food access.

Based on the findings above, we can tangentially begin to make connections to how these interactions may be reflected within the human GM in response to food insecurity. For example, recent evidence reports in extreme FI conditions, such as migration/forced displacement, that certain coping strategies may develop in order to satisfy hunger or to ensure other members of the household are receiving adequate nutrition (e.g., children, elderly). One of these coping strategies includes consuming expired or contaminated food products, which may lead to the increased production, circulation, and excretion of PA from the human body due to increased

microbial activity ^(59,60). Finally, the positive correlation observed in this study between *F. prausnitzii D* and nicotinic acid (vitamin B3) is intriguing. While several microbial species obtain the vitamin B3 biosynthesis pathway, the role of vitamin B3 on the human GM is still considered relatively underdeveloped. Specifically investigating this relationship between *F. prausnitzii D* and vitamin B3, we can again turn to other evidence to help support this potential phenomenon. For example, studies have confirmed that the butyrate-producer *F. prausnitzii* increases with enhanced uptake of riboflavin (vitamin B2) ^(61,62). Vitamin B3 can be synthesized from the amino acid, tryptophan, where vitamin B2 serves as an important cofactor in relation to energy metabolism. While these two B vitamins share similar food sources, such as meat, dairy, eggs, and fortified foods, this positive correlation between vitamin B3 and *F. prausnitizii D* may reveal an underlying mechanism not yet well understood that warrants further investigation ⁽⁶³⁾ within the specific context of food insecurity and impacts on the human condition.

It is essential to recognize that FI extends beyond dietary factors and encompasses important social and psychological dimensions, such as psychological well-being and stress, which can influence the relationship between FI and dietary intake ⁽⁶⁴⁾. These factors may underpin the complex interplay between FI and GM structure and function ⁽¹⁸⁾. Our previous research has highlighted the influence of these psychological factors in the context of FI. For example, we reported FI students had elevations in the fecal metabolites, picolinic acid, 2-pyrrolidinone, and phosphocreatine ⁽¹⁸⁾. These metabolites are central to neurological health ⁽⁶⁵⁾, gut-axis signaling ⁽⁶⁶⁾, and energy metabolism ⁽⁶⁷⁾, respectively. Many of the microbial and metabolomic features detected in the present analysis seem to be more nutritionally orientated, though could have secondary influence in relation to the gut-brain axis. Future work will be required with expanded longitudinal sampling as mentioned previously.

Very few studies have explored associations between FI and GM composition. An impactful strength of the present work was the longitudinal nature of the data allowing for the study of persistent and intermittent periods of FI in relation to the GM and metabolome and estimated functional activities. To our knowledge, no studies have explored the longitudinal relationship between FI and GM structure in a vulnerable adolescent population. Although our sample size of 85 participants may be considered moderate for an observational study, the repeated measures and multi-omic data still provided meaningful insights into the complex interplay of social, biological, and physiological factors underlying FI. Lastly, stool samples

serve as a valuable source of objective biological data reflecting gut microbiome-host interactions, complementing self-reported dietary measures in understanding the holistic dietary and physiological context.

Despite the mentioned strengths, this study was not without limitations. Firstly, there was a sample size difference between FI and FS groups that occurred as a result of investigating a convenience sample. Although not balanced, these groups were representative of the student population under study ^(68,69) and cohorts from observational studies of food security among college students ^(3,18). Further, although 85 participants provided sufficient longitudinal data to detect notable differences, this is considered moderate for an observational study; subgroup analyses with fewer than 30 individuals inherently limit the detection of smaller effect sizes. Secondly, while dietary intake was assessed using the National Cancer Institute Dietary Screener Questionnaire, these data were not used as covariates because they did not significantly differ across the food security groups. Moreover, the screener does not capture the variety of foods consumed or potentially relevant eating behaviors (e.g., intermittent fasting), making it less sensitive in detecting nuanced dietary patterns that could affect the GM. Consequently, we could not fully capture differences in dietary diversity between groups. It is possible that FS participants had access to a wider variety of foods, contributing to the greater variability in their gut microbiome observed in this study. Thirdly, another limitation of this work was the lack of validated instruments for assessing FI in college students ⁽⁷⁰⁾, as the instrument used has only been validated in low-income adults ⁽⁷⁰⁾. Albeit, most studies of college students use a version of the measure used in the current study ⁽²⁷⁾. All of the students in this study had purchased a prepaid meal plan, which in theory minimizes food insecurity. However, our prior research has shown that even students who pre-paid for an unlimited meal plan report food insecurity ⁽⁵⁾. Potential explanations for this apparently contradictory finding include conflicts with students' schedules and dining hall hours, and student time management. Fourth, although our multi-omic design provides novel insights, we inferred microbial function using 16S rRNA gene sequencing rather than employing shotgun metagenomic approaches. Because reference databases for 16Sbased functional prediction are incomplete, these functional inferences could introduce biases and must be interpreted cautiously. Finally, while we characterized overall relationships between abundance and stability, we did not define a "core" or exceptionally stable set of taxa or

functions, as this would require threshold-specific analyses and multiple-comparisons with more frequent sampling that extend beyond the scope of the present study.

In summary, the results suggest that food security status is associated with significant shifts in the gut microbial composition, metabolite profiles, and metabolic pathways among college students. These findings underscore the complex interplay between food security, the GM, and host metabolism, highlighting the potential role of the GM in responding to dietary and environmental factors associated with food insecurity. Moving forward, future research should focus on addressing the limitations identified, such as employing more robust measures for assessing food security, dietary intake, and mental health symptoms associated with food insecurity. Additionally, diversifying the study population to include individuals not bound by dorm contracts with provided food could offer valuable insights into the broader spectrum of food security experiences among college students. Further investigation is needed to elucidate the mechanistic links between these observations and their implications for health and well-being in this population.

Acknowledgements:

The authors are grateful for the time participants generously volunteered for this project.

Data Availability:

Due to the vulnerable nature of adolescent and emerging adult populations, researchers did not seek participant consent to share study data. Therefore, participants of this study did not agree for their data to be shared publicly, and supporting data are not available.

Financial support:

This study received support from the NIH Common Fund through the Office of the Director and the Office of Behavioral and Social Sciences Research, with grant number [1DP5OD017910] awarded to M.B. as the Principal Investigator. Additionally, research presented in this publication was funded by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under Award Number T32DK137525 (A.E.M.). The views expressed in this paper are solely those of the authors and do not necessarily reflect the official perspectives of the National Institutes of Health.

Declaration of Interests:

C.M.W. serves on the scientific advisory boards for Avocado Nutrition Center's Avocado Nutrition Science Advisory and the California Walnut Commission. All other authors declare no conflicts of interest.

Authorship:

A.E.M., M.B., and C.W. formulated the research question and designed the analysis. P.J., I.v.W., H.G., M.B., and C.W. conducted research and/or provided essential materials. A.E.M. performed statistical analysis, interpreted results, and drafted the manuscript. P.J., D.B., and C.W. contributed to the statistical analysis and interpretation of results. A.E.M., P.J., I.v.W., D.B., H.G., M.B., and C.W. critically reviewed the manuscript. M.B. and C.W. secured funding and supervised the entire project. All authors were given the opportunity to read and contribute to the text and approved the final manuscript.

References

1. Department of Agriculture. Economic Research Service (2023) - Household food security in the United States in 2022. <u>https://doi.org/10.32747/2023.8134351.ers</u> (accessed Feburary 2024).

2. Coleman-Jensen A, Gregory C, & Singh A (2014) Household Food Security in the United States in 2013. *Ssrn Electron J*. doi:10.2139/ssrn.2504067.

3. Bruening M, Argo K, Payne-Sturges D, & Laska MN (2017) The Struggle Is Real: A Systematic Review of Food Insecurity on Postsecondary Education Campuses. *J Acad Nutr Diet* **117**, 1767–1791.

4. Larson N, Laska MN, & Neumark-Sztainer D (2020) Food Insecurity, Diet Quality, Home Food Availability, and Health Risk Behaviors Among Emerging Adults: Findings From the EAT 2010–2018 Study. *Am J Public Health* **110**, 1422–1428.

Woerden I van, Hruschka D, Vega-López S, Schaefer DR, Adams M, & Bruening M (2019)
 Food Insecure College Students and Objective Measurements of Their Unused Meal Plans.
 Nutrients 11, 904.

6. Bruening M, Brennhofer S, Woerden I van, Todd M, & Laska M (2016) Factors Related to the High Rates of Food Insecurity among Diverse, Urban College Freshmen. *J Acad Nutr Diet* **116**, 1450–1457.

7. Woerden I, Hruschka D, & Bruening M (2019) Food insecurity negatively impacts academic performance. *J Public Aff* **19**, e1864.

8. Gundersen C & Ziliak JP (2017) Food Insecurity And Health Outcomes. *Health Affair* **34**, 1830–1839.

9. Kendall A, Olson CM, & Frongillo EA (1996) Relationship of Hunger and Food Insecurity to Food Availability and Consumption. *J Am Diet Assoc* **96**, 1019–1024.

10. Hazzard VM, Loth KA, Hooper L, & Becker CB (2020) Food Insecurity and Eating Disorders: a Review of Emerging Evidence. *Curr Psychiat Rep* **22**, 74.

 Stinson EJ, Votruba SB, Venti C, Perez M, Krakoff J, & Gluck ME (2018) Food Insecurity is Associated with Maladaptive Eating Behaviors and Objectively Measured Overeating. *Obesity* 26, 1841–1848.

12. Nagata JM, Palar K, Gooding HC, Garber AK, Whittle HJ, Bibbins-Domingo K *et al.* (2019) Food Insecurity Is Associated With Poorer Mental Health and Sleep Outcomes in Young Adults. *J Adolesc Heal* **65**, 805–811.

13. Mohr AE, Jasbi P, Bowes DA, Dirks B, Whisner CM, Arciero KM *et al.* (2022) Exploratory analysis of one versus two-day intermittent fasting protocols on the gut microbiome and plasma metabolome in adults with overweight/obesity. *Frontiers Nutrition* **9**, 1036080.

14. Schwartzenberg RJ von, Bisanz JE, Lyalina S, Spanogiannopoulos P, Ang QY, Cai J *et al.*(2021) Caloric restriction disrupts the microbiota and colonization resistance. *Nature* 595, 272–277.

15. Fan Y, Støving RK, Ibraim SB, Hyötyläinen T, Thirion F, Arora T *et al.* (2023) The gut microbiota contributes to the pathogenesis of anorexia nervosa in humans and mice. *Nat Microbiol* **8**, 787–802.

16. Mohr AE, Sweazea KL, Bowes DA, Jasbi P, Whisner CM, Sears DD *et al.* (2024) Gut microbiome remodeling and metabolomic profile improves in response to protein pacing with intermittent fasting versus continuous caloric restriction. *Nat Commun* **15**, 4155.

17. Vila AV, Hu S, Andreu-Sánchez S, Collij V, Jansen BH, Augustijn HE *et al.* (2023) Faecal metabolome and its determinants in inflammatory bowel disease. *Gut* **72**, 1472–1485.

18. Mohr AE, Jasbi P, Wyst KBV, Woerden I van, Shi X, Gu H *et al.* (2022) Association of food insecurity on gut microbiome and metabolome profiles in a diverse college-based sample. *Sci Rep-uk* **12**, 14358.

19. Mehta RS, Abu-Ali GS, Drew DA, Lloyd-Price J, Subramanian A, Lochhead P *et al.* (2018) Stability of the human faecal microbiome in a cohort of adult men. *Nat Microbiol* **3**, 347–355.

20. Bruening M, Ohri-Vachaspati P, Brewis A, Laska M, Todd M, Hruschka D *et al.* (2016) Longitudinal social networks impacts on weight and weight-related behaviors assessed using mobile-based ecological momentary assessments: Study Protocols for the SPARC study. *Bmc Public Health* **16**, 901.

21. Sirard JR & Pate RR (2001) Physical Activity Assessment in Children and Adolescents. *Sports Med* **31**, 439–454.

22. Utter J, Neumark-Sztainer D, Jeffery R, & Story M (2003) Couch potatoes or French fries: Are sedentary behaviors associated with body mass index, physical activity, and dietary behaviors among adolescents? *J Am Diet Assoc* **103**, 1298–1305.

23. National Cancer Institute (2009) - Dietary Screener Questionnaire in the NHANES 2009–10. http://appliedresearch.cancer.gov/nhanes/dietscreen/dsq_english.pdf (accessed June 2022).

24. American College Health Association (2013) *American College Health Association-National College Health Assessment II*: Reference Group Report Spring 2013. Hanover, MD: American College Health Association.

25. Cohen S, Kamarck T, & Mermelstein R (1983) A global measure of perceived stress. *J Heal Soc Behav* 24, 385–96.

26. U.S. Household Food Security Survey Module: Three-Stage Design, With Screeners. <u>https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-us/survey-tools/#household</u> (accessed March 2023).

27. Nikolaus CJ, Ellison B, & Nickols-Richardson SM (2019) Are estimates of food insecurity among college students accurate? Comparison of assessment protocols. *Plos One* **14**, e0215161.

 28. Jasbi P, Mohr AE, Shi X, Mahmood T, Zhu Q, Bruening M *et al.* (2022) Microbiome and metabolome profiles of high screen time in a cohort of healthy college students. *Sci Rep-uk* 12, 3452. 29. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA *et al.* (2019) Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* **37**, 852–857.

30. Chaumeil P-A, Mussig AJ, Hugenholtz P, & Parks DH (2022) GTDB-Tk v2: memory friendly classification with the genome taxonomy database. *Bioinformatics* **38**, 5315–5316.

31. Kaehler BD, Bokulich NA, McDonald D, Knight R, Caporaso JG, & Huttley GA (2019) Species abundance information improves sequence taxonomy classification accuracy. *Nat Commun* **10**, 4643.

32. McKnight DT, Huerlimann R, Bower DS, Schwarzkopf L, Alford RA, & Zenger KR (2019) Methods for normalizing microbiome data: An ecological perspective. *Methods Ecol Evol* **10**, 389–400.

33. Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM *et al.* (2020) PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol* **38**, 685–688.

34. Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM *et al.* (2016) The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res* **44**, D471–D480.

35. Scieszka DP, Garland D, Hunter R, Herbert G, Lucas S, Jin Y *et al.* (2023) Multi-omic assessment shows dysregulation of pulmonary and systemic immunity to e-cigarette exposure. *Respir Res* **24**, 138.

36. Wishart DS, Guo A, Oler E, Wang F, Anjum A, Peters H *et al.* (2021) HMDB 5.0: the Human Metabolome Database for 2022. *Nucleic Acids Res* **50**, D622–D631.

37. Raghuvanshi R, Vasco K, Vázquez-Baeza Y, Jiang L, Morton JT, Li D *et al.* (2020) High-Resolution Longitudinal Dynamics of the Cystic Fibrosis Sputum Microbiome and Metabolome through Antibiotic Therapy. *Msystems* **5**, e00292-20.

38. Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL *et al.* (2013) The Long-Term Stability of the Human Gut Microbiota. *Science* **341**, 1237439.

39. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH *et al.* (2021) Multivariable association discovery in population-scale meta-omics studies. *Plos Comput Biol* **17**, e1009442.

40. Pang Z, Zhou G, Ewald J, Chang L, Hacariz O, Basu N *et al.* (2022) Using MetaboAnalyst 5.0 for LC–HRMS spectra processing, multi-omics integration and covariate adjustment of global metabolomics data. *Nat Protoc* **17**, 1735–1761.

41. Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT *et al.* (2012) Structure, Function and Diversity of the Healthy Human Microbiome. *Nature* **486**, 207–214.

42. Lima FA e, Li K, Wen W, Yan J, Nikoloski Z, Willmitzer L *et al.* (2018) Unraveling lipid metabolism in maize with time-resolved multi-omics data. *Plant J* **93**, 1102–1115.

43. Tian L, Wang X-W, Wu A-K, Fan Y, Friedman J, Dahlin A *et al.* (2020) Deciphering functional redundancy in the human microbiome. *Nat Commun* **11**, 6217.

44. Li L, Wang T, Ning Z, Zhang X, Butcher J, Serrana JM *et al.* (2023) Revealing proteomelevel functional redundancy in the human gut microbiome using ultra-deep metaproteomics. *Nat Commun* **14**, 3428.

45. Xu M, Lan R, Qiao L, Lin X, Hu D, Zhang S *et al.* (2023) Bacteroides vulgatus Ameliorates Lipid Metabolic Disorders and Modulates Gut Microbial Composition in Hyperlipidemic Rats. *Microbiol Spectr* **11**, e02517-22.

46. Vu H, Muto Y, Hayashi M, Noguchi H, Tanaka K, & Yamamoto Y (2022) Complete Genome Sequences of Three Phocaeicola vulgatus Strains Isolated from a Healthy Japanese Individual. *Microbiol Resour Announc* **11**, e01124-21.

47. Laraia BA (2013) Food Insecurity and Chronic Disease. Adv Nutr 4, 203–212.

48. Muñoz-Esparza NC, Latorre-Moratalla ML, Comas-Basté O, Toro-Funes N, Veciana-Nogués MT, & Vidal-Carou MC (2019) Polyamines in Food. *Front Nutr* **6**, 108.

49. Tofalo R, Cocchi S, & Suzzi G (2019) Polyamines and Gut Microbiota. Front Nutr 6, 16.

50. Kitada Y, Muramatsu K, Toju H, Kibe R, Benno Y, Kurihara S *et al.* (2018) Bioactive polyamine production by a novel hybrid system comprising multiple indigenous gut bacterial strategies. *Sci Adv* **4**, eaat0062.

51. Rio B del, Fernandez M, Redruello B, Ladero V, & Alvarez MA (2024) New insights into the toxicological effects of dietary biogenic amines. *Food Chem* **435**, 137558.

52. Deloyer P, Dandrifosse G, Kok N, & Delzenne NM (2000) Dietary Fructans Modulate Polyamine Concentration in the Cecum of Rats. *J Nutr* **130**, 2456–2460.

53. Noack J, Dongowski G, Hartmann L, & Blaut M (2000) The Human Gut Bacteria Bacteroides thetaiotaomicron and Fusobacterium varium Produce Putrescine and Spermidine in Cecum of Pectin-Fed Gnotobiotic Rats. *J Nutr* **130**, 1225–1231.

54. Noack J, Kleessen B, Proll J, Dongowski G, & Blaut M (1998) Dietary Guar Gum and Pectin Stimulate Intestinal Microbial Polyamine Synthesis in Rats. *J Nutr* **128**, 1385–1391.

55. Larqué E, Sabater-Molina M, & Zamora S (2007) Biological significance of dietary polyamines. *Nutrition* **23**, 87–95.

56. Ginty DD, Osborne DL, & Seidel ER (1989) Putrescine stimulates DNA synthesis in intestinal epithelial cells. *Am J Physiol-Gastrointest Liver Physiol* **257**, G145–G150.

57. McCormack SA & Johnson LR (1991) Role of polyamines in gastrointestinal mucosal growth. *Am J Physiol-Gastrointest Liver Physiol* **260**, G795–G806.

58. Bardócz S, Grant G, Brown DS, & Pusztai A (1998) Putrescine as a source of instant energy in the small intestine of the rat. *Gut* **42**, 24.

59. Deschak CI, Infante C, Mundo-Rosas V, Orjuela-Grimm M, & Aragón-Gama AC (2022) Food insecurity and coping strategies in international migrants in transit through Mexico. *J Migr Heal* **5**, 100099.

60. Martínez-Martínez OA, Gil-Vasquez K, & Romero-González MB (2023) Food insecurity and levels of marginalization: food accessibility, consumption and concern in Mexico. *Int J Equity Heal* **22**, 178.

61. Khan MT, Duncan SH, Stams AJM, Dijl JM van, Flint HJ, & Harmsen HJM (2012) The gut anaerobe Faecalibacterium prausnitzii uses an extracellular electron shuttle to grow at oxic– anoxic interphases. *ISME J* **6**, 1578–1585.

62. Khan MT, Browne WR, Dijl JM van, & Harmsen HJM (2012) How Can Faecalibacterium prausnitzii Employ Riboflavin for Extracellular Electron Transfer? *Antioxid Redox Signal* **17**, 1433–1440.

63. Wan Z, Zheng J, Zhu Z, Sang L, Zhu J, Luo S *et al.* (2022) Intermediate role of gut microbiota in vitamin B nutrition and its influences on human health. *Front Nutr* **9**, 1031502.

64. Cedillo YE, Kelly T, Davis E, Durham L, Smith DL, Kennedy RE *et al.* (2023) Evaluation of food security status, psychological well-being, and stress on BMI and diet-related behaviors among a sample of college students. *Public Heal* **224**, 32–40.

65. Davis I & Liu A (2015) What is the tryptophan kynurenine pathway and why is it important to neurotherapeutics? *Expert Rev Neurother* **15**, 719–721.

66. Baj A, Moro E, Bistoletti M, Orlandi V, Crema F, & Giaroni C (2019) Glutamatergic Signaling Along The Microbiota-Gut-Brain Axis. *Int J Mol Sci* **20**, 1482.

67. Christ M, Zange J, Janson CP, Müller K, Kuklinski P, Schmidt BM *et al.* (2001) Hypoxia modulates rapid effects of aldosterone on oxidative metabolism in human calf muscle. *J Endocrinol Investig* **24**, 587–597.

68. Bruening M, Woerden I van, Schaefer DR, Hruschka D, Brewis A, Whisner CM *et al.* (2018) Friendship as a social mechanism influencing body mass index (BMI) among emerging adults. *Plos One* **13**, e0208894.

69. Bruening M, Woerden I van, Todd M, & Laska MN (2018) Hungry to learn: the prevalence and effects of food insecurity on health behaviors and outcomes over time among a diverse sample of university freshmen. *Int J Behav Nutr Phy* **15**, 9.

70. Ellison B, Bruening M, Hruschka DJ, Nikolaus CJ, Woerden I van, Rabbitt MP *et al.* (2021) Viewpoint: Food insecurity among college students: A case for consistent and comparable measurement. *Food Policy* **101**, 102031.

Tables:

1 1				
Characteristic	Total (<i>n</i> = 85)	FI ($n = 13$)	FS ($n = 44$)	VAR (<i>n</i> = 28)
Age (years)	18.6 ± 0.8	18.5 ± 0.5	18.7 ± 0.9	18.6 ± 0.6
Sex, % (<i>n</i>)				
Male	38.8 (33)	53.8 (7)	45.5 (20)	21.4 (6)
Female	61.2 (52)	46.2 (6)	54.5 (24)	78.6 (22)
Race/ethnicity % (<i>n</i>)				
White	49.4 (42)	30.8 (4)	56.8 (25)	46.4 (13)
Black	11.8 (10)	15.4 (2)	11.4 (5)	10.7 (3)
Hispanic	21.2 (18)	23.1 (3)	20.5 (9)	21.4 (6)
Other	17.6 (15)	30.8.4 (4)	11.4 (5)	21.4 (6)
Height (cm)	168.6 ± 10.5	168.1 ± 12.8	170.3 ± 9.6	166.5 ± 10.8
Weight (kg)	71.1 ± 17.7	63.4 ± 13.3	73.1 ± 16.9	71.2 ± 19.9
Waist circumference (cm)	82.4 ± 13.5	75.9 ± 7.4	84.1 ± 13.5	82.6 ± 14.9
Waist/hip ratio	0.82 ± 0.07	0.78 ± 0.05	0.83 ± 0.07	0.82 ± 0.07
BMI (kg/m ²)	24.8 ± 5.1	22.2 ± 2.1	25.1 ± 4.9	25.5 ± 6.1
BMI Categories (kg/m ²), %				
<i>(n)</i>	5.9 (5)	7.7 (1)	4.5 (2)	7.1 (2)
$<18.5 \text{ kg/m}^2$	57.6 (49)	84.6 (11)	50.0 (22)	57.1 (16)
18.5-24.9 kg/m ²	24.7 (21)	7.7 (1)	31.8 (14)	21.4 (6)
25-29.9 kg/m ²	11.8 (10)	-	13.6 (6)	14.3 (4)
$\geq 30 \text{ kg/m}^2$				
MVPA (min/day)	48.2 ± 31.1	56.9 ± 33.7	46.7 ± 30.7	46.8 ± 31.2
Stress score	8.0 ± 2.3	8.9 ± 1.8	7.6 ± 2.4	8.2 ± 2.2
Depression score	1.9 ± 0.8	$2.3 \pm 0.7*$	1.7 ± 0.7	2.1 ± 0.8
Hours of nightly sleep ¹	7.4 ± 0.9	6.8 ± 1.3	7.6 ± 0.9	7.4 ± 0.9
Alcohol intake ²	3.2 ± 5.5	$7.6\pm10.0^*$	1.8 ± 2.9	3.7 ± 5.1
Fruit/vegetable intake ³	2.5 ± 0.8	2.5 ± 0.8	2.7 ± 0.9	2.2 ± 0.7

Table 1. Baseline demographic, anthropometric, and behavioral characteristics of study

 participants

Whole grain intake ⁴	0.8 ± 0.3	0.7 ± 0.4	0.8 ± 0.3	0.7 ± 0.3
Dairy intake ³	2.0 ± 0.9	2.1 ± 0.9	2.2 ± 0.9	1.8 ± 0.6
Red/processed meat intake ⁴	0.9 ± 0.8	0.9 ± 0.8	0.9 ± 0.8	0.9 ± 0.7
Daily sugar intake (g)	19.1 ± 7.2	21.5 ± 8.5	17.7 ± 5.8	20.3 ± 8.3
Daily fiber intake (g)	16.4 ± 3.6	16.8 ± 4.2	17.1 ± 3.5	15.0 ± 3.3

Data displayed as mean \pm SD, unless stated otherwise. Abbreviations: FI, food insecure; FS,

food secure; VAR, food variable; BMI, body mass index; MVPA, Moderate-to-vigorous

physical activity.

¹Average of combined weekday and weekend nightly sleep hours.

²Number of beverages over the last 7 days.

³Expressed as daily cup equivalents.

⁴Daily number of servings.

*Significant difference between FI and FS (pairwise comparison), P < 0.05.

Figure Legends:



Fig. 1. Overview of the study design and sample collection periods used in the present analysis.



Fig. 2. Gut microbiome and metabolome of college students by food security status. (A) Beta dispersion of Bray-Curtis dissimilarity for food insecure (FI), food secure (FS), and variable food security (VAR) classifications. NS = non-significant, ** $P \le 0.005$. Intra-individual shifts over the duration of the study period showing the 1st (timepoint 1 vs 2) and 2nd (timepoint 1 vs 3) time comparisons for the (B) microbiome, (C) estimated functional pathways, and (D) fecal metabolome. Scatter plots show Spearman's rank correlation coefficients and linear regression slopes by FS status for (E) gut microbiome, (F) estimated functional pathway, and (G) fecal metabolome stability of each feature (1.0 = most stable, 0 = least stable) vs. log-transformed abundance.



Fig. 3. Volcano plots display differential abundance of significant microbial species for (A) FI and (B) VAR and estimated functional pathways for (C) FI and (D) VAR with FS as the reference group (Q < 0.10). Fecal metabolome analysis showing (E) significant between-group metabolites as determined by GLM adjusted for time, sex, and BMI. Significant metabolites are displayed over the three time periods by log10 abundance and logFC values (ordered by FI positive to negative change) between the first and last sample for each FS classification. (F) Significantly altered metabolic pathways and (G) metabolite sets at the main chemical class level comparing FI to FI status (Q < 0.10). Note: Pathway impact scores summarize normalized topology measures of differential metabolites in each pathway. The enrichment ratio is computed by number of observed hits / expected number of hits.



Fig. 4. Grid-fused least absolute shrinkage and selection operator (GFLASSO) regression of fecal metabolites that best predicted species abundance of significant microbes associated with food insecure (FI) status.



Fig. 5. Exploratory analysis of the gut microbiome and metabolome of college students with variable food security (VAR) status. (A) Overview of VAR participants transition between food insecure (FI) and food secure (FS) states over the academic year. (B) Intra-individual shifts from FS to FI status in 19 individuals at the microbial species, estimated functional pathway, and fecal metabolome level. Alpha diversity metrics, (C) observed amplicon sequence variants and (D) phylogenetic diversity, capturing the shift from FS to FI status in the 19 individuals. (E) Relative abundance of the most dominant family level features of the 19 VAR participants at the FS and FI state. Families with a mean relative abundance of less than 1% are collapsed in the category "Other". NS = non-significant, ** P ≤ 0.01 , **** $P \leq 0.001$.