

- quel B, et al. Strain identification of *Mycobacterium tuberculosis* by DNA fingerprinting: recommendations for a standardized methodology. *J Clin Microbiol* 1993;31:406-409.
9. Bauer J, Yang Z, Poulsen S, Andersen AB. Results from 5 years of nationwide DNA fingerprinting of *Mycobacterium tuberculosis* complex isolates in a country with a low incidence of *M tuberculosis* infection. *J Clin Microbiol* 1998;36:305-308.
 10. Witte W, Cuny C, Braulke C, Heuck D, Klare I. Widespread dissemination of epidemic MRSA in German hospitals. *Eurosurveillance* 1997;2:25-28.
 11. Givney R, Vickery A, Holliday A, Pegler M, Benn R. Evolution of an endemic methicillin-resistant *Staphylococcus aureus* population in an Australian hospital from 1967 to 1996. *J Clin Microbiol* 1998;36:552-556.
 12. Cookson BD, Aparicio P, Deplano A, Struelens M, Goering R, Marples R. Inter-centre comparison of pulsed-field gel electrophoresis for the typing of methicillin-resistant *Staphylococcus aureus*. *J Med Microbiol* 1996;44:179-184.
 13. van Belkum A, Kluytmans J, van Leeuwen W, Bax R, Quint W, Peters E, et al. Multicenter evaluation of arbitrarily primed PCR of typing of *Staphylococcus aureus* strains. *J Clin Microbiol* 1995;33:1537-1547.
 14. Grundmann HJ, Towner KJ, Dijkshoorn L, Gerner-Smidt P, Maher M, Seifert H, et al. Multicenter study using standardized protocols and reagents for evaluation of reproducibility of PCR-based fingerprinting of *Acinetobacter* spp. *J Clin Microbiol* 1997;35:3071-3077.
 15. DelVecchio VG, Petroziello JM, Gress MJ, McCleskey FK, Melcher GP, Crouch HK, et al. Molecular genotyping of methicillin-resistant *Staphylococcus aureus* via fluorophore-enhanced repetitive-sequence PCR. *J Clin Microbiol* 1995;33:2141-2144.
 16. Deplano A, Vaneechoutte M, Verschraegen G, Struelens MJ. Typing of *Staphylococcus aureus* and *Staphylococcus epidermidis* strains by PCR analysis of inter-IS256 spacer length polymorphisms. *J Clin Microbiol* 1997;35:2580-2587.
 17. Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, et al. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 1995;23:4407-4414.
 18. Mazurek GH, Reddy V, Marston BJ, Haas WH, Crawford JT. DNA fingerprinting by infrequent-restriction-site amplification. *J Clin Microbiol* 1996;34:2386-2390.
 19. Riffard S, Lo Presti F, Vandenesch F, Forey F, Reyrolle M, Etienne J. Comparative analysis of infrequent-restriction-site PCR and pulsed-field gel electrophoresis for epidemiological typing of *Legionella pneumophila* serogroup 1 strains. *J Clin Microbiol* 1998;36:161-167.
 20. Fields PI, Blom K, Hughes HJ, Helsel LO, Feng P, Swaminathan B. Molecular characterization of the gene encoding H antigen in *Escherichia coli* and development of a PCR-restriction fragment length polymorphism test for identification of *E coli* O157:H7 and O157:NM. *J Clin Microbiol* 1997;35:1066-1070.
 21. Nachamkin I, Ung H, Patton CM. Analysis of HL and O serotypes of *Campylobacter* strains by the flagellin gene typing system. *J Clin Microbiol* 1996;34:277-281.
 22. Newcombe J, Dyer S, Blackwell L, Cartwright K, Palmer WH, McFadden J. PCR-single stranded conformational polymorphism analysis for non-culture-based subtyping of meningococcal strains in clinical specimens. *J Clin Microbiol* 1997;35:1809-1812.
 23. Harrington CS, Thomson-Carter FM, Carter PE. Evidence for recombination in the flagellin locus of *Campylobacter jejuni*: implications for the flagellin gene typing scheme. *J Clin Microbiol* 1997;35:2386-2392.
 24. van Belkum A, Riewerts Eriksen N, Sijmons M, van Leeuwen W, VandenBergh M, Kluytmans J, et al. Are variable repeats in the spa gene suitable targets for epidemiological studies of methicillin-resistant *Staphylococcus aureus* strains? *Eur J Clin Microbiol Infect Dis* 1996;15:768-769.
 25. Hoefnagels-Schuermans A, Peetermans WE, Struelens MJ, Van Lierde S, Van Eldere J. Clonal analysis and identification of epidemic strains of methicillin-resistant *Staphylococcus aureus* by antibiotyping and determination of protein A gene and coagulase gene polymorphisms. *J Clin Microbiol* 1997;35:2514-2520.
 26. Dean D, Schachter J, Dawson CR, Stephens RS. Comparison of the major outer membrane protein variant sequence regions of B/Ba isolates: a molecular epidemiologic approach to *Chlamydia trachomatis* infections. *J Infect Dis* 1992;166:383-392.
 27. Calderwood SB, Baker MA, Carroll PA, Michel JL, Arbeit RD, Ausubel FM. Use of cleaved amplified polymorphic sequences to distinguish strains of *Staphylococcus epidermidis*. *J Clin Microbiol* 1996;34:2860-2865.
 28. Perea Mejia LM, Stockbauer KE, Pan X, Cravioto A, Musser JM. Characterization of group A streptococcus strains recovered from Mexican children with pharyngitis by automated DNA sequencing of virulence-related genes: unexpectedly large variation in the gene (*sic*) encoding a complement-inhibiting protein. *J Clin Microbiol* 1997;35:3220-3224.
 29. van Leeuwen W, Sijmons M, Sluijs J, Verbrugh H, van Belkum A. On the nature and use of randomly amplified DNA from *Staphylococcus aureus*. *J Clin Microbiol* 1996;34:2770-2777.
 30. Kamerbeek J, Schoult L, Kolk A, van Achtveld M, van Soelingen D, Kuijper S, et al. Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J Clin Microbiol* 1997;35:907-914.
 31. Chee M, Yang R, Hubbell E, Berno A, Huang XC, Stern D, et al. Accessing genetic information with high-density DNA arrays. *Science* 1996;274:610-614.

Nosocomial Postoperative Endophthalmitis

Gina Pugliese, RN, MS
Martin S. Favero, PhD

Aaberg and coworkers evaluated the incidence of acute-onset (within 6 weeks after surgery) postoperative endophthalmitis between January 1, 1984, and December 30, 1994, at the Anne Bates Leach Eye Hospital, University of Miami Medical Center. The overall 10-year incidence of acute-onset postoperative endophthalmitis after intraocular surgery was 0.093% (54/58,123). Incidences by surgical category were cataract surgery with or without intraocular lens (IOL), 0.082% (34/41,654); pars plana vitrectomy (PPV), 0.046% (3/6,557); penetrating keratoplasty, 0.178% (5/2,805); secondary IOL place-

ment, 0.366% (5/1,367); glaucoma surgeries, 0.124% (4/3,233); combined trabeculectomy and cataract surgery, 0.114% (2/1,743); and combined penetrating keratoplasty and cataract surgery, 0.194% (1/515). The median visual acuity after endophthalmitis treatment was 20/200. Median acuities by procedure were cataract surgery with or without IOL, 20/133; PPV, no light perception; penetrating keratoplasty, 20/200; secondary IOL implantation, 20/40; glaucoma surgery, 20/80; and combined trabeculectomy and cataract surgery with or without IOL, 20/150.

The incidence of endophthalmitis was higher after secondary IOL implantation than after cataract extraction ($P=.008$). After treatment, visual acuity outcomes

were worse in the patients who developed endophthalmitis after PPV than after cataract extraction, glaucoma procedures, or secondary IOL implantation (analysis of variance [ANOVA], $P<.05$), and were better among the patients with secondary IOL implantation than after penetrating keratoplasty or PPV ($P<.05$, ANOVA). The results of this 10-year review from a large teaching center can serve as a source of comparison for other centers and future studies.

FROM: Aaberg TM Jr, Flynn HW Jr, Schiffman J, Newton J. Nosocomial acute-onset postoperative endophthalmitis survey. A 10-year review of incidence and outcomes. *Ophthalmology* 1998;105:1004-1010.