54. RADIAL VELOCITIES OF EARLY-TYPE STARS

J. P. KAUFMANN

Technische Universität, Berlin, Germany

With the Fehrenbach objective prism radial velocities of about 700 stars of type B0 to A0 were determined in two fields of the Southern Milky Way ($l^{\text{II}}=295^\circ$ $b^{\text{II}}=-0.6^\circ$; $l^{\text{II}}=320^\circ$ $b^{\text{II}}=-2.5^\circ$), with a mean error of ± 20 km s $^{-1}$. An additional photographic *UBV*-photometry with plates of the ADH-telescope at Boyden Observatory was accomplished. Minimum distances for the stars resulted from absorption-corrected magnitudes and a MK-spectral classification. About 200 stars lay at distances greater than 1.5 kpc from the sun. The largest distances determined were 5 kpc. From the radial velocities and distances circular velocities were derived and plotted against galactocentric distances R . Even within the possible error limits a positive velocity gradient showed up in the range $8 \text{ kpc} < R < 9.5 \text{ kpc}$, which French authors had already found for the region $10.5 \text{ kpc} < R < 12.5 \text{ kpc}$. If there do not exist significant deviations from circular motion for these stars, a conformity with Schmidt's 1965 model cannot be obtained.