

4. Sepkowitz KA, Raffalli J, Riley L, Kiehn TE, Armstrong D. Tuberculosis in the AIDS era. *Clin Microbiol Rev* 1995;8:180-199.
5. Wurtz R. Administrative controls for TB: 'Keep doing what you've always done, and you'll get what you always got.' *Infect Control Hosp Epidemiol* 1996;17:409-411.
6. Pegues CF, Johnson DC, Pegues DA, Spencer M, Hopkins CC. Implementation and evaluation of an algorithm for isolation of patients with suspected pulmonary tuberculosis. *Infect Control Hosp Epidemiol* 1996;17:412-418.
7. Mathur P, Sacks L, Auten G, Sall R, Levy C, Gordin F. Delayed diagnosis of pulmonary tuberculosis in city hospitals. *Arch Intern Med* 1994;154:306-310.
8. Counsell SR, Tan JS, Dittus RS. Unsuspected pulmonary tuberculosis in a community teaching hospital. *Arch Intern Med* 1989;149:1274-1278.
9. Centor R, Schwartz J. An evaluation of methods for estimating the area under the receiver operating characteristic (ROC) curve. *Med Decis Making* 1985;5:149-156.
10. Hanley JA, McNeil BL. The meaning and use of the area under the receiver operating characteristic (ROC) curve. *Radiology* 1982;143:29-36.
11. Wasson JH, Sox HC, Neff RK, Goldman L. Clinical prediction rules. Applications and methodological standards. *N Engl J Med* 1985;313:793-799.
12. Scott B, Schmid M, Nettleman MD. Early identification and isolation of inpatients at high risk for tuberculosis. *Arch Intern Med* 1994;154:326-330.
13. Cohen R, Muzaffar S, Capellan J, Azar H, Chinikamwala M. The validity of classic symptoms and chest radiographic configuration in predicting pulmonary tuberculosis. *Chest* 1996;109:420-423.
14. Antonucci G, Girardi E, Raviglione MC, Ippolito G. Risk factors for tuberculosis in HIV-infected persons. A prospective cohort study. *JAMA* 1995;274:143-148.
15. El-Solh A, Mylotte J, Sherif S, Serghani J, Grant BJB. Validity of a decision tree for predicting active pulmonary tuberculosis from clinical and radiographic data. *Am J Respir Crit Care Med* 1997;155:1711-1716.
16. Centers for Disease Control and Prevention. Tuberculosis morbidity—United States, 1995. *MMWR* 1996;45:365-370.
17. Brobrowitz ID. Active tuberculosis undiagnosed until autopsy. *Am J Med* 1982;72:650-658.
18. Couser JI, Glassroth J. Tuberculosis. An epidemic in older adults. *Clinics Chest Med* 1993;14:491-499.
19. Pitlik SD, Fainstein V, Bodey GP. Tuberculosis mimicking cancer: a reminder. *Am J Med* 1984;76:822-825.
20. Fischl MA, Daikos GL, Uttamchandani RB, et al. Clinical presentation and outcome of patients with HIV infection and tuberculosis caused by multiple-drug-resistant bacilli. *Ann Intern Med* 1992;117:184-189.
21. Given MJ, Khan A, Reichman LB. Tuberculosis among patients with AIDS and a control group in an inner-city community. *Arch Intern Med* 1994;154:640-645.

---

## Nosocomial Aspergillosis Associated With Construction

**Gina Pugliese, RN, MS**  
**Martin S. Favero, PhD**

The CDC recently assisted with an investigation of an outbreak of invasive aspergillosis among patients in a Maryland hospital. A case-control study identified seven case patients; five were rheumatology patients hospitalized on two wards. Rheumatology case patients were more likely than randomly selected controls with invasive aspergillosis to have longer hospitalizations both in current and prior admission, to receive high doses of intravenous immunosuppressive agents, or to receive immunosuppressive agents for a long period of time.

An evaluation of the environment revealed that construction areas in the hospitals were not sealed off from patient-care areas nor under negative pressure in relation to patient-care areas. In addition, the air flow from patient rooms was not positive in relation to the hallway, and the rooms had only 1.6 air exchanges per hour. The risk of aspergillosis among highly immune-suppressed patients exposed to air from hospital construction has been well documented. This study documents that rheumatology patients, particularly those receiving high-dose intravenous immunosuppressive therapy, should be considered at increased risk of invasive *Aspergillus* species.

Measures must be taken to hospitalize these patients away from construction activities. Additional measures to reduce risk include the creation of barriers to seal off all construction areas and housing high-risk patients in rooms with positive pressure and high-efficiency particulate air filtration. The CDC has published guidelines for the prevention of nosocomial pneumonia that contain additional recommendations.

FROM: Garrett DO, Jochimsen E, Banerjee C, et al. Invasive *Aspergillus* species infections in rheumatology patients. Presented at the 46th Annual Epidemic Intelligence Service Conference; April 14-18, 1997; Atlanta, GA.