

Editorial

Technological innovations in mental healthcare: harnessing the digital revolution

Chris Hollis, Richard Morriss, Jennifer Martin, Sarah Amani, Rebecca Cotton, Mike Denis and Shôn Lewis

**Summary**

Digital technology has the potential to transform mental healthcare by connecting patients, services and health data in new ways. Digital online and mobile applications can offer patients greater access to information and services and enhance clinical management and early intervention through access to real-time patient data. However, substantial gaps exist in the evidence base underlying these technologies. Greater patient and clinician involvement is needed to evaluate digital technologies and ensure they target unmet needs, maintain public trust and improve clinical outcomes.

Declaration of interest

C.H. receives research funding from Shire Pharmaceuticals paid to his employing institution. R.C. reports that Janssen Cilag Ltd have provided a grant to support an upcoming Mental Health Network event on the subject of technology and mental health services. S.L. is chief investigator on the Clintouch project. C.H., R.M. and J.M. are all involved in the National Institute of Health Research (NIHR) MindTech Healthcare Technology Co-operative.

Chris Hollis (pictured) is Professor of Child and Adolescent Psychiatry at the University of Nottingham, and Director of the National Institute of Health Research (NIHR) MindTech Healthcare Technology Co-operative (HTC); his research interests are in attention-deficit hyperactivity disorder, neurodevelopmental disorders and the evaluation and implementation of digital technologies in mental healthcare. Richard Morriss is Professor of Psychiatry at the University of Nottingham, Mood Disorders lead for the NIHR MindTech HTC, Director of Research for NIHR CLAHRC East Midlands and leads externally funded research in randomised controlled trials of video-based psychological treatment for health anxiety and self-harm, internet programmes for managing early warning signs of bipolar disorder and facial recognition to objectively measure the severity of depression. Jennifer Martin is NIHR MindTech HTC programme manager; prior to joining MindTech she was a senior research fellow in human factors at the University of Nottingham where her research was focused on the development and evaluation of healthcare technology. Sarah Amani is a mental health nurse with a special interest in early intervention and youth mental health. Her recent work includes the co-development of mobile health apps, telehealth solutions and piloting the use of social media in mental health services. Rebecca Cotton is Director of Policy for the Mental Health Network, part of the NHS Confederation. She has a strong interest in e-health and has previously written on the subject. Mike Denis has recently joined Oxford Academic Health Science Centre/Network bringing successful experience in electronic medical records and patient health records design and leading an Oxford, Cambridge, London research informatics programme. Shôn Lewis is Director of the Institute of Brain, Behaviour and Mental Health at the University of Manchester and Mental Health Academic Lead for the Manchester Academic Health Sciences Centre. He leads an European Union- and Medical Research Council-funded programme of connected health research for people with severe mental illness.

The digital revolution

Advances in technology and particularly mobile digital information and communication technology continue at an exponential rate, making it possible to communicate, obtain information and access and buy goods and services in new ways. In 2013, 36 million adults in the UK (73%) accessed the internet every day and 72% of all adults bought goods or services online.¹ Access to the internet using a mobile phone more than doubled between 2010 and 2013 to 53%. In total, 51% of UK adults now own a smartphone (almost doubled from 2011) and 24% own a tablet. Also, 43% of adults now use the internet to seek health-related information, an increase from 18% in 2007.¹ Significant sections of society are becoming increasingly familiar and comfortable with using technology for a wide range of transactions.

E-health and m-health (increasingly collectively referred to as 'connected health', see Appendix) describe the delivery of healthcare by electronic means via the internet using a variety of devices including mobile phones, remote monitoring devices and other wireless devices. These digital technologies can greatly improve access to mental healthcare and treatment adherence by enabling services to be delivered more flexibly and tailored to individual patient needs. Recent developments in sensor technology, online psychological therapy and remote video consultation, mobile applications ('apps') and gaming all present real opportunities to engage and empower patients and create novel approaches to both assessment and intervention for mental health problems.

The drivers for technological transformation in mental health services

The growth in demand for mental healthcare exceeds available UK National Health Service (NHS) resources, and this gap is likely to increase up to 2020. Cost pressures require that more is done for less and providers therefore must find innovative ways to deliver services. The UK Government's mental health strategy 'No health without mental health'² recommends the increased use of information and communication technology (ICT) to improve care and access to services. UK Government initiatives such as 'Digital First' aim to reduce unnecessary face-to-face contact between patients and healthcare professionals with NHS trusts encouraged to replace these unnecessary face-to-face contacts with video-based remote consultations. However, service users consistently report dissatisfaction with a 'top-down' one-size-fits-all approach. Connected health innovations have the potential to offer great flexibility and to be more patient-centred.

Digital mental healthcare

E-mental health is not just about technology, but represents a cultural change in mental healthcare by empowering patients to exercise greater choice and control. For example, the provision of online psychological interventions, potentially accessible 24 h a day, 7 days a week, is increasing rapidly using synchronous video/voice or asynchronous text communication. Examples of providers to the NHS include Xenzone (www.xenzone.com),

PsychologyOnline (www.psychologyonline.co.uk) and Big White Wall (BWW) (www.bigwhitewall.com)

Mental health professionals have historically been reliant on face-to-face consultations in clinic settings away from the normal lives of their patients. Mobile apps are now available that allow patients to record their mood, behaviour and activities in real-time using well-validated measures such as the PHQ-9 depression scale.³ Patients can track their condition using their own mobile device over time and share this information with their clinician. Potential benefits to patients include greater engagement in their care and earlier detection of problems, more timely adjustment of treatment and shared decision-making. Sensors such as accelerometers, gyroscopes, microphones and cameras, which are now standard parts of smartphones and other mobile devices, mean that it will be possible to continuously and passively collect objective data that can give additional insight into a person's behaviour and activities. For example, detecting subtle changes in sleep and activity patterns constitute important early warning signs of relapse and can prompt early intervention in depression and bipolar disorder.⁴

In mental healthcare, assessment and diagnosis is still largely based on subjective clinical judgements about symptoms and behaviour. Technological innovations have the potential to bring more objectivity and reliability to these processes of assessment, diagnosis and monitoring. Affective computing is a branch of computer science that aims to develop automated assessment of a person's mood by analysis of their facial expression, speech rate and tone of voice.⁵ Automated analysis of facial expression has been shown to correspond closely with clinical ratings of depression⁵ and could potentially augment clinical assessment in the near future.

In addition, ICT can address issues such as social isolation and the relative inaccessibility of mental health services to young people. The rapid increase in popularity of social media platforms means that people are increasingly able to access information and support from peers and professionals in new and informal ways. Mental health problems disproportionately affect younger people, and this group is also the biggest user of ICT; social media may widen access for groups who find it difficult to access traditional services.

It is critical, however, to consider the challenges of introducing new technology to mental healthcare. The so-called 'digital divide', describing a gulf between those who have ready access to a computer and the internet and those who do not, means that some groups of patients (for example, older adults, people who are homeless and people with intellectual disabilities) could experience barriers to accessing mental healthcare delivered using ICT. A study of patients with mental health disorders found that their use of technology (computers, mobile and smartphones) was similar to the general population, with older people reporting less familiarity, access and confidence with these technologies.⁶ Older people experiencing psychosis report a significant desire to increase their use of computers, suggesting that a cost and skills gap, rather than indifference, is the reason for lower rates of computer usage.⁶ Tailoring the device to the needs of the individual is particularly important, as witnessed by the rapid uptake of tablet devices by older people.

Data protection, privacy and security: establishing public and patient trust

Maintaining public trust when handling and sharing personal health data is paramount and this requires serious public engagement over issues of consent, data security and privacy.

There is an even greater need to address these ethical issues in the field of mental healthcare as the data are often highly sensitive personal information. Although technological developments hold great promise, the lessons from the aborted initial roll out of *Care.data* by NHS England⁷ demonstrate that public acceptance and large-scale population utilisation are unlikely unless steps are taken to safeguard the legal and ethical rights of the public and patients.

M-health apps

The past 5 years have seen a rapid increase in the number of m-health apps, including many targeted at mental health and well-being. Some of these are aimed at healthcare professionals but the majority are targeted at the general public, although many are developed without significant user (patient or clinician) involvement.⁸ The majority of apps are sold via commercial marketplaces but the NHS has also established a health apps library, which contains (as of 1 February 2015) 23 apps categorised under mental health, although the safety and efficacy of many of these appears unproven.

The rapid emergence of apps combined with the relative low cost and speed with which they can be developed and placed on the market, has resulted in the NHS and regulatory bodies struggling to keep pace with the evolving field of m-health. Guidance on the regulatory requirements for m-health apps was only published by the Medicines and Healthcare products Regulatory Agency (MHRA) in March 2014⁹ and NHS England is currently working with the US Food and Drug Administration on a bilateral framework for regulation of m-health apps. Evaluating the safety and efficacy of m-health interventions should be subject to the same rigour as evaluations required for new drug or psychological treatments.

Some examples of apps that have been developed for mental health include ClinTouch (www.clintouch.com), My Journey (www.sabp.nhs.uk/eiip/app), Buddy App (www.buddyapp.co.uk) and WellHappy (apps.nhs.uk/app/wellhappy/). Typically, these apps include a symptom tracker and diary function, appointment and medication reminders and motivational prompts. These examples have been developed with user and clinician involvement and incorporate evidence-based principles of care (for example, early intervention improves outcomes in psychosis). Although early-stage pilot evaluations suggest that these apps are safe and their use is acceptable to patients and clinicians, further research is needed to demonstrate clinical and cost-effectiveness in routine NHS settings. The task of building an evidence base and framework for evaluating the rapidly increasing number of m-health apps and other digital products is being led by NIHR MindTech Healthcare Technology Co-operative and should be a priority area for NHS applied health research.

The 'big data' challenge for mental health

Digital health technologies, including, electronic medical records, imaging, e-prescribing and the increasingly prevalent use of connected devices to monitor health, have the potential to generate large amounts of data that could be of considerable benefit to researchers and policymakers.

There are three processes required to achieve these benefits: unlocking value in electronic medical records (EMRs); new forms of access that allow patients direct control (patient health records, PHRs); and pervasive computing and connected devices (m-health) to enable expansion of the 'real-time' patient data.

Increasingly, data accessed through the EMR is being used within clinical, epidemiological and public health research. In each case use of the EMR enables scaling of such studies; sometimes by orders of magnitude in both size and speed. Challenges to the use of the EMR in such research are not trivial and include technical (can data be accessed and linked in a format that enables analysis from multiple data sources?), governance (security, privacy, access control) and scientific (is the data of sufficient quality?); however, these challenges are being met both for meta-data such as diagnosis and hospital episodes, and for patient level-data including complex text or narrative data.¹⁰

An example of the use of complex EMR data is the Clinical Record Interactive Search (CRIS) system established by the Maudsley Biomedical Research Centre in London.¹⁰ CRIS allows researchers to view de-identified records provided that research studies are approved by an oversight committee with full user engagement. The system has been used in many studies, an example being the extraction of data from a single NHS trust to show substantially increased mortality in people with serious mental illness.¹¹

Conclusion

Despite the clear potential of digital technology to connect people and health data in new ways, there is currently insufficient evidence to suggest that this potential is being fully realised, with uptake being limited and outcomes being largely anecdotal and unpublished. There are a number of key challenges: first, ensuring that patients and their needs remain at the centre of technology development and implementation; second, rapidly increasing the evidence base for the clinical effectiveness of digital technology; third, ensuring that the opportunity provided by data sharing between patients, carers and clinicians does not threaten privacy and undermine public trust. Finally, patients, clinicians and NHS commissioners require an agreed framework to evaluate the core features of new technologies including usability, content, safety, clinical- and cost-effectiveness.

Chris Hollis, PhD, MRCPsych, **Richard Morriss**, MD, FRCPsych, **Jennifer Martin**, BSc, PhD, NIHR MindTech Healthcare Technology Co-operative, Institute of Mental Health, University of Nottingham, Nottingham; **Sarah Amani**, PGDip Mental Health Nursing, PGDip CBT, Oxford Academic Health Science Network, Oxford; **Rebecca Cotton**, BSc, MRes, Mental Health Network – NHS Confederation, London; **Mike Denis**, MSc, Oxford Academic Health Science Centre/Network, Oxford; **Shôn Lewis**, MD, FMedSci, Institute of Brain, Behaviour and Mental Health, University of Manchester, Manchester Academic Health Science Centre, Manchester Mental Health and Social Care Trust, Manchester, UK

Correspondence: Chris Hollis, NIHR MindTech Healthcare Technology Co-operative, Institute of Mental Health, University of Nottingham, Jubilee Campus, University of Nottingham Innovaton Park, Triumph Road, Nottingham NG7 2TU, UK. Email: chris.hollis@nottingham.ac.uk

First received 8 Aug, accepted 15 Nov 2014

Appendix

Definitions in digital healthcare

E-health

The transfer of health resources and healthcare by electronic means.¹² It encompasses three main areas:

- the delivery of health information, for health professionals and health consumers, through the internet and telecommunications;
- using the power of information technology and e-commerce to improve public health services, for example, through the education and training of health workers;
- the use of e-commerce and e-business practices in health systems management.

M-health

Medical and public health practice supported by mobile devices, such as mobile phones, patient-monitoring devices, personal digital assistants and other wireless devices.¹³ M-health involves the use and capitalisation on a mobile phone's core utility of voice and short messaging service (SMS) as well as more complex functionalities and applications including general packet radio service (GPRS), third- and fourth-generation mobile telecommunications (3G and 4G systems), global positioning system (GPS) and Bluetooth technology.

E-mental health

The use of ICT to support and improve mental health, including the use of online resources, social media and smartphone applications. Two types of e-mental health are commonly referred to: web interventions and mobile applications.

Telemental health

A subset of telehealth that uses video-conferencing technology to provide mental health services from a distance. It includes telepsychology, telepsychiatry, telemental health nursing and telebehavioural health.

Funding

C.H. receives research funding from the National Institute of Health Research (NIHR) and European Union paid to his employing institution. R.M. receives research funding from the NIHR Collaboration for Leadership in Applied Health Research and Care East Midlands paid to his employing institution. R.C. is employed by the Mental Health Network, which receives funding from providers of NHS-funded mental health services in England by way of membership fees. S.L. is chief investigator on the Medical Research Council-funded Clintouch project. This editorial is based on research conducted by the NIHR MindTech Healthcare Technology Co-operative, which is funded by the National Institute of Health Research. The views represented are the views of the authors alone and do not necessarily represent the views of the Department of Health in England or the National Institute for Health Research.

References

- Ofcom. *Communications Market Report 2013*. Ofcom, 2013.
- HM Government. *No Health Without Mental Health*. Department of Health, 2011.
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; **16**: 606–13.
- Prociow PA, Crowe JA. Towards personalised ambient monitoring of mental health via mobile technologies. *Technol Health Care* 2010; **18**: 275–84.
- Valstar M, Schuller B, Smith K, Eyben F, Jiang B, Bilakhia S, et al. *AVEC 2013: The Continuous Audio/Visual Emotion and Depression Recognition Challenge. Proceedings of the 3rd ACM International Workshop on Audio/Visual Emotion Challenge*. ACM, 2013.
- Ennis L, Rose D, Denis M, Pandit N, Wykes T. Can't surf, won't surf: the digital divide in mental health. *J Ment Health* 2012; **21**: 395–403.
- Vallance C. NHS Care.data information scheme 'mishandled'. *BBC News* 2014; April 18.
- Craven MP, Selvarajah K, Miles R, Schnädelbach H, Massey A, Vedhara K, et al. User requirements for the development of smartphone self-reporting applications in healthcare. In *Human-Computer Interaction Applications and Services* (ed. M Kurosu). Springer, 2013.
- Medicines and Healthcare Regulatory Authority. *Guidance on Medical Device Stand-Alone Software (Including Apps)*. MHRA, 2014 (<http://www.mhra.gov.uk/Howweregulate/Devices/Software/index.htm>).
- Stewart R, Soremekun M, Perera G, Broadbent M, Callard F, Denis M, et al. The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry* 2009; **9**: 51.
- Chang CK, Hayes RD, Perera G, Broadbent MT, Fernandes AC, Lee WE, et al. Life expectancy at birth for people with serious mental illness and other major disorders from a secondary mental health care case register in London. *PLoS One* 2011; **6**: e19590.
- World Health Organization. *Glossary of Health Terms*. WHO, no date (<http://www.who.int/trade/glossary/story021/en/>).
- World Health Organization. *mHealth New Horizons for Health through Mobile Technologies. Global Observatory for eHealth Series – Volume 3*. WHO, 2013.