

Pathways to sympathies for violent protest and terrorism

Kamaldeep Bhui, Maria Joao Silva, Raluca A. Topciu and Edgar Jones

Background

Radicalisation is proposed to explain why some individuals begin to support and take part in violent extremism. However, there is little empirical population research to inform prevention, and insufficient attention to the role of psychiatric vulnerabilities.

Aims

To test the impact of depressive symptoms, adverse life events and political engagement on sympathies for violent protest and terrorism (SVPT).

Method

A cross-sectional survey of a representative sample of Pakistani and Bangladeshi men and women from two English cities. Weighted, multivariable, logistic regression yielded population estimates of association (odds ratio (OR) and 95% confidence intervals) against a binary outcome of SVPT derived from a three-group solution following cluster analysis.

Results

Depressive symptoms were associated with a higher risk of

SVPT (OR=2.59, 95% CI 1.59–4.23, $P<0.001$), but mediated little of the overall effects of life events and political engagement, which were associated with a lower risk of SVPT (death of a close friend: OR=0.24, 95% CI 0.07–0.74; donating money to a charity: OR=0.52, 95% CI 0.3–0.9).

Conclusions

Independent of SVPT associations with depressive symptoms, some expressions of social connectedness (measured as life events and political engagement) are associated with a lower risk of SVPT.

Declaration of interest

K.B. is Editor of the *BJPsych* and was not involved in any editorial decisions about this manuscript. The data collection was commissioned from Ipsos MORI. K.B. and E.J. are trustees of Careif (who part funded the statistical analysis).

Copyright and usage

© The Royal College of Psychiatrists 2016.

In the aftermath of attacks on civilians in Western cities, psychiatrists, psychologists and criminal justice agencies have sought to understand the role of mental illness in terrorist offending.^{1–3} An association has been reported between severe mental illness and terrorists who operate independently of others; 30–40% of these so-called ‘lone wolves’ appear to show signs of mental illness, isolation and marginalisation, which may make them suggestible and vulnerable to persuasion by terrorist ideology.⁴ Mental illnesses are 13 times more likely to occur in ‘lone wolves’ than in group-based terrorists, but mostly as a result of severe mental illnesses such as psychoses rather than depression.^{4,5} However, even lone wolves are not always isolated, suggesting there is no uniform profile.⁵ Compared with lone wolves, those who conduct school attacks and assassins are more likely to have signs of depression, despair and suicidal ideas, and a history of violence.⁶

By contrast, terrorist plots and attacks in the UK, France, USA and Canada were, on the whole, organised by people without obvious symptoms of mental illnesses.⁷ The perpetrators were born and educated in the countries that they attacked, and they seemed to be socially integrated. Without evidence of previous criminal activities or adverse life events, they fall into the category of offenders called ‘late starters.’⁷ Links with organised terrorist groups are not easily identified, but communications through social media and websites, as well as exposure to extremist ideology, are often revealed during criminal investigations to have contributed to adopting extremist ideology.⁷ Whether hidden or subthreshold mental illness plays a part in the recruitment of this group of ostensibly ordinary individuals is underresearched, but radicalisation is the process that is proposed by governments to explain this phenomenon.

The term radicalisation was first used following the Madrid 2004 and London 2005 bombings,⁸ although its definition

continues to evolve. The UK *Prevent* policy defined radicalisation as the process by which a person comes to support terrorism and forms of extremism leading to terrorism. We adopt a broader definition: a social and psychological process by which ordinary individuals come to sympathise with, and then make a commitment to, terrorist activities.^{9,10} However, all definitions are explicit that radicalisation can exist without violence and extremist behaviour. Indeed, the 2011 revision of *Prevent* includes a broadening of what is considered radical, encompassing vocal or active opposition to fundamental British values, democracy, the rule of law, individual liberty, mutual respect and tolerance of different faiths and beliefs.¹¹ Despite the evolving shift in preventive frameworks and terminology, there is little empirical research into the process of radicalisation, and how this might differ in populations and specific groups, or about the role of psychological factors or common mental illnesses. The dominant explanation about radicalisation is that poverty, unemployment, discrimination, political isolation and cultural marginalisation lead to grievances, which in turn foster increased receptivity to political violence as a solution.⁷ Adverse life events and poor civic engagement are associated with depression^{12,13} and poor health^{14,15} and are reported to engender extremism,^{7,16–18} suggesting some shared aetiologies for depression and extremism.

Sympathies for violent protest and terrorism (SVPT) are regarded as a ‘pre-radicalisation’ phase when individuals are vulnerable to recruitment to terrorist causes. We developed a measure of SVPT as a marker of susceptibility to engagement with extremist groups and actions.¹⁹ We previously found an association between depressive symptoms and SVPT,¹⁹ suggesting that these symptoms may drive cognitive biases leading to the adoption of extremist ideology and violence.¹ The lack of hope and pessimism that characterise depression may increase the appeal

of potent ideologies that promote agency and empowerment, and give purpose and meaning, even if related to criminal actions.^{1,20} Further evidence in support of a potential role for depression comes from a recent meta-analysis that shows a three-fold increase in the risk of violence among those with depression.²¹ Given the associations that exist between depression, social adversity²² and marginalisation,^{23,24} we hypothesised that depressive symptoms mediate relationships with SVPT.

Method

Participants

The study included 608 people of Pakistani and Bangladeshi family origin, aged between 18 and 45, of Muslim heritage and living in Bradford and East London. Bradford, an industrial town in northeast England, is home to a significant proportion of the Muslim population who live in isolated and traditional communities associated with deprivation.^{19,20} East London has a substantial and well-established Muslim population living in a region of greater religious and cultural diversity with wider opportunities for employment.

Participants were recruited by proportional quota sampling. This is a standard method that sets quotas for participants on a range of demographic factors and ensures that the sample interviewed is representative of the target population. Quota sampling offers an alternative to probability sampling and is often used in market research and national surveys as an efficient sampling strategy.²⁵ Using UK Census 2001 data, quotas were set for each region to reflect the key demographic variables of those living there. Target quotas were set for age (18–30 years and 31–45 years) gender, work status (working full time, not working full time) and ethnicity (Pakistani and Bangladeshi). Data were collected from Pakistani and Bangladeshi men and women of Muslim heritage, given the concerns expressed in the media and in counter-terrorism responses focused on South Asians and people of Muslim heritage. In addition, these UK communities experience social adversity and marginalisation, and in our preliminary community discussions endorsed the need for more empirical research to inform preventive actions. Individuals living within a sampling unit were identified by door knocking and offered a computer-assisted interview if they gave informed consent. Flash cards were used to simplify the process of answering questions with choices.

Data collection was undertaken by Ipsos MORI Social Research Institute. All questions were refined following eight pilot interviews to check wording, sensitivity and questioning styles. Interviewers from Ipsos MORI were recruited from the local population, and had significant experience of research into sensitive topics including religion and terrorism. Questions were asked in a computer-assisted format with prompts and cues so that sensitive questions could be answered anonymously, out of sight of the interviewer. Piloting and the main study itself found that language or religious matching were not requested or necessary, although available. Informed consent was recorded by checking an appropriate box before proceeding with the survey. Ethical approval was received from Queen Mary University of London Research Ethics Committee.

Measuring SVPT

The 16-item measure designed to assess early signs of radicalisation asked about support for, or condemnation of, acts of protest characterised by differing levels of violence and extremist behaviour.¹⁹ Sympathies are regarded as an early phase of vulnerability to radicalisation.¹⁰ The wording and items were

developed through participatory discussions.¹⁸ We consulted Muslim and non-Muslim researchers and members of local community panels (consisting of local charities and mental health and educational organisations and religious institutions) about how to measure radicalisation.¹⁹ The 16 core questions identified for inclusion had been proposed by and then reviewed by the community panel, and tested in pilot interviews.¹⁹ The questions were specifically aimed at being inclusive, rather than focusing on specific religious, cultural or ethnic groups as respondents. The responses were in the form of a seven-item Likert scale, ranging from –3 (completely condemn) to +3 (completely sympathise). For all but two items a higher score indicated greater support for violent protest and terrorism. These two items, which asked about sympathies for or condemnation of the UK government's decision to send British troops to Afghanistan and Iraq, were reverse-scored as condemnation might reflect a more radicalised perspective. The 16-item scale was found to be highly reliable, with a Cronbach's $\alpha = 0.81$.

A cluster analysis of the 16-item measure of SVPT produced a three-group solution: a group that was least sympathetic (group 1, $n = 93$), a large intermediary group (group 2, $n = 423$) and a most sympathetic group (group 3, $n = 92$). The methods for generating clusters are already published;²⁰ a specific method of cluster analysis, a classification likelihood method, was applied to the 16 items.^{26,27} The Bayesian information criterion was used to determine the number of clusters. The clustering was carried out on the principal component scores from a principal components analysis of the original 16-item scores. The clustering was carried out using different numbers of principal component scores, and the most stable solution found was the one with the three groups.

Depressive symptoms were associated with membership of group 3 (which contains individuals who show most sympathy towards violent protest and terrorism) when compared with groups 1 and 2 combined, or group 2 alone. Depressive symptoms were not associated with membership of group 1 when compared with group 2, or with groups 2 and 3 combined. Therefore, we aggregated groups 1 and 2 to form a reference group to offer a comparison with group 3. Conceptually, this comparison is the most important in terms of understanding radicalisation.

Employment and education

Employment status was grouped into a three-level variable: employed (full time, part time or self-employed), unemployed, and an aggregated group who reported as retired, unwell or a housewife. Educational status included those having no qualifications *v.* any qualifications below degree level (GCSE/O-level/CSE, vocational qualifications such as NVQ1+2, A-level or equivalent such as NVQ3), and those having a degree (bachelor, master or doctorate).

Adverse life events

The measure of adverse life events included injury, bereavement, separations, loss of job, financial crisis, problems with the police or courts, theft and other major stressful events in the preceding 12 months.²⁸ For each adverse life event, a binary variable (yes/no) was derived.

Political engagement

The questions to assess political engagement were drawn from the UK Department of Communities and Local Government Citizenship Survey.²⁹ These questions addressed voting in local council elections, political discussions, signing a petition,

donations to a charity or campaigning organisation, payment of membership fees to a charity or campaigning organisation, voluntary work, a boycott for political, ethical, environmental or religious reasons, political views expressed online, attendance at a political meeting, donations to or membership of a political party, and participation in a demonstration or march.²⁹ For each specific item of political engagement, a binary variable (yes/no) was derived.

Depressive symptoms

Depressive symptoms were measured by the Patient Health Questionnaire (PHQ-9), a screening measure commonly used in primary care and specialist mental health services, with well-established validated thresholds for indicating risks of clinical depression.³⁰ For the analysis, the total PHQ-9 score was classified into the following categories: PHQ score < 5 and PHQ score \geq 5, where the latter indicates 'probable clinical depression'.

Statistical analysis

A binary measure of sympathies for violent protest and terrorism was used in univariable and multivariable logistic regression models weighted for the sampling strategy and for non-response, thus yielding estimates attributable to the population from which the sample was drawn.

- (a) All sociodemographic, life event and political engagement variables were assessed for associations with the binary outcome of SVPT and probable clinical depression. This information was used to undertake two further analyses.
- (b) All variables significantly associated with the binary SVPT in the univariable analyses were included in the multivariable logistic regression models with one model for each life event and for each action of political engagement. These models were adjusted for age, gender, employment status, education level and depression.

- (c) If specific life events and political engagement actions were significantly associated with both depressive symptoms and with SVPT, mediation models were employed to assess to what extent depressive symptoms explained the associations of life event and political engagement with SVPT.³¹ Where conditions of mediation analyses were met, we estimated what proportion of the direct relationship was explained by the indirect relationship through depressive symptoms.

The cluster analysis was implemented using the *mclust* package in *R*. All other analyses were performed in Stata 14. Statistical significance was considered at $P < 0.05$.

Results

Demographic, health and social characteristics

Tables 1 and 2 show the distribution of demographic, social and health characteristics. The sample is primarily composed of 26 to 35-year olds, most of whom are employed and educated; 61% of this sample have a personal income between £5000 and £24 999. A total of 10% of the participants had experienced the death of a close friend or relative, and 10% had encountered a serious problem with a close friend, neighbour or relative; 62% of the sample voted in the last local council election, 41% donate money to charity and 19% undertake voluntary work. Only 1.4% reported a problem with the police or courts and 6% had been seeking a job for one month or more in the preceding year. A fifth (22%) reported a PHQ-9 score indicating probable clinical depression.

Univariable analyses

Table 3 shows that, contrary to expectation, those who had experienced the death of a close friend, a serious problem with a close friend, neighbour or relative, or another major event were less likely to have SVPT. People who had problems with the police

Table 1 Demographic characteristics by sympathies for violent protest and terrorism (weighted)

Characteristic	n (%)		
	Groups 1 and 2 (n = 488)	Group 3 (n = 120)	All (n = 608)
Age groups, ^a years			
18–25	113 (23.68)	42 (35.29)	155 (25.98)
26–35	249 (52.14)	42 (35.82)	291 (48.91)
36–45	116 (24.18)	24 (28.89)	150 (25.11)
Gender			
Men	272 (55.79)	59 (49.03)	331 (54.44)
Women	216 (44.21)	61 (50.97)	277 (45.56)
Ethnicity			
Pakistani	223 (45.63)	61 (50.78)	284 (46.65)
Bangladeshi	265 (54.37)	59 (49.22)	324 (53.35)
Employment			
Employed	247 (50.45)	59 (49.48)	306 (50.26)
Unemployed	97 (19.98)	30 (24.94)	127 (20.97)
Retired/ill/housewife	144 (29.57)	31 (25.57)	175 (28.78)
Education			
No qualifications	93 (19.13)	26 (21.63)	119 (19.62)
< Bachelor degree	239 (49.24)	65 (55.31)	304 (50.43)
Bachelor, Master, PhD	153 (31.63)	27 (23.06)	180 (29.95)
Income ^b			
< £5000	75 (23.12)	9 (15.01)	84 (21.86)
£5000–24 999	199 (60.96)	34 (58.05)	234 (60.51)
£25 000–49 999	31 (9.45)	12 (19.91)	43 (11.07)
> £50 000	21 (6.48)	4 (7.03)	25 (6.56)

a. n = 596.
b. n = 386.

Table 2 Social and health characteristics by sympathies for violent protest and terrorism (weighted)

	n (%)		
	Groups 1 and 2 (n = 488)	Group 3, % (n = 120)	All, % (n = 608)
Life events			
Serious illness, injury or assault to a relative	24 (4.89)	1 (1.10)	25 (4.14)
Death of a partner, spouse, parent or child	16 (3.25)	1 (0.42)	17 (2.69)
Death of a close friend or relative	57 (11.65)	4 (3.64)	61 (10.07)
Separation due to marital differences	4 (0.86)	0 (0.00)	4 (0.69)
The end of a regular and steady relationship	23 (4.69)	3 (2.66)	26 (4.29)
A serious problem with a close friend, neighbour or relative	58 (11.81)	6 (4.88)	64 (10.44)
Unemployment or seeking work unsuccessfully for 1 month or more	28 (5.77)	10 (8.51)	38 (6.32)
Lost a job (fired, asked to leave)	18 (3.68)	1 (1.05)	19 (3.16)
Major financial crisis	24 (5.03)	8 (6.38)	32 (5.30)
Problem with the police or a court appearance	4 (0.81)	5 (3.81)	9 (1.41)
Something valuable to you was lost or stolen	23 (4.81)	0 (0.02)	23 (3.86)
Another major event that you found stressful not listed above	35 (7.08)	1 (0.88)	36 (5.85)
Political engagement			
Voted in the last local council election	316 (64.92)	60 (49.48)	376 (61.86)
Discussed politics or political news with someone else	122 (24.98)	23 (19.34)	145 (23.86)
Signed a petition	125 (25.69)	17 (13.79)	142 (23.33)
Donated money to a charity or campaigning organisation	222 (45.56)	26 (21.97)	248 (40.89)
Paid a membership fee to a charity or campaigning organisation	30 (6.06)	3 (2.73)	33 (5.40)
Done voluntary work	104 (21.30)	10 (8.66)	114 (18.80)
Boycotted certain products for political, ethical or environmental reasons	24 (4.84)	2 (2.07)	26 (4.29)
Boycotted certain products for religious reasons	43 (8.95)	1 (0.40)	44 (7.26)
Expressed my political opinions online	16 (3.41)	5 (3.98)	21 (3.52)
Been to any political meeting	10 (2.05)	0 (0.13)	10 (1.67)
Donated money or paid a membership fee to a political party	17 (3.46)	4 (3.12)	21 (3.39)
Take part in a demonstration, picket or march	19 (4.00)	1 (0.71)	20 (3.35)
Depression^a			
Patient Health Questionnaire score <5	339 (80.93)	56 (62.09)	395 (77.57)
Patient Health Questionnaire score ≥5	80 (19.07)	34 (37.91)	114 (22.43)

a. n = 509.

Table 3 Simple regression models: associations between sympathies for violent protest and terrorism and depression with social and health variables (weighted)^a

	Sympathies for violent protest and terrorism		Probable clinical depression	
	Odds ratio (OR) (95% CI)	P	OR (95% CI)	P
Life events (no reference group)				
Serious illness, injury or assault to a relative	0.22 (0.04–1.26)	0.089	1.00 (0.38–2.64)	0.999
Death of a partner, spouse, parent or child	0.12 (0.01–2.08)	0.147	13.15 (3.83–45.17)	<0.001
Death of a close friend or relative ^b	0.29 (0.11–0.77)	0.014	2.16 (1.18–3.94)	0.012
Separation due to marital differences	–	–	0.39 (0.01–14.89)	0.611
The end of a regular and steady relationship	0.56 (0.17–1.82)	0.332	0.64 (0.16–2.58)	0.532
A serious problem with a close friend, neighbour or relative	0.38 (0.16–0.92)	0.031	1.39 (0.73–2.65)	0.311
Unemployment or seeking work unsuccessfully for 1 month or more	1.52 (0.72–3.20)	0.271	1.07 (0.45–2.54)	0.876
Lost a job (fired, asked to leave)	0.28 (0.05–1.71)	0.167	4.69 (1.70–12.95)	0.003
Major financial crisis	1.29 (0.56–2.97)	0.555	2.68 (1.12–6.44)	0.027
Problem with the police or a court appearance	4.84 (1.24–18.86)	0.023	5.15 (0.81–32.91)	0.084
Something valuable to you was lost or stolen	–	–	1.79 (0.72–4.46)	0.209
Another major event that you found stressful not listed above ^b	0.12 (0.02–0.81)	0.030	4.72 (2.00–11.12)	<0.001
Political engagement (no reference group)				
Voted in the last local council election	0.53 (0.35–0.79)	0.002	1.11 (0.72–1.72)	0.625
Discussed politics or political news with someone else	0.72 (0.44–1.18)	0.195	0.52 (0.31–0.89)	0.016
Signed a petition ^b	0.46 (0.27–0.81)	0.007	1.59 (1.00–2.52)	0.048
Donated money to a charity or campaigning organisation	0.34 (0.21–0.54)	<0.001	0.67 (0.44–1.04)	0.076
Paid a membership fee to a charity or campaigning organisation	0.43 (0.14–1.38)	0.158	0.86 (0.34–2.17)	0.750
Done voluntary work	0.35 (0.18–0.69)	0.002	1.46 (0.89–2.40)	0.131
Boycotted certain products for political, ethical or environmental reasons	0.42 (0.11–1.56)	0.193	2.17 (0.92–5.10)	0.076
Boycotted certain products for religious reasons	0.04 (0.00–0.70)	0.028	0.82 (0.37–1.83)	0.621
Expressed my political opinions online	1.17 (0.42–3.31)	0.761	3.40 (1.31–8.88)	0.012
Been to any political meeting	0.06 (0.00–8.77)	0.274	8.07 (2.07–31.42)	0.003
Donated money or paid a membership fee to a political party	0.90 (0.29–2.80)	0.852	1.85 (0.68–5.03)	0.225
Take part in a demonstration, picket or march	0.17 (0.02–1.51)	0.112	1.38 (0.50–3.85)	0.537
Depression (Patient Health Questionnaire score <5 as reference group)				
Patient Health Questionnaire score ≥5	2.59 (1.59–4.23)	<0.001	–	–

a. See online Table DS1 for odds ratios for demographic characteristics.

b. Potential mediating effect of depression as associated with life events and political engagement and sympathies for violent protest and terrorism carried forward for mediation analyses, see Fig. 2.

or made a court appearance were more likely to report SVPT. As predicted, people who voted in the last election, signed a petition, donated money to charity, provided voluntary work or boycotted products for religious reasons were less likely to report SVPT. See online Table DS1 for odds ratios for demographic characteristics.

Multivariable analyses

Figure 1 shows the relationship between SVPT, specific life events and acts of political engagement, with one model for each of the items. On the whole, the effects of life events and political engagement on SVPT were independent of probable clinical depression. Adjusted analyses (Fig. 1) suggest that death of a close friend (odds ratio (OR) = 0.24, 95% CI 0.07–0.74, $P = 0.014$), signing a petition (OR = 0.32, 95% CI 0.15–0.66, $P = 0.002$), donating money to a charity (OR = 0.52, 95% CI 0.3–0.9, $P = 0.018$), voluntary work (OR = 0.31, 95% CI 0.14–0.66, $P = 0.003$) and boycotting religious products (OR = 0.04, 95% CI 0–0.78, $P = 0.033$) are associated with a lower risk of SVPT. Another variable, major life events (not specifically defined by the questionnaire), falls just short of a significantly lower risk (OR = 0.01, 95% CI 0–1.05, $P = 0.053$), whereas contact with the police and courts falls just short of a significantly higher risk of SVPT (OR = 6.49, 95% CI 0.96–43.85, $P = 0.055$).

The aggregation of cluster groups in the analysis was driven by the association of a higher risk of depressive symptoms in group 3 compared with groups 1 and 2 combined. However, in order to aid interpretation of the findings, univariable analyses of life event and political engagement items by specific cluster groups were also undertaken. These compared group 1 (condemning) with group 2 (intermediate as reference) and group 3 (sympathetic). Boycotting religious products, signing a petition and voluntary work were associated with (lower risk) membership of group 3 compared with group 2 as the reference, but these items were also associated with a lower risk of membership of group 1 compared with group

2, suggesting that those who expressed most sympathies and most condemnation had lower levels of political engagement. In contrast, voting in the last council elections, donating money to a charity, and all the life event items were not associated with membership of group 1 compared with group 2, but showed an association (lower risk) with membership of group 3 compared with group 2.

Mediation analyses

Three items were potential mediators, showing significant associations with both probable clinical depression and SVPT (Table 3): death of a close friend or relative, another major life event, and signing a petition. Expressing a problem with the police or criminal justice agencies was strongly associated with SVPT, but less so with probable clinical depression. As a result, the effects were unlikely to be mediated by depression and this possible association was not considered further.

In the absence of depression, experiencing the death of a close friend or relative, another major life event, and having signed a petition are all associated with a lower direct risk of SVPT. Yet, when the three variables are accompanied by symptoms of depression, the risk of SVPT increases by a small amount. We found a lower overall risk of SVPT associated with political engagement and life events even in the presence of depressive symptoms (Fig. 2).

Discussion

Pathways to SVPT

Specific life events are strongly associated with a lower risk of SVPT, whereas the effects are mostly independent of depression. Only contact with police or the courts carried a higher risk of SVPT, perhaps explained by past criminality or a heightened sense of injustice, leading to grievance and support for extremism. Yet,




Fig. 1 Multivariable analyses: association between sympathies for violent protest and terrorism and demographic, social and health variables (odds ratios, weighted).

Logistic regression model for each life event or political engagement action in separate models (adjusted for age, gender, employment status, education level, depression; weighted). Most sympathetic group ($n = 92$) compared with least sympathetic and intermediary groups ($n = 516$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. No-Ref: 'no' responses used as reference.




Fig. 2 Mediation analyses for the role of depressive symptoms in explaining the relationship between life events, political engagement and sympathies for violent protest and terrorism: logistic regression showing direct and indirect pathways.

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. OR, odds ratio.

relatively few participants reported involvement with the police and courts and the variable was not strongly associated with probable clinical depression. Political engagement was also associated with a lower risk of SVPT, which is encouraging given the current UK emphasis on policies to promote political literacy and civic participation. These life events and acts of political engagement should not be used as markers of SVPT, as some types of political engagement (boycotting religious products, signing a petition and voluntary work) do not distinguish those at high and low risk of SVPT.

An association between adverse life events and depression is well established, invoking feelings of entrapment or humiliation,^{22,32,33} underpinned by biological mechanisms of heightened amygdala activity and altered brain connectivity.^{34,35} We found depressive symptoms are associated with SVPT. However, the finding that life events appear to reduce the risk of SVPT is surprising, as adversity and inequality are often proposed to explain extreme beliefs and violent behaviour.⁷ It is possible that losing a friend or relative might teach about the value of life and what it means to others to suffer a bereavement or loss, thereby deterring SVPT. Alternatively, adverse life events may cause people to draw on pre-existing social networks as a means of emotional support, creating opportunities to resolve disaffection and isolation. Yet, *post hoc* adjustments to the regression models for social support and the proportion of people from the same ethnic group made no difference to the estimates. This suggests that social support does not explain the effect, although there may be residual unmeasured or unknown influences.

Depression and violence

A recent systematic review suggested that depression predisposes participants to later offending.²¹ Depression is also associated with impulsivity and suicidal behaviour, and these in turn are associated with risk of violence more generally.^{36,37} In a previous paper, we found that the effect of depressive symptoms on SVPT is sustained when the analysis is re-run without the suicide item from the PHQ.²⁰ This suggests that the association between SVPT and depressive symptoms is not because of suicidal thinking.

Alternatively, depressive symptoms may serve as a proxy for a number of other social concerns and psychiatric disorders.^{33,38} Further research into these possibilities is needed. Preventing depressive responses to adverse life events and poor political engagement (or poor civic participation) may marginally reduce the risk of SVPT, but our findings suggest that promoting political engagement and social connectedness are more likely to have a larger impact.

Criminal justice system contact

The association between problems with the police or courts and SVPT suggests a subsample who have offended or come to the attention of law enforcement agencies. Violent offending linked with early exposure to adversity, such as material disadvantage and harsh or absent parenting in childhood, produces so-called 'early starters' who use substances, join gangs and offend.^{39,40} However, such influences have not been reported among the families of the recent perpetrators of terrorist attacks in the

European Union and North America, where young men and women involved in terrorist actions appear to fall into the group called 'late starters'; that is, they are relatively high functioning and offend after having encountered political ideologies or developed grievances or, less frequently, become violent because of developing mental illness.

Strengths and limitations

SVPT do not measure actual violence or terrorist offending, but their importance lies in the finding that such sympathies can create or accentuate vulnerability to persuasion and the adoption of the narratives of extremist groups.¹⁰ In recognition of the importance of cognitive rather than behavioural violence,⁹ recent definitions of radicalisation include attitudes towards democracy, British values, and respect for the law and human dignity. Studies of terrorist offending and the emergence of extreme beliefs are important but ethically challenging, given the dilemma and risks of observing behaviour of increasing radicalisation. Furthermore, levels of support for terrorism fluctuate and are influenced by high-profile events and selection bias in sampling. For example, after the Charlie Hebdo attacks in France, 27% of a sample of British Muslims endorsed an item showing sympathy for the motives behind the attacks.⁴¹

There are other reasons for trying to reduce SVPT. Sympathisers may serve as a pool for sustaining infectious ideas that, even if in the minority, polarise whole populations.^{42,43} Radical ideas may be transformed into a practical threat if those who are sympathetic offer resources to terrorist groups.⁴³ Reductions of the population prevalence of SVPT may be effected by encouraging political engagement and social inclusion in order to shift public opinion. Achieving this in young people and public institutions accords with the UK Terrorism Act of 2000, which mandates safeguarding duties for all citizens. Although this study suggests that depression may be a key pathway; more needs to be discovered about specific mechanisms of developing extremist ideas, preferably using longitudinal designs. Although cross-sectional data are not ideal for studying partial mediation,⁴⁴ the bias serves to overestimate apparent effects. Our study found little support for mediation and, as longitudinal studies would reveal more conservative or no effect, these would be consistent with our findings.

Given the global importance of terrorism and the relative lack of research into the process of radicalisation, further studies are needed of other populations, and replication of the existing methods in different country contexts. Alternative sampling strategies, such as probability sampling, may be useful but would be expensive as large numbers of participants would be needed to complete the preliminary consent and screening procedure to assess their suitability for the study. For such a sensitive topic, this would raise ethical questions if the same research questions can be answered using quota samples. We did not assess personality disorders, which may be important correlates of offending behaviour, especially antisocial personality disorder. However, the notion of measuring personality across cultures is contested and diagnostic thresholds may differ across cultural groups.^{45,46} Future research will need to grapple with these methodological dilemmas.

Kamaldeep Bhui, BSc, MBBS, MSc, MD, FRCPsych, **Maria Joao Silva**, MSc, **Raluca A. Topciu**, MD, MA, PhD, Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Queen Mary University of London, London; **Edgar Jones**, MA, PhD, DPhil, King's Centre for Military Health Research, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

Correspondence: Kamaldeep Bhui, Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Charterhouse Square, London EC1M 6BQ, UK. Email: k.s.bhui@qmul.ac.uk

First received 18 Mar 2016, final revision 19 Jun 2016, accepted 20 Jun 2016

Funding

The study received no grant funding. The data collection was commissioned from Ipsos MORI. The statistical analyses were part funded by Careif, www.careif.org, an international mental health charity. Careif had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Victoroff J. The mind of the terrorist: a review and critique of psychological approaches. *J Confl Resolut* 2005; **49**: 39.
- Lord Alderdice. The individual, the group and the psychology of terrorism. *Int Rev Psychiatry* 2007; **19**: 201–9.
- Stevens MJ. What is terrorism and can psychology do anything to prevent it? *Behav Sci Law* 2005; **23**: 507–26.
- Corner E, Gill P. A false dichotomy? Mental illness and lone actor terrorism. *Law Hum Behav* 2014; **39**: 11.
- Gill P, Horgan J, Deckert P. Bombing alone: tracing the motivations and antecedent behaviors of lone-actor terrorists. *J Forensic Sci* 2014; **59**: 425–35.
- McCauley C, Moskalenko S, Van Son B. Characteristics of lone-wolf violent offenders: a comparison of assassins and school attackers. *Perspect Terrorism* 2013; **7**: 4–24.
- Home Affairs Committee. *The Roots of Violent Radicalisation*. House of Commons, 2012.
- Christmann K. *Preventing Religious Radicalisation and Violent Extremism*. Youth Justice Board, 2012.
- Horgan J. *Walking Away from Terrorism: Accounts of Disengagement from Radical and Extremist Movements*. Routledge, 2009.
- Silber M, Bhatt A. *Radicalization in the West: the Homegrown Threat*. New York City Police Department, 2007.
- HM Government. *Prevent Strategy*. TSO, 2011.
- Tennant C. Life events, stress and depression: a review of recent findings. *Aust NZ J Psychiatry* 2002; **36**: 173–82.
- Viswanath K, Randolph Steele W, Finnegan JR, Jr. Social capital and health: civic engagement, community size, and recall of health messages. *Am J Public Health* 2006; **96**: 1456–61.
- Gourion D. Events of life and links with severe depression at different ages [in French]. *L'Encephale* 2009; **35** (suppl 7): S250–6.
- Gluckman PD, Hanson MA, Beedle AS. Early life events and their consequences for later disease: a life history and evolutionary perspective. *Am J Hum Biol* 2007; **19**: 1–19.
- Sousa CA. Political violence, collective functioning and health: a review of the literature. *Med Confl Surv* 2013; **29**: 169–97.
- Canetti D, Hall BJ, Greene T, Kane JC, Hobfoll SE. Improving mental health is key to reduce violence in Israel and Gaza. *Lancet* 2014; **384**: 493–4.
- Ghosh P. *Violent Radicalisation and Recruitment to Terrorism: Perspectives of Wellbeing and Social Cohesion of Citizens of Muslim Heritage*. Scientific Research Publishing, 2013.
- Bhui K, Warfa N, Jones E. Is violent radicalisation associated with poverty, migration, poor self-reported health and common mental disorders? *PLoS ONE* 2014; **9**: e90718.
- Bhui K, Everitt B, Jones E. Might depression, psychosocial adversity, and limited social assets explain vulnerability to and resistance against violent radicalisation? *PLoS ONE* 2014; **9**: e105918.
- Fazel S, Wolf A, Chang Z, Larsson H, Goodwin GM, Lichtenstein P. Depression and violence: a Swedish population study. *Lancet Psychiatry* 2015; **2**: 224–32.
- Brown GW. Genetic and population perspectives on life events and depression. *Soc Psychiatry Psychiatr Epidemiol* 1998; **33**: 363–72.
- Targosz S, Bebbington P, Lewis G, Brugha T, Jenkins R, Farrell M, et al. Lone mothers, social exclusion and depression. *Psychol Med* 2003; **33**: 715–22.
- Jobst A, Sabass L, Palagyi A, Bauriedl-Schmidt C, Mauer MC, Sarubin N, et al. Effects of social exclusion on emotions and oxytocin and cortisol levels in patients with chronic depression. *J Psychiatr Res* 2015; **60**: 170–7.
- Rubin GJ, Amlot R, Page L, Wessely S. Methodological challenges in assessing general population reactions in the immediate aftermath of a terrorist attack. *Int J Methods Psychiatr Res* 2008; **17** (suppl 2): S29–35.
- Everitt B, Landau S, Leese M, Stahl M. *Cluster Analysis*. John Wiley & Sons Ltd, 2011.
- Banfield JD, Raftery AE. Model-based Gaussian and non-Gaussian clustering. *Biometrics* 1993; **49**: 803–21.

- 28 Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med* 1985; **15**: 189–94.
- 29 Department for Communities and Local Government. Local Government and Communities Citizenship Survey. *SN7111-Citizenship Survey 2010/11*. Department for Communities and Local Government, IPSOS Mori, 2012.
- 30 Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; **16**: 606–13.
- 31 MacKinnon DP, Fairchild AJ, Fritz MS. Mediation analysis. *Ann Rev Psychol* 2007; **58**: 593.
- 32 Parker G, Paterson A, Hadzi-Pavlovic D. Emotional response patterns of depression, grief, sadness and stress to differing life events: a quantitative analysis. *J Affect Disord* 2015; **175**: 229–32.
- 33 Mumford DB, Nazir M, Jilani FU, Baig IY. Stress and psychiatric disorder in the Hindu Kush: a community survey of mountain villages in Chitral, Pakistan. *Br J Psychiatry* 1996; **168**: 299–307.
- 34 Anand A, Li Y, Wang Y, Wu J, Gao S, Bukhari L, et al. Activity and connectivity of brain mood regulating circuit in depression: a functional magnetic resonance study. *Biol Psychiatry* 2005; **57**: 1079–88.
- 35 Swartz JR, Williamson DE, Hariri AR. Developmental change in amygdala reactivity during adolescence: effects of family history of depression and stressful life events. *Am J Psychiatry* 2015; **172**: 276–83.
- 36 Witt K, Hawton K, Fazel S. The relationship between suicide and violence in schizophrenia: analysis of the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) dataset. *Schizophr Res* 2014; **154**: 61–7.
- 37 Apter A, Plutchik R, van Praag HM. Anxiety, impulsivity and depressed mood in relation to suicidal and violent behavior. *Acta Psychiatr Scand* 1993; **87**: 1–5.
- 38 Bhui K, Stansfeld S, McKenzie K, Karlsen S, Nazroo J, Weich S. Racial/ethnic discrimination and common mental disorders among workers: findings from the EMPIRIC Study of Ethnic Minority Groups in the United Kingdom. *Am J Public Health* 2005; **95**: 496–501.
- 39 Simpson AI, Grimbos T, Chan C, Penney SR. Developmental typologies of serious mental illness and violence: evidence from a forensic psychiatric setting. *Aust NZ J Psychiatry* 2015; **49**: 1048–59.
- 40 Ferguson CJ, San Miguel C, Hartley RD. A multivariate analysis of youth violence and aggression: the influence of family, peers, depression, and media violence. *J Pediatr* 2009; **155**: 904–8.e3.
- 41 Hodges D. Over a quarter of British Muslims have sympathy for the Charlie Hebdo terrorists. That is far too many. *Telegraph* 2015; Feb 25 (<http://www.telegraph.co.uk/news/religion/11434695/Over-a-quarter-of-British-Muslims-have-sympathy-for-the-Charlie-Hebdo-terrorists.-That-is-far-too-many.html>).
- 42 Deffuant G, Amblard F, Weisbuch H, Fauris T. How can extremism prevail? A study based on the relative agreement interaction model. *J Artif Soc Soc Simul* 2002; **5**: 1 (<http://jasss.soc.surrey.ac.uk/5/4/1.html>).
- 43 Johnson D, Madin J. Population models and counterinsurgency strategies. In: *Natural Security* (eds R Sagarin, T Taylor): 159–85. London University of California, 2008.
- 44 Maxwell SE, Cole DA. Bias in cross-sectional analyses of longitudinal mediation. *Psychol Methods* 2007; **12**: 23–44.
- 45 McGilloway A, Hall RE, Lee T, Bhui KS. A systematic review of personality disorder, race and ethnicity: prevalence, aetiology and treatment. *BMC Psychiatry* 2010; **10**: 33.
- 46 Ryder AG, Sunohara M, Kirmayer LJ. Culture and personality disorder: from a fragmented literature to a contextually grounded alternative. *Curr Opin Psychiatry* 2015; **28**: 40–5.



psychiatry in literature

Natasha Rostova: a spiritual recovery from depression in *War and Peace*

Hannah Marcarian, Paul O. Wilkinson

War and Peace was written at a time when the term 'depression' was not used, although the symptoms were recognised and referred to as 'melancholy'. It is believed that Tolstoy himself had suffered from depression, and in this work he is able to powerfully convey its symptoms through Natasha Rostova. Interestingly, her father develops depression later in the book, reflecting the frequent presence of a family history.

When Natasha is well, she is charming, lively and spontaneous, and enjoys singing. She is attractive and has many suitors, becoming engaged to Prince Andrew. When Prince Andrew leaves to go travelling, womaniser Anatole seduces Natasha. She makes plans to elope with him but her family foil these and shame Natasha for her actions. Her initial reaction is one within the normal range for someone suffering an adverse life event: she does not move from the sofa, sobbing that everyone despises her, and does not sleep that night. Following the news that Anatole is in fact already married, Natasha feels foolish and humiliated. She attempts suicide by poisoning herself with arsenic, but she quickly becomes frightened and wakes her cousin, who administers an antidote. Again, we sometimes see suicide attempts in adolescents with love life troubles that they soon regret and seek help for.

Natasha's condition worsens over the following months. She is permanently low in spirits, cannot eat or sleep and grows thinner. She avoids any activities that previously brought her pleasure, such as balls and concerts as well as singing and laughing, as these feel like 'blasphemy in face of her sorrow'. She rarely leaves the house and only feels comfortable with her brother Petya and slight happiness if close friend Pierre visits. Natasha feels that there is no future for her and no joy in life; she continues to feel guilty. She thus meets ICD and DSM criteria for depression, with five depressive symptoms lasting longer than 2 weeks and associated functional impairment. Anger from her family is replaced by them medicalising her problems and paying doctors to try to treat her 'illness'. In 1812, when events described in the book take place, depression was not an identified illness and there were no effective treatments. Doctors tried several medications in order to appear to be helping and although Natasha knows they will not work, she seems to appreciate the effort. Her family recognising her as being ill and not apportioning blame, but trying to treat her, seem like helpful and supportive reactions.

Natasha recovers when a friend encourages her to attend church regularly. Praying for forgiveness satisfies her spiritual discomfort after doing wrong. She is also given a sense of routine and purpose, the basis of activity scheduling therapy for depression. Natasha obtains peace and happiness through her sense of forgiveness from and closeness to God, but the doctors think that their treatments have made her get better.

As 21st-century doctors, we must not assume it is always our treatments that lead to our patients' recovery – like for Natasha, this may in fact be due to changes in their social situation or their spiritual needs being met. This also illustrates the importance of thinking about our patients holistically. We teach students and trainees the importance of considering (and trying to address) physical, psychological and social aetiology factors. We should also think about spiritual issues that may be contributing to illness and encourage our patients to seek expert help, as addressing these problems may be crucial for recovery.