September 1961

Vol. LXXV No. 9

The Journal of Laryngology and Otology

EDITED BY WALTER HOWARTH ASSISTANT EDITOR

G. H. BATEMAN

Contents

TREATMENT OF INTRINSIC CARCINOMA OF THE LARYNX RECENT Advances in the Treatment of Intrinsic

CINOMA OF THE LARYNX BY IMPLANTATION OF RADIUM OR TANTALUM WIRE

MOBILIZATION OF A "FIXED " MALLEUS FOLLOWING ON HEAD INJURY

Comments on the Term Epipharynx . . . Laryngeal Stridor in Infants and Children Clinical Records—

- A CASE OF TRACHEAL OBSTRUCTION BY AN ENDOTRACHEAL KELOIDAL PAPILLOMA, FOLLOWING TRACHEOSTOMY FOR NEONATAL LARYNGEAL INSUFFICIENCY. EXCISION AFTER FIVE YEARS OF TRACHEOSTOMY LIFE .
- THE CHANGING SCENE IN OTOLARYNGOLOGY: A 21-YEAR SURVEY

PAGET'S DISEASE OF THE ETHMOIDS

Two Cases of Foreign Bodies in the Nasopharynx

Societies Proceedings-

ROYAL SOCIETY OF MEDICINE—SECTION OF LARYNGOLOGY

GENERAL NOTES

London Headley Brothers

109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13

C. P. WILSON

R. S. LEWIS

FRANK ELLIS

P. W. R. M. ALBERTI and J. D. K. DAWES

> W. Doyle Kelly V. E. Negus D. A. Birch

W. STIRK ADAMS

Ronald Macbeth A. P. Fuller

G. M. TANEJA and G. S. KOHLI

Monthly, 10/- net post free

tos://doi.org/10.1017/S0022215100058424 Published online by Cambridge University Press

ADVERTISEMENTS

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

WALTER HOWARTH

ASSISTANT EDITOR G. H. BATEMAN

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of

every reference before submitting a paper for publication.
3. Galley proofs and engraver's proofs of illustrations are sent to the author, Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within s days.
4. Illustration blocks are provided free up to the limit of £ to per article; beyond this authors are expected to pay half the cost. Coloured illustrations will

be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2. 5. Orders for reprints must be sent when returning galley proofs, and for this

purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English,

French, Italian, or German) it will be welcomed. 7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2. 8 The annual subscription is four guineas sterling (U.S.A. \$13) post free,

and is payable in advance.

9. Single copies will be on sale at ros. od. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this Journal may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East roth Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2, England.

O Journal of Laryngology and Otology, 1960

	PAGE
TREATMENT OF INTRINISIC CARCINOMA OF THE LARYNX. C. P. Wilson	
(London)	78 5
RECENT ADVANCES IN THE TREATMENT OF INTRINSIC CANCER OF	
THE LARYNX. R. S. Lewis (London)	794
A New Method of Treatment of Intrinsic Carcinoma of the	
Larynx by Implantation of Radium or Tantalum	
WIRE. Frank Ellis (Oxford)	811
NECROSIS OF THE LENTICULAR PROCESS OF THE INCUS AFTER STAPES	
SURGERY AND ITS TREATMENT, P. W. R. M. ALBERTI AND	
J. D. K. Dawes (Newcastle upon Tyne)	821
MOBILIZATION OF A "FIXED" MALLEUS FOLLOWING ON HEAD INJURY.	
W. Doyle Kelly (Dublin)	826
Comments on the Term Epipharynx, V. E. Negus (London) .	828
LARYNGEAL STRIDOR IN INFANTS AND CHILDREN. D. A. BIRCH	
(London)	833
CLINICAL RECORDS—	
A Case of Tracheal Obstruction by an Endotracheal Keloidal	
Papilloma, etc. W. Stirk Adams (Birmingham)	841
The Changing Scene in Otolaryngology: A 21-Year Survey.	
Ronald Macbeth (Oxford)	852•
Paget's Disease of the Ethmoids. A. P. Fuller (London)	86 0
Two Cases of Foreign Bodies in the Nasopharynx. G. M. Taneja	
and G. S. Kohli (Amritsar, India)	864
Societies' Proceedings-R.S.MSection of Laryngology	867
General Notes	872

WHICH DEAF AID?

Your patient can compare a complete range of aids to hearing at no extra cost.

Mr. D. C. ALLEN, a member of the Society of Hearing Aid Audiologists, will advise your patients impartially and demonstrate a selection of the leading makes of hearing aids without obligation.




FOR YOUR OWN GUIDANCE -

Send now for an attractive, specially prepared Desk Folder which will tell you at a glance "which deaf aid suits which kind of deafness," complete with a reply-paid referral pad. (Over 50 different fittings are listed.)



HEARING COUNSELLORS LTD 53 GEORGE STREET, LONDON, W.I Tel.: WELbeck 8493 Signatories to the Code of Commercial Practice of the Hearing Aid Industry approved by the National Institute for the Deof—Member of the Hearing Aid Manufacturers and Suppliers Association.

ii



MA 8254

Please mention The Journal of Laryngology and Otology when replying to advertisements

First clinical audiometer to provide 100% accurate audiograms



AMPLIVOX MODEL 82

world's most fully-developed clinical audiometer

It is now known that the disc type of earphone, used in conventional audiometers for air conduction testing and the application of masking noise, transfers the sound by bone conduction to the opposite ear at a level only 40 dB below the signal level. This means that up to 50% of audiograms taken at the present time with other audiometers may be invalid.

The Amplivox Model 82 Audiometer provides, for the first time, the means of taking accurate audiograms in cases of severe unilateral deafness through the provision of narrow band masking applied by insert earphone. This new masking technique is of special significance in bone conduction audiometry.

In addition to the exclusive narrow band masking feature, the Model 82 provides loudness balance test facilities, and *all* the other tests essential to proved pure tone audiometry.

AMPLIVOX LTD Medical Acoustic Division 80 New Bond St, London W.1. Tel: HYDe Park 9888



Containing full information on the latest audiometric techniques in air and bone conduction.

Reference: Hearing Hids = refer patients to INGRAMS **Remarks**: they have the widest selection of heaving aids Ingrams in the country and are ······PLEASE TEAR THIS OFF AND KEE completely independent of all makers. They really do give patients impartial advice. They charge only the makers' list prices. nothing extra-Ingrams Hearing Aids Ltd. 2 Shepherd Street, Shepherd Market, London, W.I. phone-Hyde Park 9041


* Members of the Society of Hearing Aid Audiologists. Resident representatives in all parts of the country.

CASE 1.

Fitting HEARING AIDS

in

DIFFICULT CASES



Case 1: LORD E. AGE 75

History: Progressive deterioration in hearing over many years. Presbyacusis. Tried several hearing aids without success.

Hearing aid fitting: Amplivox Model AV, 9W wide range phone, amplifier with reduced low tone response. Maximum AVC setting.

Maximum Intelligibility: With hearing aid in left ear-90%

Remarks: Has expressed great satisfaction with clarity of tone and the substantial assistance obtained in all situations.

In measuring the performance of a hearing aid, no yardstick is more indicative than the results achieved in 'difficult cases'. The above case history is only one of the many successes which Amplivox have had over the past quarter-century.

Contributing to this success are the care and thoroughness with which the Amplivox Hearing Advisory Service fit a hearing aid. Air and bone conduction pure tone audiograms, loudness tolerance levels and phonetically balanced word tests are the basis of testing. It is this extreme care in testing and fitting which, coupled with expert knowledge of the instrument characteristics, ensures optimum hearing in all cases.

Amplivox incorporate the world's finest hearing aids in their range, including head-borne and body-worn aids (many incorporating AVC) from the Super 'A', for the profoundly deaf, to the latest all-behind-the-ear model, Secrette. There are permanent Amplivox Hearing


Advisory Centres in principal cities throughout the country, providing a reliable and thorough service for the hard-of-hearing.

For further information and descriptive literature please telephone the local Amplivox Centre or write to:

AMPLIVOX LTD.

80 New Bond Street, London, W.1 Tel: Hyde Park 9888

CENTRES AT: BIRMINGHAM, BOURNEMOUTH, BRISTOL, CARDIFF, EDINBURGH, GLASGOW, HULL, LEEDS, LEICESTER, LIVERPOOL, MANCHESTER, NEWCASTLE, DUBLIN



'OTOSPORIN' clears infected ears

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' brand Drops.

'Otosporin' is an aqueous suspension containing both neomycin and polymyxin **B** sulphate which are together effective against practically all the bacteria found in ear infections. They are unlikely to induce bacterial resistance or skin sensitisation, and neither of them gives rise to cross-resistance or cross-sensitisation to penicillin and other antibiotics. Their effectiveness is visibly enhanced by the hydrocortisone in 'Otosporin'; this, by reducing inflammation, not only relieves pain, but provides greater access for the antibiotics.

'OTOSPORIN¹ drops

issued in bottles of 5 c.c.

