- 4 Hoekstra PJ, Dietrich A, Edwards MJ, Elamin I, Martino D. Environmental factors in Tourette syndrome. *Neurosci Biobehav Rev* 2013; **37**: 1040–9.
- 5 Scharf JM, Yu D, Mathews CA, Neale BM, Stewart SE, Fagerness JA, et al. Genome-wide association study of Tourette's syndrome. *Mol Psychiatry* 2013; **18**: 721–8.
- 6 Plessen KJ, Bansal R, Peterson BS. Imaging evidence for anatomical disturbances and neuroplastic compensation in persons with Tourette syndrome. *J Psychosom Res* 2009; **67**: 559–73.
- 7 Ganos C, Roessner V, Munchau A. The functional anatomy of Gilles de la Tourette syndrome. *Neurosci Biobehav Rev* 2013; **37**: 1050–62.
- 8 Munafò MR, Flint J. How reliable are scientific studies? *Br J Psychiatry* 2010; **197**: 257–8.
- 9 Murad MH, Montori VM. Synthesizing evidence: shifting the focus from individual studies to the body of evidence. *JAMA* 2013; **309**: 2217–8.
- 10 Santer BD, Wigley TM, Taylor KE. The reproducibility of observational estimates of surface and atmospheric temperature change. *Science* 2011; **334**: 1232–3.
- 11 Mathews CA, Scharf JM, Miller LL, Macdonald-Wallis C, Lawlor DA, Ben-Shlomo Y. Association between pre- and perinatal exposures and Tourette syndrome or chronic tic disorder in the ALSPAC cohort. *Br J Psychiatry* 2014; **204**: 40–5.
- 12 Lebel C, Mattson SN, Riley EP, Jones KL, Adnams CM, May PA, et al. A longitudinal study of the long-term consequences of drinking during pregnancy: heavy in utero alcohol exposure disrupts the normal processes of brain development. *J Neurosci* 2012; **32**: 15243–51.
- 13 Lin H, Katsoyich L, Ghebremichael M, Findley DB, Grantz H, Lombroso PJ, et al. Psychosocial stress predicts future symptom severities in children and adolescents with Tourette syndrome and/or obsessive-compulsive disorder. *J Child Psychol Psychiatry* 2007; **48**: 157–66.
- 14 Lin H, Williams KA, Katsoyich L, Findley DB, Grantz H, Lombroso PJ, et al. Streptococcal upper respiratory tract infections and psychosocial stress predict future tic and obsessive-compulsive symptom severity in children and adolescents with Tourette syndrome and obsessive-compulsive disorder. *Biol Psychiatry* 2010; **67**: 684–91.
- 15 Eisenberg L. The social construction of the human brain. *Am J Psychiatry* 1995; **152**: 1563–75.
- 16 Spencer N, Thanh TM, Louise S. Low income/socio-economic status in early childhood and physical health in later childhood/adolescence: a systematic review. *Matern Child Health J* 2013; **17**: 424–31.
- 17 Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *JAMA* 2007; **298**: 1685–7.

100
words

Gilles de la Tourette syndrome

Andrea E. Cavanna

Gilles de la Tourette syndrome was first described by Parisian doctor Georges Albert Édouard Brutus Gilles de la Tourette in 1885. Originally considered a rare medical curiosity, the syndrome of multiple tics is now regarded as a fairly common neurodevelopmental condition, affecting up to 1% of school-age children. Gilles de la Tourette described the clinical triad of tics, echolalia and coprolalia. The last is relatively rare and no longer features as a diagnostic criterion, despite the media obsession with obscene manifestations. French colleagues react with horror when they see their syndrome's name shortened to Tourette syndrome, even in 100-word texts!

The British Journal of Psychiatry (2014)
204, 8. doi: 10.1192/bjp.bp.112.121467