



An Extension of the Dirichlet Density for Sets of Gaussian Integers

L. C. Rêgo and R. J. Cintra

Abstract. Several measures for the density of sets of integers have been proposed, such as the asymptotic density, the Schnirelmann density, and the Dirichlet density. There has been some work in the literature on extending some of these concepts of density to higher dimensional sets of integers. In this work, we propose an extension of the Dirichlet density for sets of Gaussian integers and investigate some of its properties.

1 Introduction

Several measures for the density of sets of integers have been discussed in the literature [23], [6], [9], [10], [11], [8], [2]. Presumably the most employed of such measures is the asymptotic density, also referred to as natural density [20], [23]. For a given set of integers A , its asymptotic density is expressed by

$$d(A) = \lim_{n \rightarrow \infty} \frac{\|A \cap \{1, 2, 3, \dots, n\}\|}{n},$$

provided that such a limit does exist. The symbol $\|\cdot\|$ returns the cardinality of its argument.

In [6], Bell and Burris provide a good exposition on the Dirichlet density, which is defined as follows.

Definition 1.1 The Dirichlet density of a subset A of the positive integers is given by

$$\partial(A) \triangleq \lim_{s \downarrow 1} \frac{\sum_{n \in A} \frac{1}{n^s}}{\zeta(s)},$$

if the limit does exist, for real $s > 1$. The quantity $\zeta(\cdot)$ denotes the Riemann zeta function [16].

If the asymptotic density is well defined, then the Dirichlet density does also exist and assumes the same value [5, p. 10]. Since the converse is not always true, the Dirichlet density is a more encompassing tool when compared to the asymptotic

Received by the editors April 8, 2010; revised July 1, 2010.

Published electronically August 3, 2011.

The second author acknowledges partial financial support from the Department of Foreign Affairs and International Trade (DFAIT), Canada, and the *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq), Brazil. Part of the second author's work was done during his sabbatical at the University of Calgary, Calgary, AB, Canada.

AMS subject classification: 11B05, 11M99, 11N99.

Keywords: Gaussian integers, Dirichlet density.

density [5, p. 11]. Dirichlet density also admits lower and upper versions, which have been explored along with other densities to characterize primitive sets [1], [3], [4].

Gaussian integers are simply complex numbers of the form $m + in$, where m and n are integers. Despite the considerable amount of development addressing densities for sets of positive integers [15], densities for sets of Gaussian integers appear to be an overlooked topic. However, a seminal paper by Cheo [7] investigated the question, suggesting an extension of the Schnirelmann density [9], [10]. Such extended definition applies to subsets of the nonzero Gaussian integers inclusively confined in the first quadrant of the complex plane.

Generalizations of Schnirelmann density for the n -dimensional case were proposed in [18], [19]. Additionally, a modified Schnirelmann density was introduced in [22] and was generalized in [13] years later. In a comparable venue, Freedman [12], [14] advanced the concept of asymptotic density to higher dimensions.

In this context, the aim of the present work is to advance a method for evaluating the density of sets of Gaussian integers. To address this problem, a density based on Dirichlet generating functions is proposed.

For ease of notation, henceforth we identify a Gaussian integer $m + in$ with the pair of integers (m, n) . All considered Gaussian integers and their sets are defined in \mathbb{P}^2 , where \mathbb{P} is the set of strictly positive integers.

2 Definition and General Properties

The Gaussian integers can be realized as points over a square lattice in the complex plane. The square lattice is composed by an infinite array of Gaussian integers, set up in rows and columns. In addition, each lattice row or column can furnish sets of integers according to the following constructions: $A_{*,n} = \{m \in \mathbb{P} : (m, n) \in A\}$ and $A_{m,*} = \{n \in \mathbb{P} : (m, n) \in A\}$.

Our goal is to investigate the properties of the following density for Gaussian integers, which we show to be a generalization of the Dirichlet density for sets of integers.

Definition 2.1 Let A be a set of Gaussian integers. Admit $I_{A_{*,n}}(\cdot)$ and $I_{A_{m,*}}(\cdot)$ to be the indicator functions of the sets $A_{*,n}$ and $A_{m,*}$, for $m, n \in \mathbb{P}$, respectively. The proposed density for A is given by

$$\text{dens}(A) \triangleq \lim_{s \downarrow 1} \frac{1}{\zeta^2(s)} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{I_{A_{*,n}}(m) I_{A_{m,*}}(n)}{(mn)^s},$$

provided that the limit exists.

From now on, we only consider sets whose referred densities are well-defined, *i.e.*, the implied limits exist. Thus, we restrict ourselves to indicating in every instance that the results are valid only when the discussed limits exist. On account of the proposed definition, a series of consequences is listed below.

Proposition 2.2 Let A and B be two sets of Gaussian integers. The following assertions hold true:

- (i) $\text{dens}(A) \geq 0$,
- (ii) $\text{dens}(\mathbb{P}^2) = 1$,
- (iii) if $A \cap B = \emptyset$, then $\text{dens}(A \cup B) = \text{dens}(A) + \text{dens}(B)$,
- (iv) $\text{dens}(\emptyset) = 0$,
- (v) $\text{dens}(B - A) = \text{dens}(B) - \text{dens}(A \cap B)$, where $B - A$ is the relative complement of A in B .
- (vi) $\text{dens}(A^c) = 1 - \text{dens}(A)$, where A^c is the complement of A ,
- (vii) if $A \subset B$, then $\text{dens}(A) \leq \text{dens}(B)$,
- (viii) $\text{dens}(A \cup B) = \text{dens}(A) + \text{dens}(B) - \text{dens}(A \cap B)$.

Proof The proof follows directly from the definition. ■

The first three properties stated in the previous proposition are the same conditions that form an axiomatic definition of a probability measure, except for the σ -additivity axiom.

Proposition 2.3 (Cartesian Product) *Let A and B be two subsets of \mathbb{P} . Then the density of the Cartesian product $A \times B$ satisfies*

$$\text{dens}(A \times B) = \partial(A) \partial(B).$$

Proof We have that

$$\begin{aligned} \text{dens}(A \times B) &= \lim_{s \downarrow 1} \frac{\sum_{(m,n) \in A \times B} \frac{1}{(mn)^s}}{\zeta^2(s)} \\ &= \lim_{s \downarrow 1} \frac{\sum_{m \in A} \frac{1}{m^s} \sum_{n \in B} \frac{1}{n^s}}{\zeta^2(s)} \\ &= \partial(A) \partial(B). \end{aligned} \quad \blacksquare$$

Corollary 2.4 (Dirichlet Density) *For a set A of positive integers, $\text{dens}(A \times \mathbb{P}) = \partial(A)$.*

Proof This result is a direct consequence of the fact that $\partial(\mathbb{P}) = 1$ [15]. ■

Given any set A of Gaussian integers, let the horizontal and vertical axis sections be denoted by $\text{sup}_h(A) = \bigcup_{n=1}^\infty A_{*,n}$ and $\text{sup}_v(A) = \bigcup_{m=1}^\infty A_{m,*}$, respectively.

Proposition 2.5 *Let A be a set of Gaussian integers. If*

$$\partial(\text{sup}_h(A)) = 0 \quad \text{or} \quad \partial(\text{sup}_v(A)) = 0,$$

then $\text{dens}(A) = 0$.

Proof For instance, assume that $\partial(\text{sup}_h(A)) = 0$. Note that $A \subset \text{sup}_h(A) \times \mathbb{P}$. Due to the monotonicity property, it follows that $\text{dens}(A) \leq \text{dens}(\text{sup}_h(A) \times \mathbb{P})$. Moreover, the property of the density of Cartesian products allows us to write $\text{dens}(A) \leq \partial(\text{sup}_h(A)) \partial(\mathbb{P})$. Applying the hypothesis, the result follows. The proof would be analogous in the case that $\partial(\text{sup}_v(A)) = 0$. ■

Corollary 2.6 (Finite Axis Section) *If a set A of Gaussian integers has a finite axis section, then $\text{dens}(A) = 0$.*

Proof It is enough to observe that any finite set of integers has null Dirichlet density [15]. ■

As a consequence, a finite set of Gaussian integers has null density, since both of its axis sections are finite. In particular, the density of a singleton is zero. On the other hand, nonzero density subsets must have infinite axis sections.

Let

$$U_{m_0, n_0} = \{(m, n) \in \mathbb{P}^2 : m \geq m_0 \text{ and } n \geq n_0\}$$

$$L_{m_0, n_0} = \{(m, n) \in \mathbb{P}^2 : m < m_0 \text{ and } n < n_0\}.$$

The next proposition states that, for density evaluation, the only relevant set elements are those located in the region defined by U_{m_0, n_0} for any choice of m_0 and n_0 .

This means that the “weight” of the set is located on its “tail”. Nevertheless, we need the result of the following lemma first.

Lemma 2.7 *The set U_{m_0, n_0} has unit density.*

Proof Let U_{m_0, n_0}^c be the complement of U_{m_0, n_0} . Therefore, the set \mathbb{P}^2 can be partitioned into $\mathbb{P}^2 = U_{m_0, n_0} \cup U_{m_0, n_0}^c$. Then $\text{dens}(U_{m_0, n_0}) = 1 - \text{dens}(U_{m_0, n_0}^c)$. Notice also that $U_{m_0, n_0}^c = L_{m_0, \infty} \cup L_{\infty, n_0}$. The union property allows us to state that $\text{dens}(U_{m_0, n_0}^c) = \text{dens}(L_{m_0, \infty}) + \text{dens}(L_{\infty, n_0}) - \text{dens}(L_{m_0, n_0})$. Since $L_{m_0, \infty}$, L_{∞, n_0} , and L_{m_0, n_0} have each at least one finite axis section, it follows that $\text{dens}(U_{m_0, n_0}^c) = 0$. ■

Proposition 2.8 (Heavy Tail) *Let A be a set of Gaussian integers. Then for any two given nonnegative integers m_0 and n_0 , we have*

$$\text{dens}(A) = \text{dens}(A \cap U_{m_0, n_0}).$$

Proof Observe that $A = A \cap (U_{m_0, n_0} \cup U_{m_0, n_0}^c) = (A \cap U_{m_0, n_0}) \cup (A \cap U_{m_0, n_0}^c)$. Since we have a partition of A , it follows that $\text{dens}(A) = \text{dens}(A \cap U_{m_0, n_0}) + \text{dens}(A \cap U_{m_0, n_0}^c)$. But, $A \cap U_{m_0, n_0}^c \subset U_{m_0, n_0}^c$, then $\text{dens}(A \cap U_{m_0, n_0}^c) \leq \text{dens}(U_{m_0, n_0}^c) = 0$. ■

Proposition 2.9 (Axis Independence) *If there is a pair (m_0, n_0) such that, for every $m \geq m_0$ and $n \geq n_0$, the functions $I_{A_{m,*}}(n)$ and $I_{A_{*,n}}(m)$ are independent of m and n , respectively, then*

$$\text{dens}(A) = \partial(A_{m_0,*}) \partial(A_{*,n_0}).$$

Proof Because of the assumed independence, we can write $I_{A_{m,*}}(n) = I_{A_{m_0,*}}(n)$ and $I_{A_{*,n}}(m) = I_{A_{*,n_0}}(m)$, for $m \geq m_0$ and $n \geq n_0$, respectively. Thus, for $m \geq m_0$ and $n \geq n_0$, the set A is indistinguishable of $A_{m_0,*} \times A_{*,n_0}$. But the heavy tail property implies that

$$\begin{aligned} \text{dens}(A) &= \text{dens}(A \cap U_{m_0, n_0}) \\ &= \text{dens}((A_{m_0,*} \times A_{*,n_0}) \cap U_{m_0, n_0}) \\ &= \text{dens}(A_{m_0,*} \times A_{*,n_0}) \\ &= \partial(A_{m_0,*}) \partial(A_{*,n_0}). \end{aligned}$$

■

Given a set A of Gaussian integers and a Gaussian integer (m_0, n_0) , let

$$A \oplus (m_0, n_0) \triangleq \{(m + m_0, n + n_0) \mid (m, n) \in A\}.$$

This process is called a translation of A by (m_0, n_0) units [21, p. 49]. Now our goal is to show that the proposed density is translation invariant, *i.e.*, $\text{dens}(A \oplus (m_0, n_0)) = \text{dens}(A)$, $m_0 \geq 0$ and $n_0 \geq 0$. However, the proof that we will supply requires the following lemma.

Lemma 2.10 (Unitary Translation) *Let A be a set of Gaussian integers, such that $\text{dens}(A) > 0$. Then*

$$\text{dens}(A \oplus (1, 0)) = \text{dens}(A \oplus (0, 1)) = \text{dens}(A).$$

Proof It suffices to show that $\text{dens}(A \oplus (1, 0)) = \text{dens}(A)$, being the other case analogous. First, note that since

$$\sum_{(m,n) \in A} \frac{1}{(mn)^s} - \sum_{(m,n) \in A} \frac{1}{((m+1)n)^s} \geq 0,$$

it follows that $\text{dens}(A) - \text{dens}(A \oplus (1, 0)) \geq 0$. Also observe that

$$\frac{s}{m^{s+1}} \geq \int_m^{m+1} \frac{s}{x^{s+1}} dx = \frac{1}{m^s} - \frac{1}{(m+1)^s} \geq 0.$$

Thus, we have that

$$\begin{aligned} \sum_{(m,n) \in A} \frac{1}{(mn)^s} - \sum_{(m,n) \in A} \frac{1}{((m+1)n)^s} &= \sum_{(m,n) \in A} \frac{1}{n^s} \left(\frac{1}{m^s} - \frac{1}{(m+1)^s} \right) \\ &\leq \sum_{(m,n) \in A} \frac{1}{n^s} \frac{s}{m^{s+1}} \\ &\leq \sum_{n \in \mathbb{P}} \frac{1}{n^s} \sum_{m \in \text{sup}_h(A)} \frac{s}{m^{s+1}}. \end{aligned}$$

Dividing both sides by $\zeta^2(s)$ and letting $s \downarrow 1$, since the last series is convergent as $s \downarrow 1$, yields

$$\text{dens}(A) - \text{dens}(A \oplus (1, 0)) \leq 0. \quad \blacksquare$$

Proposition 2.11 (Translation Invariance) *Let A be a set of Gaussian integers. Then*

$$\text{dens}(A \oplus (m, n)) = \text{dens}(A),$$

where m and n are nonnegative integers.

Proof We have already proven that $\text{dens}(A \oplus (1, 0)) = \text{dens}(A \oplus (0, 1)) = \text{dens}(A)$. Therefore, we have that

$$\begin{aligned} \text{dens}(A \oplus (m + 1, n + 1)) &= \text{dens}\left(\left((A \oplus (m, n)) \oplus (1, 0)\right) \oplus (0, 1)\right) \\ &= \text{dens}\left((A \oplus (m, n)) \oplus (1, 0)\right) \\ &= \text{dens}(A \oplus (m, n)) \\ &= \text{dens}(A). \end{aligned} \quad \blacksquare$$

Corollary 2.12 *The proposed density is not σ -additive.*

Proof This result follows directly from Propositions 2.2 and 2.11. ■

Now consider the set operation defined by

$$(a, b) \otimes A \triangleq \{(am, bn) \mid (m, n) \in A\},$$

where (a, b) is a Gaussian integer. This construction can be interpreted as a dilation on the elements of A . The following proposition relates the density of a set of Gaussian integers with the density of its dilated form.

Proposition 2.13 (Dilation) *Let A be a set of Gaussian integers and let (a, b) be any Gaussian integer. Then*

$$\text{dens}((a, b) \otimes A) = \frac{1}{ab} \text{dens}(A).$$

Proof This result follows directly from the definition of the proposed density:

$$\begin{aligned} \text{dens}((a, b) \otimes A) &= \lim_{s \downarrow 1} \frac{\sum_{(m,n) \in A} \frac{1}{(abn)^s}}{\zeta^2(s)} \\ &= \lim_{s \downarrow 1} \frac{\frac{1}{(ab)^s} \sum_{(m,n) \in A} \frac{1}{(mn)^s}}{\zeta^2(s)} \\ &= \frac{1}{ab} \text{dens}(A). \end{aligned} \quad \blacksquare$$

3 Density of Particular Sets

In this section, we evaluate the density of some particular sets of Gaussian integers.

3.1 Cartesian Product of Arithmetic Progressions

Let p be an integer. The set $M_p = \{m \in \mathbb{P} : m \equiv 0 \pmod{p}\}$ constitutes an arithmetic progression with Dirichlet density $\partial(M_p) = 1/p$. Furthermore, the Cartesian product of two arithmetic progressions generates a rectangular lattice denoted by $M_{(p,q)} \triangleq M_p \times M_q$, where p and q are positive integers. Then it follows from Proposition 2.3 that $\text{dens}(M_{(p,q)}) = \partial(M_p) \partial(M_q)$. Let us investigate the density of sets that are intersections of particular Cartesian products of arithmetic progressions.

Proposition 3.1 (Intersection) For any positive integers p, q, s and t , we have that

$$\text{dens}(M_{(p,q)} \cap M_{(s,t)}) = \text{dens}(M_{(\text{lcm}(p,s), \text{lcm}(q,t))}) = \frac{1}{\text{lcm}(p, s) \text{lcm}(q, t)},$$

where $\text{lcm}(\cdot, \cdot)$ denotes the least common multiple of its arguments.

Proof First, note that $M_{(p,q)} = (p, q) \otimes \mathbb{P}^2$. Therefore,

$$\begin{aligned} M_{(p,q)} \cap M_{(s,t)} &= ((p, q) \otimes \mathbb{P}^2) \cap ((s, t) \otimes \mathbb{P}^2) \\ &= (\text{lcm}(p, s), \text{lcm}(q, t)) \otimes \mathbb{P}^2. \end{aligned}$$

Applying $\text{dens}(\cdot)$ on both sides of above equation and invoking the dilation property, we obtain the desired result. ■

Corollary 3.2 Let (m, n) be a Gaussian integer. Admit also that $\text{gcd}(p, s) = 1$ and $\text{gcd}(q, t) = 1$, where $\text{gcd}(\cdot, \cdot)$ returns the greatest common divisor of its arguments. Then

$$\text{dens}(M_{(mp,nq)} \cap M_{(ms,nt)}) = \frac{1}{mn} \text{dens}(M_{(p,q)} \cap M_{(s,t)}).$$

Proof Follows directly from Proposition 3.1. ■

3.2 Sets Delimited by Functions

Let us consider a set of Gaussian integers defined as $C = \{(m, n) \in \mathbb{P}^2 : f(m) \leq n \leq g(m)\}$, where $f(\cdot)$ and $g(\cdot)$ are functions such that $g(m) \geq f(m) \geq 1$ for every integer m . Functions f and g delimit the set C , confining the set elements in between. Figure 1 illustrates a possible configuration for the set C . By definition, the proposed density of C is given by

$$\text{dens}(C) = \lim_{s \downarrow 1} \frac{1}{\zeta^2(s)} \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{n^s},$$

where $\lceil \cdot \rceil$ and $\lfloor \cdot \rfloor$ represent the usual ceiling and floor functions, respectively.

Let us establish upper and lower bounds for the double summation. Initially, notice that the inner summation satisfies the following bounds:

$$\begin{aligned} \sum_{n=\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{n^s} &= \frac{1}{\lceil f(m) \rceil^s} + \sum_{n=\lceil f(m) \rceil+1}^{\lfloor g(m) \rfloor} \frac{1}{n^s} \\ &\leq 1 + \int_{\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{x^s} dx \\ &\leq 1 + \int_{f(m)}^{g(m)} \frac{1}{x^s} dx \\ &= 1 + \frac{1}{-s+1} (g(m)^{-s+1} - f(m)^{-s+1}). \end{aligned}$$




Figure 1: A set upper and lower bounded by two functions.

Thus, an upper bound for the double summation is expressed by

$$\begin{aligned} \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{n^s} &\leq \sum_{m=1}^{\infty} \frac{1}{m^s} \left(1 + \frac{1}{-s+1} (g(m)^{-s+1} - f(m)^{-s+1}) \right) \\ &= \zeta(s) + \frac{1}{-s+1} \sum_{m=1}^{\infty} \frac{1}{m^s} (g(m)^{-s+1} - f(m)^{-s+1}). \end{aligned}$$

Performing analogous manipulations, we obtain the following lower bound for the inner summation:

$$\begin{aligned} \sum_{n=\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{n^s} &\geq -\frac{1}{(\lceil f(m) \rceil - 1)^s} + \int_{\lceil f(m) \rceil - 1}^{\lfloor g(m) \rfloor + 1} \frac{1}{x^s} dx \\ &\geq -1 + \int_{f(m)}^{g(m)} \frac{1}{x^s} dx \\ &= -1 + \frac{1}{-s+1} (g(m)^{-s+1} - f(m)^{-s+1}). \end{aligned}$$

This implies that the double summation is lower bounded by

$$\begin{aligned} \sum_{m=1}^{\infty} \frac{1}{m^s} \sum_{n=\lceil f(m) \rceil}^{\lfloor g(m) \rfloor} \frac{1}{n^s} &\geq \sum_{m=1}^{\infty} \frac{1}{m^s} \left(-1 + \frac{1}{-s+1} (g(m)^{-s+1} - f(m)^{-s+1}) \right) \\ &= -\zeta(s) + \frac{1}{-s+1} \sum_{m=1}^{\infty} \frac{1}{m^s} (g(m)^{-s+1} - f(m)^{-s+1}). \end{aligned}$$

The upper and lower bounds present similar formulations, inviting an application of the squeeze theorem. Thus, after dividing both expressions by $\zeta^2(s)$ and taking the limit as $s \downarrow 1$, minor manipulations furnish

$$\text{dens}(C) = \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}).$$

Now let us analyze the density of a set C in the light of the asymptotic behavior of the delimiting functions.

Proposition 3.3 (Asymptotics) *Let $u(m)$ and $v(m)$ be delimiting functions that are always greater or equal to one. If $f(m) = \Theta(u(m))$ and $g(m) = \Theta(v(m))$, then*

$$(3.1) \quad \text{dens}(C) = \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{\infty} \frac{1}{m^s} (u(m)^{-s+1} - v(m)^{-s+1}).$$

Proof By the definition of the Θ -notation [17, p. 434], there exists a quantity m_0 such that, for every $m \geq m_0$, both functions f and g satisfy:

$$\begin{aligned} c_1 u(m) &\leq f(m) \leq c_2 u(m), \\ c_3 v(m) &\leq g(m) \leq c_4 v(m), \end{aligned}$$

where c_1, c_2, c_3 , and c_4 are positive constants. Moreover, notice that

$$\begin{aligned} \text{dens}(C) &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}) \\ &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \left[\sum_{m=1}^{m_0-1} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}) \right] \\ &\quad + \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \left[\sum_{m=m_0}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}) \right] \\ &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=m_0}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}). \end{aligned}$$

Thus, for $m \geq m_0$, we have that

$$\begin{aligned} &\sum_{m=m_0}^{\infty} \frac{1}{m^s} \left((c_2 u(m))^{-s+1} - (c_3 v(m))^{-s+1} \right) \\ &\leq \sum_{m=m_0}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}) \\ &\leq \sum_{m=m_0}^{\infty} \frac{1}{m^s} \left((c_1 u(m))^{-s+1} - (c_4 v(m))^{-s+1} \right). \end{aligned}$$

Now we show that after dividing by $\zeta(s)$ and letting $s \downarrow 1$, both upper and lower bounds above have the same limit. Since $u(m) \geq 1$ and $v(m) \geq 1$, it follows that for arbitrary positive constants k_1 and k_2

$$\begin{aligned} & \sum_{m=m_0}^{\infty} \frac{1}{m^s} \left((k_1 u(m))^{-s+1} - (k_2 v(m))^{-s+1} \right) \\ &= k_1^{-s+1} \sum_{m=m_0}^{\infty} \frac{1}{m^s} u(m)^{-s+1} - k_2^{-s+1} \sum_{m=m_0}^{\infty} \frac{1}{m^s} v(m)^{-s+1}. \end{aligned}$$

Thus, since both k_1^{-s+1} and k_2^{-s+1} tend to one as $s \downarrow 1$, we have that

$$\begin{aligned} & \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=m_0}^{\infty} \frac{1}{m^s} \left((k_1 u(m))^{-s+1} - (k_2 v(m))^{-s+1} \right) \\ &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=m_0}^{\infty} \frac{1}{m^s} (u(m)^{-s+1} - v(m)^{-s+1}). \end{aligned}$$

Therefore, we maintain that

$$\begin{aligned} & \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=m_0}^{\infty} \frac{1}{m^s} (f(m)^{-s+1} - g(m)^{-s+1}) \\ &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=m_0}^{\infty} \frac{1}{m^s} (u(m)^{-s+1} - v(m)^{-s+1}). \end{aligned}$$

Finally, since

$$\lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{m_0-1} \frac{1}{m^s} (u(m)^{-s+1} - v(m)^{-s+1}) = 0,$$

the proposition is proven. ■

We now supply two examples. But first, a simple lemma is needed.

Lemma 3.4 For $\alpha \geq 0$, $\lim_{s \downarrow 1} \zeta((\alpha + 1)s - \alpha)(s - 1) = (1 + \alpha)^{-1}$.

Proof Taking into account the substitution $t = (\alpha + 1)s - \alpha$, it follows that:

$$\begin{aligned} \lim_{s \downarrow 1} \zeta((\alpha + 1)s - \alpha)(s - 1) &= \lim_{t \downarrow 1} \zeta(t) \left(\frac{t + \alpha}{1 + \alpha} - 1 \right) \\ &= \lim_{t \downarrow 1} \zeta(t) \frac{t - 1}{1 + \alpha} \\ &= \frac{1}{1 + \alpha}. \end{aligned} \quad \blacksquare$$

Example 3.5 Let us examine the density of the set $C_{\text{pow}} = \{(m, n) \in \mathbb{P}^2 : f(m) \leq n \leq g(m)\}$, where $g(m) = \Theta(m^\beta)$ and $f(m) = \Theta(m^\alpha)$, for real quantities $\beta \geq \alpha > 0$. In order to compute such density we need the previous lemma. Thus, invoking equation (3.1), it follows that the sought density is given by

$$\begin{aligned} \text{dens}(C_{\text{pow}}) &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{\infty} \frac{1}{m^s} ((m^\alpha)^{(-s+1)} - (m^\beta)^{(-s+1)}) \\ &= \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \left(\zeta((\alpha + 1)s - \alpha) - \zeta((\beta + 1)s - \beta) \right) \\ &= \frac{1}{1 + \alpha} - \frac{1}{1 + \beta}. \end{aligned}$$

In particular, if $\alpha\beta = 1$, we have $\text{dens}(C_{\text{pow}}) = \frac{\beta-1}{\beta+1}$.

Example 3.6 Consider the set $C_{\text{exp}} = \{(m, n) \in \mathbb{P}^2 : n \leq g(m)\}$, where $g(m) = \Theta(a^m)$, for $a > 1$. Thus, by equation (3.1),

$$\text{dens}(C_{\text{exp}}) = \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \left(\zeta(s) - \sum_{m=1}^{\infty} \frac{(a^m)^{-s+1}}{m^s} \right).$$

Then note that for each $\beta > 0$, there is a quantity M such that $m \geq M$ implies that $a^m \geq m^\beta$. By Example 1, we know that

$$\lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{\infty} \frac{(m^\beta)^{-s+1}}{m^s} = \frac{1}{1 + \beta}.$$

Moreover, since $\lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=1}^{M-1} \frac{(m^\beta)^{-s+1}}{m^s} = 0$, it follows that

$$\lim_{s \downarrow 1} \frac{1}{\zeta(s)} \sum_{m=M}^{\infty} \frac{(m^\beta)^{-s+1}}{m^s} = \frac{1}{1 + \beta}.$$

Thus,

$$\begin{aligned} \text{dens}(C_{\text{exp}}) &\geq \lim_{s \downarrow 1} \frac{1}{\zeta(s)} \left(\zeta(s) - \left(\sum_{m=1}^{M-1} \frac{(a^m)^{-s+1}}{m^s} + \sum_{m=M}^{\infty} \frac{(m^\beta)^{-s+1}}{m^s} \right) \right) \\ &= 1 - \left(0 + \frac{1}{1 + \beta} \right) = \frac{\beta}{\beta + 1}. \end{aligned}$$

Finally, letting $\beta \rightarrow \infty$ yields $\text{dens}(C_{\text{exp}}) = 1$.

References

- [1] R. Ahlswede and L. H. Khachatryan, *Classical results on primitive and recent results on cross-primitive sequences*. In: The mathematics of P. Erdős, Vol. 1, Algorithms and Combinatorics B. Springer-Verlag, Berlin, 1996, 104–116.
- [2] ———, *Number theoretic correlation inequalities for Dirichlet densities*. J. Number Theory **63**(1997), 34–46. <http://dx.doi.org/10.1006/jnth.1997.2072>
- [3] R. Ahlswede, L. H. Khachatryan and A. Sárközy, *On the counting function for primitive sets of integers*. J. Number Theory **79**(1999), 330–344. <http://dx.doi.org/10.1006/jnth.1999.2427>
- [4] R. Ahlswede, L. H. Khachatryan and A. Sárközy, *On the density of primitive sets*. J. Number Theory **109**(2004), 319–361. <http://dx.doi.org/10.1016/j.jnt.2004.07.011>
- [5] P. T. Bateman and H. G. Diamond, *Analytic Number Theory*. World Scientific Publishing Co., Hackensack, NJ, 2004.
- [6] J. P. Bell and S. N. Burris, *Dirichlet density extends global asymptotic density in multiplicative systems*. Preprint, <http://www.math.uwaterloo.ca/~snburris/htdocs/MYWORKS/PREPRINTS/odelta.pdf>.
- [7] L. Cheo, *On the density of sets of Gaussian integers*. Amer. Math. Monthly **58**(1951), 618–620. <http://dx.doi.org/10.2307/2306357>
- [8] H. Davenport and P. Erdős, *On sequences of positive integers*. J. Indian Math. Soc. **15**(1951), 19–24.
- [9] R. L. Duncan, *Some continuity properties of the Schnirelmann density*. Pacific J. Math. **26**(1968), 57–58.
- [10] ———, *Some continuity properties of the Schnirelmann density II*. Pacific J. Math. **32**(1970), 65–67.
- [11] P. Erdős, *On the density of some sequences of integers*. Bull. Amer. Math. Soc. **54**(1948), 685–692. <http://dx.doi.org/10.1090/S0002-9904-1948-09057-7>
- [12] A. R. Freedman, *On asymptotic density in n -dimensions*. Pacific J. Math. **29**(1969), 95–113.
- [13] ———, *On the generalization of the modified Schnirelmann density*. J. Number Theory **2**(1970), 97–105. [http://dx.doi.org/10.1016/0022-314X\(70\)90009-0](http://dx.doi.org/10.1016/0022-314X(70)90009-0)
- [14] ———, *On the additivity theorem for n -dimensional asymptotic density*. Pacific J. Math. **49**(1973), 357–363.
- [15] A. Fuchs and R. G. Antonini, *Théorie générale des densités*. Rend. Accad. Naz. Sci. XL Mem. Mat. **14**(1990), 253–294.
- [16] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*. Fourth edition, Academic Press, New York, 1965.
- [17] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*. Addison-Wesley Publishing Company, Reading, MA, 1989.
- [18] B. Kvarda, *On densities of sets of lattice points*. Pacific J. Math. **13**(1963), 611–615.
- [19] B. Kvarda and R. Killgrove, *Extensions of the Schnirelmann density to higher dimensions*. Amer. Math. Monthly **73**(1966), 976–979. <http://dx.doi.org/10.2307/2314499>
- [20] Ivan Niven, *The asymptotic density of sequences*. Bull. Amer. Math. Soc. **57**(1951), 420–434. <http://dx.doi.org/10.1090/S0002-9904-1951-09543-9>
- [21] Walter Rudin, *Real and Complex Analysis*. McGraw-Hill, New York–Toronto–London, 1966.
- [22] R. Stalley, *A modified Schnirelmann density*. Pacific J. Math. **5**(1955), 119–124.
- [23] X. G. Sun and J. H. Fand, *On the density of integers of the form $(p - 1)2^{-n}$ in arithmetic progression*. Bull. Aust. Math. Soc. **78**(2008), 431–436. <http://dx.doi.org/10.1017/S0004972708000804>

Departamento de Estatística, Universidade Federal de Pernambuco, Cidade Universitária, 50740-540, Recife, PE, Brazil

e-mail: leandro@de.ufpe.br rjds@stat.ufpe.org