Polach, H A, 1979, Correlation of <sup>14</sup>C activity of NBS oxalic acid with Arizona 1850 wood and ANU sucrose standards, *in* Berger, Rainer and Suess H E, eds, Radiocarbon dating, Internatl conf on radiocarbon dating, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 115-124.

Rhodes, E. G., Polach, H. A., Thom, B. G., and Wilson, S. R., 1980, Age structure of Holocene coastal sediments: Gulf of Carpentaria, Australia, in Stuiver, Minze and Kra, Renee, eds, Internatl radiocarbon conf. 10th, Proc. Radiocarbon, v. 22, no. 3,

p 718-727

Thom, B G, Bowman, G M, Gillespie, R, Temple, R, and Barbetti, M, 1981, Radiocarbon dating of Holocene beach-ridge sequences in South-East Australia: Dept Geog, Fac Military Studies, Univ New South Wales, Duntroon, ACT, Australia, Mon no. 11, 84 p.

Thom, B G, Polach, H A, and Bowman, G M, 1978, Holocene age structure of coastal sand barriers in New South Wales, Australia: Dept Geog, Fac Military Studies,

Univ New South Wales, Duntroon, ACT, Australia, 86 p.

Wilson, S R and Ward G K, 1981, Evaluation and clustering of radiocarbon age determinations: procedures and paradigms: Archaeometry, v 23, p 19-39.

## **ERRATUM**

## MARINE RESOURCES RESEARCH INSTITUTE RADIOCARBON DATES IV

## THOMAS D MATHEWS and FONDA L KEARNS

Marine Resources Research Institute, P O Box 12559, Charleston, South Carolina 29412

(R, 1981, v 23, p 222-226). Due to an error in the value for NBS oxalic acid (as entered into the MRRI computer program for age and standard deviation calculations) the dates reported in this date list and repeated in the index are all incorrect. Each date should be younger by 2960 years, resulting in finite ages for most of the samples. The exceptions are MRRI -223, -239, -249, -263, -264, -267, -269, and -274-277, which are all modern in age or in the future. We interpret these future dates as indications of contamination with bomb <sup>14</sup>C. No other dates were affected in any way since the dates in this list were calculated as a group.