

## ASSESSING LOCK-IN DEPTH AND ESTABLISHING A LATE HOLOCENE PALEOMAGNETIC SECULAR VARIATION RECORD FROM THE MONGOLIAN ALTAI

Marcel Bliedtner<sup>1\*</sup>  • Torsten Haberzettl<sup>2</sup> • Norbert Nowaczyk<sup>3</sup> • Enkhtuya Bazarradnaa<sup>4</sup> • Roland Zech<sup>1</sup> • Paul Strobel<sup>1</sup>

<sup>1</sup>Institute of Geography, Friedrich Schiller University Jena, 07743 Jena, Germany

<sup>2</sup>Physical Geography, Institute for Geography and Geology, University of Greifswald, 17489 Greifswald, Germany

<sup>3</sup>Helmholtz Centre Potsdam GFZ, Section Climate Dynamics and Landscape Evolution, 14473 Potsdam, Germany

<sup>4</sup>Institute of Plant and Agricultural Sciences, Mongolian University of Life Sciences, 45047 Darkhan, Mongolia

**ABSTRACT.** Although paleomagnetic secular variations (PSV) often corroborate radiocarbon (<sup>14</sup>C)-based lacustrine sediment chronologies, this is not the case at the high-altitude site Khar Nuur in the Mongolian Altai Mountains. Our results show that the inclination pattern resembles those from a regional reference record from Shireet Naiman Nuur and global geomagnetic field models very well, but with a constant offset of  $730 \pm 90$  yr. Possible reservoir effects from terrestrial pre-aging and hardwater effects can be excluded as the cause of the  $\sim 730$ -yr offset because the different dated compounds correspond very well to each other, and modern reservoir effects are negligible. Instead, the constant  $\sim 730$ -yr offset in the PSV pattern is likely the result of a constant lock-in depth of  $26 \pm 2$  cm below the sediment-water interface at Khar Nuur. This assumption is supported by comparison of paleoclimatological proxies from Shireet Naiman Nuur, where similarities are obvious for the <sup>14</sup>C-based chronology of Khar Nuur without a  $\sim 730$ -yr adjustment. Therefore, the previously published <sup>14</sup>C-based chronology of Khar Nuur provides a reliable age control. Accepting the lock-in depth of  $26 \pm 2$  cm, the good consistency in inclination between Khar Nuur and global geomagnetic field models highlights the reliability of the latter even in a paleomagnetically understudied area.

**KEYWORDS:** lake sediments, paleoenvironmental changes, paleomagnetic secular variations, radiocarbon dating, semi-arid Mongolia.

### INTRODUCTION

Precise age control is an important precondition for robust paleoenvironmental reconstructions from lacustrine sediments. However, in semi-arid regions such as Mongolia, chronological control of the existing paleoenvironmental lacustrine records is often imprecise because most chronologies rely on few <sup>14</sup>C-dates, almost exclusively made on bulk organic material (e.g., Sun et al. 2013; Unkelbach et al. 2019; Rudaya et al. 2021). Unfortunately, bulk organic carbon often needs to be dated in semi-arid regions due to the absence of terrestrial macrofossils in lake sediments. Compared to terrestrial macrofossils, which are assumed to be rapidly deposited in lakes (Hajdas et al. 1995), bulk organic carbon can be “pre-aged” because organic material accumulates in the catchment over hundreds to thousands of years. Consequently, bulk organic carbon is possibly too old when finally ending up in the lake, overestimating its “true” deposition age (Haberzettl et al. 2013; Wüdsch et al. 2014; Gierga et al. 2016; Douglas et al. 2018; Haas et al. 2019). Moreover, lake sediments comprise organic material from terrestrial and aquatic sources and the radiocarbon (<sup>14</sup>C)-signal from aquatic contributions can additionally be affected by the so-called “hardwater effect” due to incorporation of carbonate from old calcareous bedrock and/or re-dissolved carbon from in-lake carbonate precipitation (Reinwarth et al. 2013). Therefore, the

\*Corresponding author. Email: [marcel.bliedtner@uni-jena.de](mailto:marcel.bliedtner@uni-jena.de)



reliability of paleoenvironmental reconstructions can be largely limited due to chronological uncertainties of the potentially reservoir affected  $^{14}\text{C}$ -signal.

In this context, paleomagnetic secular variation (PSV) stratigraphy is a valuable chronostratigraphic tool to evaluate and potentially refine  $^{14}\text{C}$ -based chronologies (Barletta et al. 2010; Ólafsdóttir et al. 2013; Haberzettl et al. 2015, 2019, 2021; Bliedtner et al. 2022). PSV changes are ideally suited to study short-term changes especially throughout the Holocene. The principle behind PSV in lacustrine sediments is that magnetic particles align themselves with the surrounding geomagnetic field during deposition and preserve their direction when locked-in during ongoing deposition (Merrill and McFadden 2007; Roberts and Turner 2013). The recorded pattern of inclination and declination can therefore serve as valuable chronostratigraphic indicators. For the semi-arid regions of Mongolia, we recently proposed a dual chronological approach in the high-altitude Shireet Naiman Nuur (Nuur = lake) sediments (46°31'55.04"N, 101°49'16.23"E; Khangai Mountains, central Mongolia), where we (1) established a high-resolution  $^{14}\text{C}$ -chronology for the past 7.4 cal. ka BP by dating terrestrial macrofossils and (2) corroborated the  $^{14}\text{C}$ -based chronology by a newly obtained PSV record from the lake (Bliedtner et al. 2022). The PSV record from Shireet Naiman Nuur very well confirms our  $^{14}\text{C}$ -based chronology and provides a valuable regional master record for using PSV stratigraphy in semi-arid Mongolia to evaluate and potentially refine  $^{14}\text{C}$ -chronologies when compared to our record. Therefore, if successful, the combination of  $^{14}\text{C}$ -dating and corroboration by PSV data is a valuable chronological approach to establish more reliable chronologies in semi-arid regions, especially regarding the large age uncertainties of bulk organic carbon dating often has in such regions.

Here we present a newly obtained PSV record from the high-altitude Khar Nuur in the Mongolian Altai Mountains. As in this case PSV data do not corroborate the existing good age control published by Bliedtner et al. (2021), we aim to detect causes for age offsets and establish a reliable continuous inclination and declination record for the last 4.2 cal. ka BP from the lake.

## MATERIAL AND METHODS

### Study Area

Khar Nuur is of para-glacial origin and situated at high-altitude (2486 m a.s.l.) in the Mongolian Altai (48°37'22.9"N, 88°56'42.5"E; Figure 1). The lake was previously described in detail by Strobel et al. (2021, 2022). The endorheic Khar Nuur basin was formed during the Last Glacial Maximum by glacial advances from the Tsengel Khairkhan Massif, leaving prominent moraine lobes in the main valley (Walther et al. 2017). The lake has a small hydrological catchment (44.8 km<sup>2</sup>) with quite steep slopes, which are mostly composed of friable black clay shales and granitic moraine deposits. Due to the continental position of Khar Nuur, mean annual temperatures and precipitation are low with  $-2.4 \pm 0.7^\circ\text{C}$  and  $145 \pm 35$  mm, respectively (station Tolbo sum; period 2009–2019; DWD Climate Data Center 2020). Moreover, the lake is ice-covered for 8–9 months per year (2016–2019; Planet Team 2017).

### Chronostratigraphy of the Khar Nuur Sediments

The investigated lake sediments of Khar Nuur and its chronostratigraphy were previously described in detail by Bliedtner et al. (2021). The 141-cm-long sediment core from Khar




Figure 1 Overview of semi-arid Mongolia located in central Asia. The investigated site at Khar Nuur (KN) in the Mongolian Altai Mountains is indicated by the yellow star and the regional comparison record from Shireet Naiman Nuur (SNN) (Bliedtner et al. 2022) in the Mongolian Khangai Mountains is indicated by the white star. (Please see online version for color figures.)




Figure 2 Chronostratigraphy of the Khar Nuur sediments previously published by Bliedtner et al. (2021). Inclination, declination, maximum angular deviation (MAD), median destructive field (MDF) and natural remanent magnetization (NRM) of the Khar Nuur sediments.

Nuur consists of well laminated silty sediments, which are characterized by alternating phases of increased lake primary productivity in the brown and blackish dark sediments and reduced productivity in the bluish-gray sediments (Figure 2). The chronology is based on  $^{14}\text{C}$ -datings of 12 bulk organic carbon samples, 8 aquatic macrofossil samples and 1 ostracod sample. Additionally, we accounted for possible terrestrial reservoir and hardwater effects by  $^{14}\text{C}$ -dating a surface sediment bulk organic carbon sample and one modern water plant at the coring location.  $^{14}\text{C}$ -analyses were carried out at the Laboratory for the Analysis of

Radiocarbon with AMS (LARA AMS) of the University of Bern, Switzerland (Szidat 2014) using a Mini CARbon DAting System (MICADAS) AMS coupled online to an Elemental Analyzer (Ruff et al. 2010; Wacker et al. 2010). Results were reported as fraction modern ( $F^{14}\text{C}$ ) and corrected for cross and constant contamination after the contamination drift model of Salazar et al. (2015). Resulting  $^{14}\text{C}$ -ages were calibrated with the IntCal20 calibration curve (Reimer et al. 2020) using OxCal (Bronk Ramsey 2009), except the surface sediment bulk organic carbon and modern water plant  $^{14}\text{C}$ -age, which were calibrated with the bomb 13 NH1 calibration curve (Hua et al. 2013) likewise using OxCal.

All  $^{14}\text{C}$  results can be found in the Supplementary Table S1.  $^{14}\text{C}$ -dating of the surface sediment gave a slight terrestrial pre-aging with a mean  $^{14}\text{C}$ -age of  $129 \pm 85$  cal. BP (age range:  $-7$  to 279 cal. BP [95.4%]). However, no hardwater effect was presently indicated by modern  $^{14}\text{C}$ -ages of the modern water plant (Strobel et al. 2021; Bliedtner et al. 2022). Based on the  $^{14}\text{C}$ -ages, a chronology was established for the Khar Nuur sediments by Bayesian age-depth modeling using the Bacon 2.3.4 package in R (Blaauw and Christen 2011). During Bayesian age-depth modeling, the slight terrestrial pre-aging indicated by the surface sediment bulk organic carbon sample was subtracted from all conventional bulk organic carbon  $^{14}\text{C}$ -ages. Bayesian age-depth modeling gave a modeled basal median age of  $4.2^{+0.4}_{-0.3}$  cal. ka BP with all aquatic macrofossil  $^{14}\text{C}$ -ages being in stratigraphic position, likely indicating the timing of sediment deposition (Figure 2). Terrestrial pre-aging corrected bulk organic carbon  $^{14}\text{C}$ -ages very well correspond to the aquatic macrofossil  $^{14}\text{C}$ -ages, except for four ages at 90, 40, 30, and 20 cm sediment depth, which were pre-aged and overestimated their timing of deposition. Those samples contain an erosive signal from aged terrestrial organic carbon in the catchment soils and we therefore excluded them from Bayesian age-depth modeling (Figure 2).

### Paleomagnetic Measurements

Natural remanent magnetization (NRM) was continuously measured at 1-cm intervals on a u-channel from the Khar Nuur sediment core with a 2G Enterprise DC-4K liquid helium-free magnetometer at the GFZ Potsdam. NRM was measured using stepwise (11 steps) alternating field (AF) demagnetization with peak AFs of 0, 5, 10, 15, 20, 30, 40, 50, 65, 80, and 100 mT. Inclination and declination of the characteristic remanent magnetization and maximum angular deviation (MAD) were derived from principal component analyses (Kirschvink 1980). The median destructive field (MDF) was also calculated. We have to note that declination data are relative and centered to 0 because the azimuth could not be controlled during coring. Because of the width of the response function of the SQUID sensors ( $\sim 7.8$  cm), some smoothing occurs in the data. In order to eliminate edge effects, the upper and lowermost 5 cm are plotted in gray.

## RESULTS

NRM intensities range from 0.6 to  $17.6 \text{ mA m}^{-1}$  in the Khar Nuur sediments and MDF values from 28.5 to only 51.7 mT at the very top of the sediment core (Figure 2), indicating a low coercivity mineral such as magnetite as carrier of the natural remanent magnetization. Well preservation and stable single-component magnetization of the sediments is indicated by low MAD values (Stoner and St-Onge 2007) below  $5.6^\circ$ . Orthogonal vector plots confirm stable single-component magnetizations, which trend to the origin during AF treatments (see Supplementary Figure SI 1). Distinct changes are shown by the declination and




Figure 3 Comparison of (A) inclination and (B) declination pattern of the Khar Nuur sediments with the regional reference sediments from Shireet Naiman Nuur (Bliedtner et al. 2022) and the spherical harmonic geomagnetic field model HFM.OL1.A1 (Constable et al. 2016).

inclination pattern throughout the sediment core. Declination has a wide relative range from  $-53$  to  $23^\circ$  and stretches over  $76^\circ$ . Inclination ranges from  $-3.3$  to  $62.6^\circ$ , however, especially inclination seem to be offset in the uppermost part of the sediment core due to disturbed sediment and we therefore plotted the PSV pattern as a dotted gray line there. By excluding the potentially reworked uppermost part of the sediment core, inclination narrows and ranges from  $40.6$  to  $62.6^\circ$  (Figure 2). Compared to the expected inclination derived from the geocentric axial dipole model ( $GAD = 66.22^\circ$ ), inclination in the Khar Nuur sediments is flattened and did not intersect with the expected one for the latitude of this site. This has repeatedly been noticed in lake sediments and attributed to postdepositional compaction processes or compaction during coring (Tauxe et al. 2008; Henkel et al. 2016).

## DISCUSSION

Distinct changes in inclination during the past 4.2 cal. ka BP in the Khar Nuur sediments are similar to the regional reference record from Shireet Naiman Nuur (Bliedtner et al. 2022) and various outputs of spherical harmonic geomagnetic field models (see Supplementary Figure SI 2 for comparison with different models, exemplarily shown here is the HFM.OL1.A1 model by Constable et al. [2016]; Figure 3A). Despite the similarities in inclination, the Khar Nuur inclination pattern appears to have a temporal offset of  $730 \pm 90$  yr. An adjusted Khar Nuur chronology of  $\sim 730$  yr, i.e., where we subtracted  $\sim 730$  yr from the original  $^{14}\text{C}$ -chronology, corresponds very well to the Shireet Naiman Nuur reference record and geomagnetic field model, now also temporally resembling their pattern (Figure 3A). Declination also shows no similarities to the regional reference and model pattern without an adjustment of the Khar Nuur  $^{14}\text{C}$ -chronology for  $\sim 730$  yr (Figure 3B). However, also with an adjustment, similarities are not obvious. Less similarities between declination records have also been reported from lakes from

the Tibetan Plateau (Haberzettl et al. 2015). Moreover, it has often been reported that unwanted and unknown core rotation could largely affect the declination pattern (Nilsson et al. 2018), which might explain the observed differences in the Khar Nuur sediment core. However, if the large shift in declination at 50 cm sediment depth stretching over  $\sim 50^\circ$  is attributed to core rotation and hence neglected, some similarities in small scale variations can be observed between Khar Nuur and the HFM.OL1.A1 model. Unfortunately, these small-scale declination changes are missing in the Shireet Naiman Nuur sediments which might be explained by the lower sedimentation rates of the Shireet Naiman Nuur sediments (Ohlendorf et al. 2014).

The  $\sim 730$ -yr offset in inclination at Khar Nuur could be explained by remaining reservoir effects in the bulk organic carbon and aquatic macrofossil  $^{14}\text{C}$ -ages, potentially biasing the  $^{14}\text{C}$ -based chronology. However, we have good indications that reservoir effects, i.e., from terrestrial pre-aging and hardwater effects, are negligible in the Khar Nuur sediments. While we corrected the bulk organic carbon  $^{14}\text{C}$ -ages for a slight terrestrial pre-aging using a surface sediment sample from the coring location with a  $^{14}\text{C}$ -age of  $129 \pm 85$  cal. BP (age range:  $-7$  to 279 cal. BP [95.4%]), no hardwater effects were found by modern  $^{14}\text{C}$ -ages of a modern water plant at the coring location (Bliedtner et al. 2021; Strobel et al. 2021). Although reservoir effects are not recently present in the surface sediments of Khar Nuur, they can still variably occur through time. However, the very good agreement of the bulk organic carbon and aquatic macrofossil  $^{14}\text{C}$ -ages questions variable reservoir effects, especially since the bulk organic carbon  $^{14}\text{C}$ -ages, corrected for terrestrial pre-aging, should include hardwater effects to a lesser degree than the aquatic macrofossils and would therefore be younger. Moreover, the bulk organic carbon and aquatic macrofossil  $^{14}\text{C}$ -age at 10 cm sediment depth gave younger modeled  $^{14}\text{C}$ -ages ( $0.6 \pm 0.3$  and  $0.6^{+0.3}_{-0.2}$  cal. ka BP) than the observed offset of  $\sim 730$  yr in inclination (Bliedtner et al. 2021), further providing evidence for non-existing reservoir effects. Ultimately, a constant reservoir effect of  $\sim 730$  yr over time, especially a constant hardwater effect, is rather unlikely for the Khar Nuur sediments. In the carbonate-free Khar Nuur catchment only the incorporation of bicarbonate ( $\text{HCO}_3^-$ ) could provide old carbonate for potential hardwater effects (Strobel et al. 2021). However, distinct lake primary productivity changes occur in the Khar Nuur sediments, which are related to changes in growing season temperature and the duration of ice cover (Bliedtner et al. 2021). Especially changes in the duration of ice coverage would result in a variable hardwater effect due to increased/reduced exchange with atmospheric  $^{14}\text{CO}_2$  and the reduced/increased incorporation of  $\text{H}^{14}\text{CO}_3^-$  (i.e., old carbon), respectively. Since the  $\sim 730$ -yr offset in inclination is constant throughout the Khar Nuur sediments, a bias from remaining variable reservoir effects, and especially from hard water effects, is therefore very unlikely as this would have caused unpredictable offsets between the individual inclination highs and lows in the compared records after the adjustment of  $\sim 730$  yr. Therefore, the  $^{14}\text{C}$ -based chronology provides a robust age control.

A more likely explanation for the observed age offset of  $\sim 730$  yr at Khar Nuur might be that magnetic particles become finally locked-in several centimeters below the sediment-water interface, which is the so-called “lock-in depth” (Roberts and Winklhofer 2004; Mellström et al. 2015). It has been proposed that magnetic particles can re-align to the geomagnetic field after deposition during dewatering and compaction, incorporating a post-depositional remanent magnetization (pDRM) signal which can lead to a delay and smoothing of the recorded geomagnetic signal (Roberts and Winklhofer 2004; Nilsson et al. 2018). Although the underlying physical processes are poorly understood, pDRM is thought to mostly

depend on water content, grain size, magnetic concentration, sedimentation rate and/or bioturbation (Tauxe et al. 2006; Roberts et al. 2013; Zhao et al. 2016; Valet et al. 2017). Evidence for lock-in depths in lacustrine environments is mostly from varved sediments so far, but lock-in depths up to several decimeters below the sediment-water-interface were repeatedly reported, leading to centennial to millennial delays of the geomagnetic signal (e.g., Stockhausen 1998; Snowball et al. 2013; Mellström et al. 2015; Scheidt et al. 2022). Considering the relatively constant sedimentation rate at Khar Nuur (Figure 2), a lock-in depth of  $26 \pm 2$  cm would very likely explain the constant  $730 \pm 90$  year offset in PSV.

To finally confirm that the observed age offset of  $730 \pm 90$  yr in PSV at Khar Nuur is indeed attributed to a lock-in depth of  $26 \pm 2$  cm and not to potential reservoir effects, we compare paleoclimatological proxies from the Khar Nuur sediments with those from the well-dated Shireet Naiman Nuur sediments (Bliedtner et al. 2022; see Figure 1 for location). Log ratio of calcium and titanium ( $\log \text{Ca/Ti}$ ) and center log ratios of Ti ( $\text{Ti}_{\text{clr}}$ ) are therefore plotted on both the  $^{14}\text{C}$ -based chronology and a  $\sim 730$ -yr adjusted chronology of Khar Nuur (Figure 4). Although the sample resolution slightly differs between Khar Nuur and Shireet Naiman Nuur, both lakes are ideal for direct comparison since they are located at high-altitudes ( $\sim 2400$  m a.s.l.) and have comparable environmental settings and therefore very likely respond to identical forcing mechanisms. Changes in  $\log (\text{Ca/Ti})$  are interpreted as changes in lake primary productivity controlled by growing season temperatures in both lakes, whereas  $\text{Ti}_{\text{clr}}$  indicates terrigenous minerogenic input (Bliedtner et al. 2021; Bliedtner et al. 2022). Although similarities can be found between the paleoclimatological proxies of Shireet Naiman Nuur and paleoclimatological proxies from Khar Nuur plotted on both, the  $^{14}\text{C}$ -based and  $\sim 730$  year adjusted chronology of Khar Nuur, a better fit is obvious between Shireet Naiman Nuur and the  $^{14}\text{C}$ -based chronology of Khar Nuur (Figure 4). Despite some differences in the magnitude, distinct changes in lake primary productivity and terrigenous input occur at  $\sim 4.1$ ,  $\sim 3$ , and  $\sim 1.8$  cal. ka BP. Between those distinct changes, productivity and terrigenous input follow similar characteristic trends (Figure 4). It is notable that slight temporal offsets in the order of 10–100 yr exist between Shireet Naiman Nuur and the  $^{14}\text{C}$ -based chronology of Khar Nuur, but those offsets are well within the  $2\sigma$  uncertainty of the chronologies of both records (Figure 4). Moreover, differences in the sedimentological proxies during the last  $\sim 1.5$  cal. ka BP could exist due to increased sediment erosion and relocation in the Khar Nuur catchment, potentially climatically and/or anthropogenically induced. This is also supported by the pre-aged bulk organic carbon  $^{14}\text{C}$ -ages in the sediments from  $\sim 45$  to 20 cm of the core, overestimating their true burial age due to longer residence times in the catchment soils which were hence removed from age-depth-modeling as extensively discussed by Bliedtner et al. (2021) (Figure 2).

Because of the very consistent trend of  $\log (\text{Ca/Ti})$  in the Khar Nuur and Shireet Naiman Nuur sediments, some paleoclimate interpretations can finally be made. Changes in  $\log (\text{Ca/Ti})$  have previously been interpreted in terms of changes in lake primary productivity primarily determined by growing season temperatures in those high-altitude lakes (Bliedtner et al. 2021, 2022). This is because air temperature strongly controls the duration of ice-cover and the growing season of aquatic producers at high altitudes (Willemsse and Törnqvist 1999; Mischke et al. 2010). Higher growing season temperatures are indicated in both lakes from  $\sim 4.2$ – $4.0$ ,  $3.2$ – $2.8$  and  $2.2$ – $1.9$  cal. ka BP (highlighted by reddish bars in Figure 4). In contrast, lower growing season temperatures occur from  $\sim 4.0$ – $3.2$ ,  $2.8$ – $2.2$ , and  $1.9$ – $1.6$  cal. ka BP (highlighted by bluish bars in Figure 4). Based on our well dated records from Khar Nuur and Shireet Naiman Nuur, a similar regional temperature evolution is suggested by




Figure 4 Comparison of the sedimentological proxies  $\log(\text{Ca}/\text{Ti})$  and center log-ratio  $\text{Ti}(\text{Ticlr})$  from Khar Nuur and Shireet Naiman Nuur (Bliedtner et al. 2022). Sedimentological proxies from Khar Nuur are shown both on the  $^{14}\text{C}$ -based chronology and the adjusted chronology for  $\sim 730$  yr (transparent curve). Please note that the black lines of the inorganic elements at Khar Nuur refers to a 2-point running and the black lines for the inorganic elements at Shireet Naiman Nuor refers to a 5-point running mean to create a comparable data resolution of 1 cm. Reddish (bluish) bars indicate periods of higher (lower) growing season temperature at Khar Nuur and Shireet Naiman Nuor. The timespan from 1.5 cal. ka BP until present is separated by the dashed black line since climatically and/or anthropogenically induced sediment erosion and relocation took place in the Khar Nuur catchment during this time.

the very good agreement of  $\log(\text{Ca}/\text{Ti})$  for the Altai Mountains (i.e., western Mongolia) and the Khangai Mountains (i.e., central Mongolia). Our results are further supported by previous regional studies, e.g., Bayan Nuur (Yang et al. 2020) and Telmen Nuur (Struck et al. 2022), who suggested total solar irradiance (TSI) as a main driver for temperature dynamics during the Late Holocene in the region, i.e., colder conditions during minima in TSI and warmer conditions during maxima in TSI.

## CONCLUSIONS

Inclination pattern from Khar Nuur closely resembles those from the regional reference record from Shireet Naiman Nuur and the geomagnetic field models, however, with a constant offset of  $730 \pm 90$  yr. We can exclude that the constant  $\sim 730$ -yr offset is caused by reservoir effects

from terrestrial pre-aging and hardwater effects because bulk organic carbon and aquatic macrofossil  $^{14}\text{C}$ -ages very well agree with each other, almost no modern reservoir effect was indicated by surficial sediment dating and dating a modern water plant, and the uppermost  $^{14}\text{C}$ -datings at 10-cm sediment depth gave younger ages than the offset of  $\sim 730$  yr in inclination. Instead, an almost constant lock-in depth of  $26 \pm 2$  cm below the sediment-water-interface very well explains the observed constant  $730 \pm 90$  yr offset in the PSV pattern at Khar Nuur. This is further supported by comparison of paleoclimatological proxies from Khar Nuur with Shireet Naiman Nuur, both high-altitude lakes with comparable environmental settings, where obvious similarities in  $\log(\text{Ca}/\text{Ti})$  and  $\text{Ti}_{\text{clr}}$  exist between Shireet Naiman Nuur and the  $^{14}\text{C}$ -based chronology of Khar Nuur.

We emphasize the importance of the potential lock-in of magnetic particles several centimeters below the sediment-water-interface for future studies since the lock-in depth can complicate the estimation of potential reservoir effects and finally challenge the refinement of  $^{14}\text{C}$ -based chronologies. Nevertheless, PSV will give an additional chronological indicator, and our combined approach will largely help to improve and potentially refine chronologies in semi-arid regions.

#### SUPPLEMENTARY MATERIAL

To view supplementary material for this article, please visit <https://doi.org/10.1017/RDC.2023.38>

#### ACKNOWLEDGMENTS

We would like to thank the Ernst Abbe Stiftung for financial support of the field trip to Mongolia and sediment retrieval. We further thank our logistic partners in Mongolia for their support during fieldwork. We thank two reviewers for their valuable and helpful comments on this paper.

#### AUTHOR CONTRIBUTIONS

Marcel Bliedtner: conceptualization, formal analysis, investigation, writing—original draft, writing—review and editing; Torsten Haberzettl: conceptualization, investigation, writing—review and editing; Norbert Nowaczyk: investigation, writing—review and editing; Enkhtuya Bazarradnaa: writing—review and editing; Roland Zech: writing—review and editing; Paul Strobel: conceptualization, investigation, writing—review and editing.

#### REFERENCES

- Barletta F, St-Onge G, Channell J, Rochon A. 2010. Dating of Holocene western Canadian Arctic sediments by matching paleomagnetic secular variation to a geomagnetic field model. *Quaternary Science Reviews* 29(17–18): 2315–2324. doi: [10.1016/j.quascirev.2010.05.035](https://doi.org/10.1016/j.quascirev.2010.05.035)
- Blaauw M, Christen JA. 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Analyses* 6(3):457–474. doi: [10.1214/11-BA618](https://doi.org/10.1214/11-BA618)
- Bliedtner M, Strobel P, Struck J, Salazar G, Szidat S, Nowaczyk N, Bazarradnaa E, Lloren R, Dubois N, Haberzettl T, Zech R. 2022. Holocene temperature variations in semi-arid central Mongolia—a chronological and sedimentological perspective from a 7400-year lake sediment record from the Khangai Mountains. *Frontiers in Earth Science* 10. doi: [10.3389/feart.2022.910782](https://doi.org/10.3389/feart.2022.910782)
- Bliedtner M, Struck J, Strobel P, Salazar G, Szidat S, Bazarradnaa E, Lloren R, Dubois N, Zech R. 2021. Late Holocene climate changes in the Altai region based on a first high-resolution biomarker isotope record from Lake Khar Nuur. *Geophysical Research Letters* 48(20). doi: [10.1029/2021GL094299](https://doi.org/10.1029/2021GL094299)

- Bronk Ramsey C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* 51:337–360. doi: [10.1017/S0033822200033865](https://doi.org/10.1017/S0033822200033865)
- Constable C, Korte M, Panovska S. 2016. Persistent high paleosecular variation activity in southern hemisphere for at least 10 000 years. *Earth and Planetary Science Letters* 453: 78–86. doi: [10.1016/j.epsl.2016.08.015](https://doi.org/10.1016/j.epsl.2016.08.015)
- Douglas PMJ, Pagani M, Eglinton TI, Brenner M, Curtis JH, Breckenridge A, Johnston K. 2018. A long-term decrease in the persistence of soil carbon caused by ancient Maya land use. *Nature Geoscience* 5:81. doi: [10.1038/s41561-018-0192-7](https://doi.org/10.1038/s41561-018-0192-7)
- Gierga M, Hajdas I, van Raden UJ, Gilli A, Wacker L, Sturm M, Bernasconi SM, Smittenberg RH. 2016. Long-stored soil carbon released by prehistoric land use: evidence from compound-specific radiocarbon analysis on Soppensee lake sediments. *Quaternary Science Reviews* 144:123–131. doi: [10.1016/j.quascirev.2016.05.011](https://doi.org/10.1016/j.quascirev.2016.05.011)
- Haas M, Baumann F, Castella D, Haghpor N, Reusch A, Strasser M, Eglinton TI, Dubois N. 2019. Roman-driven cultural eutrophication of Lake Murten, Switzerland. *Earth and Planetary Science Letters* 505:110–117. doi: [10.1016/j.epsl.2018.10.027](https://doi.org/10.1016/j.epsl.2018.10.027)
- Haberzettl T, Henkel K, Kasper T, Ahlborn M, Su Y, Wang J, Appel E, St-Onge G, Stoner J, Daut G, Zhu L, Mäusbacher R. 2015. Independently dated paleomagnetic secular variation records from the Tibetan Plateau. *Earth and Planetary Science Letters* 416:98–108. doi: [10.1016/j.epsl.2015.02.007](https://doi.org/10.1016/j.epsl.2015.02.007)
- Haberzettl T, Kasper T, Stoner JS, Rahobisoa JJ, Daut G. 2021. Extending and refining the paleomagnetic secular variation database for south-eastern Africa (Madagascar) to 2500 cal BP. *Earth and Planetary Science Letters* 565: 116931. doi: [10.1016/j.epsl.2021.116931](https://doi.org/10.1016/j.epsl.2021.116931)
- Haberzettl T, Kirsten KL, Kasper T, Franz S, Reinwarth B, Baade J, Daut G, Meadows ME, Su Y, Mäusbacher R. 2019. Using  $^{210}\text{Pb}$ -data and paleomagnetic secular variations to date anthropogenic impact on a lake system in the Western Cape, South Africa. *Quaternary Geochronology* 51:53–63. doi: [10.1016/j.quageo.2018.12.004](https://doi.org/10.1016/j.quageo.2018.12.004)
- Haberzettl T, St-Onge G, Behling H, Kirleis W. 2013. Evaluating Late Holocene radiocarbon-based chronologies by matching paleomagnetic secular variations to geomagnetic field models: an example from Lake Kalimpa (Sulawesi, Indonesia). *Geological Society, London, Special Publications* 373(1):245–259. doi: [10.1144/SP373.10](https://doi.org/10.1144/SP373.10)
- Hajdas I, Bonani G, Gošlar T. 1995. Radiocarbon dating the Holocene in the Gościąg Lake floating varve chronology. *Radiocarbon* 37(1): 71–74. doi: [10.1017/S0033822200014806](https://doi.org/10.1017/S0033822200014806)
- Henkel K, Haberzettl T, St-Onge G, Wang J, Ahlborn M, Daut G, Zhu L, Mäusbacher R. 2016. High-resolution paleomagnetic and sedimentological investigations on the Tibetan Plateau for the past 16 ka cal B.P.—the Tangra Yumco record. *Geochemistry, Geophysics, Geosystems* 17(3):774–790. doi: [10.1002/2015GC006023](https://doi.org/10.1002/2015GC006023)
- Hua Q, Barbetti M, Rakowski AZ. 2013. Atmospheric radiocarbon for the period 1950–2010. *Radiocarbon* 55(4):2059–2072. doi: [10.2458/azu\\_js\\_rc.v55i2.16177](https://doi.org/10.2458/azu_js_rc.v55i2.16177)
- Kirschvink JL. 1980. The least-squares line and plane and the analysis of palaeomagnetic data. *Geophysical Journal International* 62(3):699–718. doi: [10.1111/j.1365-246X.1980.tb02601.x](https://doi.org/10.1111/j.1365-246X.1980.tb02601.x)
- Mellström A, Nilsson A, Stanton T, Muscheler R, Snowball I, Suttie N. 2015. Post-depositional remanent magnetization lock-in depth in precisely dated varved sediments assessed by archaeomagnetic field models. *Earth and Planetary Science Letters* 410:186–196. doi: [10.1016/j.epsl.2014.11.016](https://doi.org/10.1016/j.epsl.2014.11.016)
- Merrill RT, McFadden PL. 2007. Paleomagnetism. In: Gubbins D, Herrero-Bervera E, editors. *Encyclopedia of geomagnetism and paleomagnetism*. Dordrecht: Springer Netherlands. p. 776–780.
- Mischke S, Rajabov I, Mustaeva N, Zhang C, Herzsuh U, Boomer I, Brown ET, Andersen N, Myrbo A, Ito E, Schudack ME. 2010. Modern hydrology and late Holocene history of Lake Karakul, eastern Pamirs (Tajikistan): a reconnaissance study. *Palaeogeography, Palaeoclimatology, Palaeoecology* 289(1–4):10–24. doi: [10.1016/j.palaeo.2010.02.004](https://doi.org/10.1016/j.palaeo.2010.02.004)
- Nilsson A, Suttie N, Hill MJ. 2018. Short-term magnetic field variations from the post-depositional remanence of lake sediments. *Frontiers in Earth Science* 6. doi: [10.3389/feart.2018.00039](https://doi.org/10.3389/feart.2018.00039)
- Ohlendorf C, Fey M, Massaferro J, Haberzettl T, Laprida C, Lücke A, Maidana N, Mayr C, Oehlerich M, Ramón Mercu J, et al. 2014. Late Holocene hydrology inferred from lacustrine sediments of Laguna Cháltel (southeastern Argentina). *Palaeogeography, Palaeoclimatology, Palaeoecology* 411:229–248. doi: [10.1016/j.palaeo.2014.06.030](https://doi.org/10.1016/j.palaeo.2014.06.030)
- Ólafsdóttir S, Geirsdóttir Á, Miller GH, Stoner JS, Channell JE. 2013. Synchronizing Holocene lacustrine and marine sediment records using paleomagnetic secular variation. *Geology* 41(5):535–538. doi: [10.1130/G33946.1](https://doi.org/10.1130/G33946.1)
- Reimer PJ, Austin WEN, Bard E, Bayliss A, Blackwell PG, Bronk Ramsey C, et al. 2020. The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon* 62:725–757. doi: [10.1017/RDC.2020.41](https://doi.org/10.1017/RDC.2020.41)

- Reinwarth B, Franz S, Baade J, Haberzettl T, Kasper T, Daut G, Helmschrot J, Kirsten KL, Quick LJ, Meadows ME, Mäusbacher R. 2013. A 700-year record on the effects of climate and human impact on the southern cape coast inferred from lake sediments of Eilandvlei, Wilderness embayment, South Africa. *Geografiska Annaler: Series A, Physical Geography* 95(4):345–360. doi: [10.1111/geoa.12015](https://doi.org/10.1111/geoa.12015)
- Roberts AP, Tauxe L, Heslop D. 2013. Magnetic paleointensity stratigraphy and high-resolution Quaternary geochronology: successes and future challenges. *Quaternary Science Reviews* 61:1–16. doi: [10.1016/j.quascirev.2012.10.036](https://doi.org/10.1016/j.quascirev.2012.10.036)
- Roberts AP, Turner G.M. 2013. Geomagnetic excursions and secular variations. In: *Encyclopedia of Quaternary Science*. Elsevier. p. 705–720.
- Roberts AP, Winklhofer M. 2004. Why are geomagnetic excursions not always recorded in sediments? Constraints from post-depositional remanent magnetization lock-in modelling. *Earth and Planetary Science Letters* 227(3–4):345–359. doi: [10.1016/j.epsl.2004.07.040](https://doi.org/10.1016/j.epsl.2004.07.040)
- Ruff M, Fahrni S, Gäggeler HW, Hajdas I, Suter M, Synal H-A, Szidat S, Wacker L. 2010. On-line radiocarbon measurements of small samples using Elemental Analyzer and MICADAS gas ion source. *Radiocarbon* 52:1645–1656. doi: [10.1017/S00382220005637X](https://doi.org/10.1017/S00382220005637X)
- Rudaya N, Nazarova L, Frolova L, Palagushkina O, Soenov V, Cao X, Syrykh L, Grekov I, Otgonbayar D, Bayarkhuu B. 2021. The link between climate change and biodiversity of lacustrine inhabitants and terrestrial plant communities of the Uvs Nuur Basin (Mongolia) during the last three millennia. *Holocene* 31(9):1443–1458. doi: [10.1177/09596836211019093](https://doi.org/10.1177/09596836211019093)
- Salazar G, Zhang YL, Agrios K, Szidat S. 2015. Development of a method for fast and automatic radiocarbon measurement of aerosol samples by online coupling of an Elemental Analyzer with a MICADAS AMS. *Nuclear Instruments and Methods in Physics Research B* 361:163–167. doi: [10.1016/j.nimb.2015.03.051](https://doi.org/10.1016/j.nimb.2015.03.051)
- Scheidt S, Lenz M, Egli R, Brill D, Klug M, Fabian K, Lenz MM, Gromig R, Rethemeyer J, Wagner B, Federov G, Melles M. 2022. A 62 kyr geomagnetic palaeointensity record from the Taymyr Peninsula, Russian Arctic. *Geochronology* 4(1):87–107. doi: [10.5194/gchron-4-87-2022](https://doi.org/10.5194/gchron-4-87-2022)
- Snowball I, Mellström A, Ahlstrand E, Haltia E, Nilsson A, Ning W, Muscheler R, Brauer A. 2013. An estimate of post-depositional remanent magnetization lock-in depth in organic rich varved lake sediments. *Global and Planetary Change* 110:264–277. doi: [10.1016/j.gloplacha.2013.10.005#](https://doi.org/10.1016/j.gloplacha.2013.10.005#)
- Stockhausen H. 1998. Geomagnetic palaeosecular variation (0–13 000 yr BP) as recorded in sediments from three maar lakes from the West Eifel (Germany). *Geophysical Journal International* 135(3):898–910. doi: [10.1046/j.1365-246X.1998.00664.x](https://doi.org/10.1046/j.1365-246X.1998.00664.x)
- Stoner JS, St-Onge G. 2007. Chapter three magnetic stratigraphy in paleoceanography: reversals, excursions, paleointensity, and secular variation. *Developments in Marine Geology* 1:99–138.
- Strobel P, Struck J, Bazarradnaa E, Zech M, Zech R, Bliedner M. 2022. Precipitation and lake water evaporation recorded by terrestrial and aquatic n-alkane  $\delta^2\text{H}$  isotopes in Lake Khar Nuur, Mongolia. *Geochemistry, Geophysics, Geosystems* 23(2). doi: [10.1029/2021GC010234](https://doi.org/10.1029/2021GC010234)
- Strobel P, Struck J, Zech R, Bliedner M. 2021. The spatial distribution of sedimentary compounds and their environmental implications in surface sediments of Lake Khar Nuur (Mongolian Altai). *Earth Surface Processes Landforms* 46(3):611–625. doi: [10.1002/esp.5049](https://doi.org/10.1002/esp.5049)
- Struck J, Bliedner M, Strobel P, Taylor W, Biskop S, Plessen B, Klaes B, Bittner L, Jamsranjav B, Salazar G, et al. 2022. Central Mongolian lake sediments reveal new insights on climate change and equestrian empires in the Eastern Steppes. *Scientific Reports* 12(1). doi: [10.1038/s41598-02206659-w](https://doi.org/10.1038/s41598-02206659-w)
- Sun A, Feng Z, Ran M, Zhang C. 2013. Pollen-recorded bioclimatic variations of the last ~22,600 years retrieved from Achit Nuur core in the western Mongolian Plateau. *Quaternary International* 311:36–43. doi: [10.1016/j.quaint.2013.07.002](https://doi.org/10.1016/j.quaint.2013.07.002)
- Szidat S, Salazar GA, Vogel E, Battaglia M, Wacker L, Synal H-A, Türlér A. 2014.  $^{14}\text{C}$  analysis and sample preparation at the new Bern Laboratory for the Analysis of Radiocarbon with AMS (LARA). *Radiocarbon* 56:561–566. doi: [10.2458/56.17457](https://doi.org/10.2458/56.17457)
- Tauxe L, Steindorf JL, Harris A. 2006. Depositional remanent magnetization: toward an improved theoretical and experimental foundation. *Earth and Planetary Science Letters* 244(3–4):515–529. doi: [10.1016/j.epsl.2006.02.003](https://doi.org/10.1016/j.epsl.2006.02.003)
- Tauxe L, Kodama KP, Kent DV. 2008. Testing corrections for paleomagnetic inclination error in sedimentary rocks: a comparative approach. *Physics of the Earth and Planetary Interiors* 169:152–165. doi: [10.1016/j.pepi.2008.05.006](https://doi.org/10.1016/j.pepi.2008.05.006)
- Unkelbach J, Kashima K, Enters D, Dulamsuren C, Punsalpaamuu G, Behling H. 2019. Late Holocene (Meghalayan) palaeoenvironmental evolution inferred from multi-proxy-studies of lacustrine sediments from the Dayan Nuur region of Mongolia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 530:1–14. doi: [10.1016/j.palaeo.2019.05.021](https://doi.org/10.1016/j.palaeo.2019.05.021)
- Valet J-P, Tanty C, Carlut J. 2017. Detrital magnetization of laboratory-redeposited sediments. *Geophysical Journal International* 210(1):34–41. doi: [10.1093/gji/ggx139](https://doi.org/10.1093/gji/ggx139)

- Wacker L, Bonani G, Friedrich M, Hajdas I, Kromer B, Němec M, Ruff M, Suter M, Synal H-A, Vockenhuber C. 2010. MICADAS: routine and high-precision radiocarbon dating. *Radiocarbon* 52:252–262. doi: [10.1017/S0033822200045288](https://doi.org/10.1017/S0033822200045288)
- Walther M, Dashtseren A, Kamp U, Temujin K, Meixner F, Pan CG, Gansukh Y. 2017. Glaciers, Permafrost and Lake Levels at the Tsengel Khairkhan Massif, Mongolian Altai, during the Late Pleistocene and Holocene. *Geosciences* 7(3):73. doi: [10.3390/geosciences7030073](https://doi.org/10.3390/geosciences7030073)
- Willemse NW, Törnqvist TE. 1999. Holocene century-scale temperature variability from west Greenland lake records. *Geology* 27(7):580–584. doi: [10.1130/0091-7613\(1999\)027<0580:hcstvf>2.3.co;2](https://doi.org/10.1130/0091-7613(1999)027<0580:hcstvf>2.3.co;2)
- Wündsche M, Biagioni S, Behling H, Reinwarth B, Franz S, Bierbaß P, Daut G, Mäusbacher R, Haberzettl T. 2014. ENSO and monsoon variability during the past 1.5 kyr as reflected in sediments from Lake Kalimpaa, Central Sulawesi (Indonesia). *Holocene* 24(12):1743–1756. doi: [10.1177/0959683614551217](https://doi.org/10.1177/0959683614551217)
- Yang Y, Ran M, Sun A. 2020. Pollen-recorded bioclimatic variations of the last ~2000 years retrieved from Bayan Nuur in the western Mongolian Plateau. *Boreas* 49(2):350–362. doi: [10.1111/bor.12423](https://doi.org/10.1111/bor.12423)
- Zhao X, Egli R, Gilder SA, Müller S. 2016. Microbially assisted recording of the Earth's magnetic 422 field in sediment. *Nature Communications* 7:10673. doi: [10.1038/ncomms10673](https://doi.org/10.1038/ncomms10673)