

EFFECT OF FOUR WEEKS OF RTMS ON BRAIN ACTIVITY IN OBSESSIVE-COMPULSIVE DISORDERS DURING AN FLANKER TASK: A RANDOMIZED DOUBLE BLIND STUDY

C. Gaudeau-Bosma^{1,2}, D. Januel², N. Jaafari³, L. Mallet¹, A. Pelissolo⁴

¹CR-ICM U 975 / UMRS 975 / UMR 7225, ²URC, G 03, EPS Ville Evrard, Neuilly-sur-Marne, Paris, ³URC, Centre Hospitalier Henri Laborit, INSERM CIC-P 0802, CHU Poitiers, Poitiers, ⁴Service de Psychiatrie Adulte, Hôpital de la Pitié Salpêtrière, AP-HP, UPMC, Paris, France

Currently, 25-30% of Obsessive-compulsive disorder (OCD) patients are considered to be resistant to conventional therapies (cognitive-behavioral therapy and medications including selective inhibitors of reuptake of serotonin and clomipramine). Among alternative therapeutic strategies for these refractory patients, repetitive Transcranial Magnetic Stimulation (rTMS) has been proposed (Greenberg et al., 1997). Few regions were choice for OCD patients and one of them could be a promising target: the Supplementary Motor Area (SMA). The aim of this project focuses on evaluate the SMA as a promising region to treat OCD patients. Forty patients were include in this double blind study, each received 20 session of rTMS (1 session / day) for 4 weeks with active or sham coil. rTMS parameters consisted of 1500 pulses/d, at 1 Hz, 5 trains of 5 min with an interval inter-stimulus of 2 min and 100% of motor threshold. To localize the SMA, neuronavigation was based on anatomical Magnetic Resonance Imaging (MRI) usingBrainsight software. The therapeutic effect of rTMS will be assessed through a psychological evaluation and a task performed in functional MRI, before and after therapeutic: The Eriksen's task. Its objective is to evaluate cerebral metabolism's modifications during a simple task likely to recruit the error detection system implicating the Anterior Cingulate Cortex without inducing obsessive-compulsive symptoms. This modification were correlated with clinical and cognitive assessment and behavioral results of the Eriksen task.