


Burden of mental disorders by gender in Pakistan: analysis of Global Burden of Disease Study data for 1990–2019

Mohsin Hassan Alvi,¹  Tehmina Ashraf,¹ Farah Naz,¹ Asif Sardar,¹ Akbar Ullah,² Anita Patel,³ Tayyeba Kiran,¹ Anil Gumber,⁴ Nusrat Husain²

BJPsych Bulletin (2023) Page 1 of 8, doi:10.1192/bjb.2023.76

¹Pakistan Institute of Living and Learning, Rawalpindi, Pakistan; ²University of Manchester, Manchester, UK; ³Anita Patel Health Economics Consulting Ltd, London, UK; ⁴Sheffield Hallam University, Sheffield, UK

Correspondence to Mohsin Hassan Alvi (mohsin.h.alvi@pill.org.pk)

First received 23 Feb 2023, final revision 31 Jul 2023, accepted 17 Aug 2023

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Aims and method We aimed to examine the burden of mental disorders in Pakistan over the past three decades. We used the crude data of disability-adjusted life-years (DALYs) obtained from the Global Burden of Disease Study database (1990–2019) to represent burden. Data were retrieved on 26 January 2021. Data for adults of reproductive age (aged 15–49 years) were analysed to discuss and interpret the disease burden. An analysis was conducted on total DALYs separately for the genders for ten mental disorders reported in Pakistan.

Results DALYs increased drastically with the onset of reproductive age. Depressive disorder was the most reported mental disorder, contributing 3.13% (95% CI 2.25–4.24) of total DALYs, and varied significantly between genders: females 3.89% (95% CI 2.73–5.29) versus males 2.37% (95% CI 1.62–3.25).

Clinical implications A nationwide high-quality epidemiological surveillance system should be implemented to monitor mental disorders and offer culturally appropriate preventive services.

Keywords Burden; mental disorder; reproductive age; gender; Pakistan.

Mental, neurological and substance use disorders (MNS) are common, highly disabling and significantly associated with premature death. These disorders are the second leading contributor to total disability-adjusted life-years (DALYs) among males (48.1%) and the leading contributor among females (51.9%).¹ According to the World Health Organization (WHO), around 13% of patients are affected by mental or neurological disorders across the globe.² Among MNS, common mental disorders are one of the leading contributors to burden of disease globally,³ responsible for 22.9% of global years lived with disability (YLDs) and 7.4% of global DALYs, and hence are the leading cause of YLDs and the fifth leading contributor to DALYs.⁴ Eighty per cent of people with mental disorders live in low- and middle-income countries (LMICs).³ Moreover, mental disorders are associated with huge economic costs, more than the cost of long-term physical health conditions such as diabetes or cancer.⁵ According to the WHO, mental disorders result in economic loss of more than US\$1 trillion per year.⁶ These disorders have been prioritised in global health policy and well-being is now included as one of the United Nation's sustainable development goals.⁷ Given the high DALYs associated with common mental disorders there is a great need to establish a national surveillance system in Pakistan to address the knowledge gap and monitor these

illnesses; such a system should include national mental health surveys based on paper and pencil and telephone interviews, personal interviews and health and nutrition examination, pregnancy risk assessment, ambulatory medical care, hospital discharge and admission, economic conditions and poverty.⁸

In South Asians, mental disorders account for 12.2% of total health problems.⁹ A review of studies from the South Asian region suggested high prevalence rates for mental disorders, including depression, anxiety, mood disorders, suicidal behaviour and self-harm, schizophrenia, substance use disorders, neurodevelopmental disorders, dementia and other mental health conditions.¹⁰ The prevalence of mental disorders in Pakistan is as high as 10%, affecting approximately 20 million Pakistanis, and mental illness is associated with huge economic burden.¹¹ Because of financial difficulties faced by most families, these disorders are difficult to manage and treat.¹²

Mental health is deeply rooted in the social, cultural, religious, spiritual, historical and holistic aspects of human lives.¹³ The risk of developing mental disorders is high among the poor, homeless, the unemployed, persons with low educational status, migrants and refugees, and indigenous populations.¹⁴ Evidence from Pakistan indicates that the risk factors for developing mental disorders are gender

(women are at higher risk),¹⁵ poverty,^{16,17} domestic violence,¹⁷ adverse childhood experiences,^{18,19} lack of social support, stressful life events^{18,20} and low educational status.²¹ Furthermore, restrictions in terms of lockdown and social distancing in response to the COVID-19 pandemic have led to economic recession, which is reflected in an increased rate of mental health conditions, including self-harm and suicide.²²

Despite the high prevalence of mental disorders, evidence on their economic burden in Pakistan is limited.²³ Therefore, the purpose of this study was to use data from the Global Burden of Diseases, Injuries and Risk Factors Study 2019 (GBD 2019)²⁴ to assess the risk factors associated with different mental disorders in adults of reproductive age and to further examine burden of disease in Pakistan from 1990 to 2019 in the context of health economics modelling.

Pakistan has one of the highest mental illness rates in the world.²⁵ According to WHO estimates, 24 million people in Pakistan require mental healthcare.²⁶ Unfortunately, the country has only 0.19 psychiatrists for every 100 000 people.²⁷ Highlighting the estimated mental health burden will advocate to policymakers to include the discussion in national health agenda. GBD is the most comprehensive database that reports mental disorders that are not reported in national reports or elsewhere.

The word 'burden' has been used in this study to reflect the technical language commonly used by the authors of GBD studies. There is no intention to express any negative connotation towards people experiencing mental health difficulties.

Method

Data sources

We analysed the GBD 2019 data-set,²⁴ which is a collaborative, comprehensive research study conducted by a global network of more than 3600 researchers from more than 145 countries. GBD 2019 measured morbidity and mortality from 369 diseases and injuries, analysing 87 risk factors in 204 countries and territories.²⁴ GBD 2019 used all available data for mortality, population, fertility, cause of death, incidence, prevalence and other epidemiological measures. To harmonise heterogeneous data and use all possible sources of data, GBD 2019 used a wide variety of statistical modelling methods to build estimates for all outcomes of interest across 204 countries and territories, including Pakistan. GBD 2019 generated a complete set of estimates for cause-specific mortality, years of life lost (YLLs), YLDs and DALYs from 1990 to 2019.

Mental disorders were discussed in GBD 2019 on the basis of the clinical diagnostic criteria from DSM-IV-TR²⁸ or ICD-10.²⁹ YLL refers to premature death in relation to the expected lifespan, measured in years.³⁰ YLDs are calculated from the product of prevalence and weightage of disability for mental health condition with adjustments for comorbidities. DALYs are the sum of premature mortality (YLLs) and loss of productivity (YLDs).⁴

For this study, we downloaded GBD 2019 statistics from the freely accessible resources on the Global Health Data

Exchange (GHDx) (ghdx.healthdata.org/) and the GBD Results Tool (vizhub.healthdata.org/gbd-results/) with additional insights taken from corresponding data visualisations (vizhub.healthdata.org/gbd-compare/). These resources are hosted by the Institute of Health Metrics and Evaluation at the University of Washington. Specifically, we obtained GBD summary results (deaths, DALYs, YLLs and YLDs) for all causes of mental illness at all levels between 1990 and 2019 via the GBD Results Tool (Fig. 1).

The downloaded data for DALYs from GBD 2019 were put into our analysis. All categories of mental disorder gathered by GBD 2019 were selected. We focused on adults of both genders of reproductive age (aged 15–49 years) because we estimated that a large proportion of total disability is associated with this age group (Fig. 2). We gathered crude annual data for Pakistan for the full GBD 2019 study period (1990 to 2019). Data on YLLs and mortality were not considered in this study. Hence, almost all the proportion of DALYs has been contributed by YLDs.

Analysis

Data were cleaned to remove typographical errors and duplicate values and manually checked to confirm authenticity prior to using them for further statistical analyses. Data were presented in tables and graphs, alongside relevant descriptive statistics. Analyses based on a previous study³¹ were carried out, with the current situation in Pakistan kept under consideration. The categories of mental disorders presented in the GBD 2019 tool were reported, along with subcategories. The gender-specific prevalence of each mental disorder and DALY rates in 2019 for Pakistan were reported. The prevalence rates of mental disorders from 1990 to 2019 for both the genders were reported, along with a gender comparison of change in DALY rate for Pakistan. The DALYs for specific mental disorders that were attributable to potential risk factors (such as bullying victimisation, intimate partner violence, childhood




Fig. 1 Data extraction flowchart. IHME, Institute for Health Metrics and Evaluation; GBD 2019, Global Burden of Diseases, Injuries and Risk Factors Study 2019; DALYs, disability-adjusted life-years.




Fig. 2 Age distribution of the population suffering with mental disorder in Pakistan, 2019.

sexual abuse and lead exposure) in 2019 were also presented. The age-standardised rates were based on the GBD global reference population. In total, 18 estimates were reported, with 95% confidence intervals. Charts were supported by bar charts and pie charts for better understanding of results.

Ethics statement

Data gathered from the Institute for Health Metrics and Evaluation's GBD 2019 data-set were presented fairly in analysis. No amendments or alterations were made to crude data, although data were checked for errors and duplication before analysis. The authors do not claim the ownership of data. All the tables and graphs presented in this paper were prepared by the authors. The economic outcomes of the data were disseminated honestly and are accessible to the lay public for awareness, critical appraisal and further research.

Results

The greatest contribution to DALYs due to mental disorders in Pakistan was from depressive disorders (3.13%, 95% CI 2.25–4.24) and major depressive disorder (2.57%, 95% CI 1.78–3.53), followed by anxiety disorders (1.7%, 95% CI 1.21–2.30), schizophrenia (0.97%, 95% CI 0.68–1.25), dysthymia (0.57%, 95% CI 0.39–0.79) and bipolar disorder (0.46%, 95% CI 0.29–0.68); attention-deficit hyperactivity disorder (ADHD; 0.039%, 95% CI 0.022–0.063) and anorexia nervosa (0.06%, 95% CI 0.036–0.096) made the least contribution to DALYs. Depressive disorders and major depressive disorder contributed to DALYs over the 20-year period for both males and females, but females showed a relatively higher burden than males. Anxiety was the second highest contributor to mental illness. The contribution of depressive disorders and eating disorders to total DALYs was substantially higher in females than in males, whereas the contribution of autism spectrum disorders and ADHD was significantly higher in males than in females (Table 1).

Childhood sexual abuse, intimate partner violence and bullying were the factors contributing to depressive and anxiety disorders. Although childhood sexual abuse and bullying victimisation also contributed to DALYs, intimate partner violence was the major risk factor, accounting for 11% of DALYs (Table 2). The proportion of DALYs contributed by childhood sexual abuse was 5.3% in females, which was significantly higher than in males (3.97%). Bullying victimisation accounted for 12.2% of DALYs for anxiety disorder. A large proportion of DALYs for developmental intellectual disability (65.85%) was attributable to lead exposure (Fig. 3).

Discussion

The analysis of data from GBD 2019 shows that the overall proportion of DALYs in Pakistan due to mental disorders increased between 1990 and 2019. The pattern and magnitude of mental disorders changed over that period with respect to gender. Several risk factors have been identified that are likely to contribute to the burden of mental disorders in Pakistan. Findings also highlighted that young and middle-aged adults (aged 15–49 years) are experiencing a higher burden of mental disorders than children and older adults.

Depression and anxiety disorders

According to the WHO, more than 322 million people globally are living with depression and rates increased between 2005 to 2015.³² The prevalence of depression is particularly high in South Asia (27%). The WHO has ranked depression as the single largest contributor to global disability, and anxiety disorders are ranked sixth.³² Similar trends have been reflected in the current study, with depression as the greatest contributor to DALYs in Pakistan, followed by anxiety disorders. These findings highlight the need for universal screening for depression and developing and testing culturally appropriate psychosocial interventions.³³ Such interventions are hugely important, considering that depression is a major risk factor for self-harm and suicide.³²

Table 1 Proportion of disability-adjusted life-years (DALYs) attributable to each mental disorder in Pakistan, 2019

Causes	Both genders, % (95% CI)	Males, % (95% CI)	Females, % (95% CI)
Depressive disorder	3.13 (2.25–4.24)	2.37 (1.62–3.25)	3.89 (2.73–5.29)
Major depressive disorder	2.57 (1.78–3.53)	1.89 (1.23–2.68)	3.25 (2.20–4.53)
Dysthymia	0.57 (0.39–0.79)	0.49 (0.32–0.69)	0.64 (0.44–0.92)
Anxiety	1.70 (1.21–2.30)	1.49 (1.03–2.09)	1.92 (1.34–2.65)
Schizophrenia	0.97 (0.68–1.25)	1.14 (0.80–1.49)	0.81 (0.57–1.07)
Bipolar disorder	0.46 (0.29–0.68)	0.50 (0.30–0.75)	0.42 (0.26–0.62)
Conduct disorder	0.21 (0.12–0.34)	0.30 (0.17–0.48)	0.13 (0.07–0.24)
Intellectual disability	0.37 (0.20–0.59)	0.47 (0.25–0.75)	0.28 (0.15–0.45)
Autism spectrum disorder	0.19 (0.13–0.26)	0.28 (0.19–0.39)	0.10 (0.07–0.14)
Eating disorder	0.21 (0.13–0.30)	0.16 (0.10–0.24)	0.25 (0.16–0.37)
Anorexia nervosa	0.06 (0.04–0.10)	0.03 (0.02–0.06)	0.09 (0.05–0.14)
Bulimia nervosa	0.15 (0.09–0.22)	0.13 (0.07–0.20)	0.17 (0.10–0.25)
ADHD	0.04 (0.02–0.06)	0.06 (0.03–0.10)	0.02 (0.01–0.03)
Other mental disorder	0.45 (0.29–0.65)	0.55 (0.35–0.83)	0.35 (0.22–0.53)

ADHD, attention-deficit hyperactivity disorder.

Gender

The burden of mental disorders was found to be higher in females than in males. This is consistent with global trends.³² Women globally have reported significantly poorer health than men in terms of health-related quality of life, depression and psychological distress.³⁴ Several risk factors, such as women's low self-esteem, high rates of stressful life events, interpersonal violence, discrimination and lack of gender equality, make women more vulnerable to developing mental health problems.³⁵ Women during the reproductive years are particularly vulnerable to developing mental disorders; a study in Nigeria reported that prevalence of mental disorders (depression and anxiety) is 60.7% among women of reproductive age.³⁶ Evidence supports that both genes and gene–environment interactions contribute to the risk of depression in a gender-specific manner.³⁶ Therefore, strategies to incorporate mental health into reproductive health services may contribute to reducing this high burden. A series paper in the *Lancet* reported that Pakistan had the highest rate of stillbirths (43.1 per 1000 total births) across the globe.³⁷ There is also a need to promote public health interventions to target women who have gynaecological conditions and who have experienced miscarriage/stillbirth to effectively reduce the burden of mental health problems in this age group.³⁸

Adverse childhood experiences and intimate partner violence

In the current study childhood sexual abuse, intimate partner violence and bullying were the major factors contributing to depressive and anxiety disorders. Studies have shown that childhood trauma is associated with a range of adverse physical, psychological and social outcomes.³⁸ Individuals with a history of childhood abuse are significantly more likely to develop major depressive disorder in

adulthood.³⁹ A meta-analysis suggests that approximately 46% of individuals with depression reported childhood abuse, which is also associated with an earlier onset, a chronic course and treatment resistance for depression.⁴⁰ Recent evidence from Pakistan shows that the majority of women with depression (58%) had had an adverse childhood experience such as home violence and neglect.^{19,41} Bullying is another type of childhood traumatic experience that is a major psychosocial concern because of its implications for subsequent behavioural and mental maladjustments over time. Individuals with history of childhood bullying are at increased risk of anxiety and depressive symptoms as well as future bullying and victimisation.⁴² These findings highlight the need for translational research that could potentially help in exploring strategies that might play a role in dealing with the impact of childhood abuse and bullying, particularly in individuals with major depressive disorder. Individualised culturally appropriate evidence-based psychosocial interventions that specifically address this subgroup are needed.⁴³ Digitally delivered psychosocial interventions such as TechMotherCare⁴⁴ may help to improve the mental health of the general population.⁴⁵

Intimate partner violence has also been reported as a risk factor for mental disorders (depression and anxiety) in this study. Intimate partner violence is a significant human rights issue and global health concern, particularly in LMICs, with numerous detrimental outcomes, including intimate partner homicide⁴⁶ and suicide.⁴⁷ According to the WHO's global estimates, on average, up to 852 million women (almost 1 in 3 women) aged 15 years or older in 2018 had experienced intimate partner violence at least once in their lifetime.⁴⁸ The prevalence of intimate partner violence in South Asia is the highest, at 35%.⁴⁵ The Pakistan Demographic and Health Survey for 2012–2013 reported that 34% of ever-married women had experienced spousal physical, sexual or emotional violence, and 7% of women who had ever been pregnant had experienced




Fig. 3 Comparison of disability-adjusted life-years (DALYs) by gender in Pakistan for 1990 and 2019.

Mental disorder	Risk factor	Proportion of total DALYs attributable to each risk factor, % (95% CI)		
		Both genders	Males	Females
Idiopathic developmental disorder	Lead exposure	65.85 (35.2–84.19)	66.57 (35.91–84.85)	64.69 (33.65–83.16)
Anxiety disorder	Bullying victimisation	12.20 (3.19–24.45)	14.07 (3.73–28.06)	10.81 (2.85–21.89)
Depressive disorder	Intimate partner violence	11.00 (0.07–22.39)	Not applicable	17.49 (0.10–35.76)
Depressive disorder	Bullying victimisation	7.06 (1.14–14.68)	8.15 (1.65–16.91)	6.42 (1.23–13.40)
Depressive disorder	Childhood sexual abuse	4.80 (2.51–7.92)	3.97 (1.78–7.23)	5.30 (1.78–7.23)

DALYs, disability-adjusted life-years.

violence during pregnancy.⁴⁹ In addition, the containment measures introduced by the government as a result of the COVID-19 pandemic led to exacerbation of psychological adversities, including increased rates of intimate partner violence.⁵⁰ Reducing gender-based violence is an indicator of one of the United Nation's sustainable development goals.⁵¹ Research conducted in high-income countries shows that mental health interventions may reduce the risk for victimisation by treating mental health problems and empowering women.⁵² However, the evidence from LMICs such as Pakistan is limited.⁵³

Addressing the burden of mental disorders: information and intervention

Mental health is being neglected in national surveys of Pakistan,^{54–56} resulting in limited robust data as well as contradictions in reports. Providing national level high-quality information is a first milestone towards strengthening the mental health surveillance system. This system will play a vital role in monitoring and reporting the national mental health burden and how much resources are needed to cover this burden. The accurate prevalence of diagnosable mental, neurological and substance use disorders (MNS) in Pakistan remains unclear⁵⁷ because of limited high-quality research evidence on mental health.⁵⁸ In addition to limited research, there are also problems at service delivery level because of limited trained staff to meet the needs of the population, contributing to the treatment gap. One of the recent initiatives to improve the health and well-being of people in Pakistan is the 'President's Programme to Promote Mental Health of Pakistanis' launched in 2019.⁵⁹ Based on the recommendations made by the Lancet Commission on Global Mental Health, the programme emphasises the potential role of early-life interventions that promote mental health and well-being and prevent MNS, and it calls for implementation of the WHO Thinking Healthy Programme for mothers⁶⁰ and the WHO School Mental Health Programme adapted for Pakistan.⁶¹ A similar programme, learning through play (LTP) plus CBT, is currently being implemented across Karachi city, which has a population of 23 million.⁴¹ The National Health Vision of Pakistan 2016–2025 is intended to focus on improving health facilities, with a particular focus on communicable and non-communicable diseases, including mental illness.⁶² The health minister of Sindh province of Pakistan has taken an initiative to implement a culturally adapted improving access to psychological therapy programme (IAPT-PK) in Dadu and Qambar Shahdadkot districts, which are most affected by recent floods, in order to treat mental health conditions among the population.⁶³

Limitations

The limitations of the GBD methods, including those for estimation of MNS, have been documented in published literature. They include underestimation of the total uncertainty for DALYs; time difference in the reporting of health data by countries and their subsequent incorporation into the GBD estimations; and the scarcity of data for particular

locations, leading to wider uncertainty intervals.⁵¹ A major limitation of this secondary analysis for Pakistan is that accurate and reliable population-level data on the prevalence of MNS are very limited across Pakistan, which may have created unknown biases in estimates reported in this paper. An estimation of the prevalence of common mental disorders has been reported in this study using data gathered from the GBD data-set that might vary from actual figures as the context is highly dynamic. The GBD estimation of burden of MNS relies on severity distribution data from high-income countries, which may not be a true reflection of the distribution in LMICs such as Pakistan.

About the authors

Mohsin Hassan Alvi is a research fellow, Department of Research, Pakistan Institute of Living and Learning, Rawalpindi, Pakistan. **Tehmina Ashraf** is a research assistant, Department of Research, Pakistan Institute of Living and Learning, Rawalpindi, Pakistan. **Farah Naz** is a research fellow, Department of Research, Pakistan Institute of Living and Learning, Rawalpindi, Pakistan. **Asif Sardar** is a research fellow, Department of Research, Pakistan Institute of Living and Learning, Rawalpindi, Pakistan. **Akbar Ullah** is a research fellow, Division of Population Health, Health Services Research & Primary Care, University of Manchester, Manchester, UK. **Anita Patel** is Director of Anita Patel Health Economics Consulting Ltd, London, UK. **Tayyeba Kiran** is Assistant Director, Department of Research, Pakistan Institute of Living and Learning, Rawalpindi, Pakistan. **Anil Gumber** is Principal Health Economist, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, UK. **Nusrat Husain** is Professor of Psychiatry and Director of Research for Global Mental Health, Division of Psychology & Mental Health, University of Manchester, Manchester, UK.

Data availability

The data that support the findings of this study are openly available on the Global Health Data Exchange at vizhub.healthdata.org/gbd-results.

Acknowledgements

We thank the Institute for Health Metrics and Evaluation (GBD 2019) at the University of Washington for providing access to data for Pakistan and the reviewers and supervisors for their support.

Author contributions

M.H.A. and T.A. developed the concept for the article. M.H.A. and A.S. designed the model and performed acquisition, analysis and interpretation of data. T.A., F.N. and T.K. contributed to the interpretation of the results and writing the manuscript. A.P., A.G. and A.U. agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. N.H. approved the version to be published. All authors reviewed the results and approved the final version of the manuscript.

Funding

This research has been supported through a grant from the Medical Research Council, the UK Department for International Development and the National Institute for Health and Care Research (MR/R022461/1) for the Youth Culturally Adapted Manual Assisted Problem Solving Training (YCMAP) project. The funders were not involved in the preparation of this paper.

Declaration of interest

None.

References

- Patel V, Chisholm D, Parikh R, Charlson FJ, Degenhardt L, Dua T, et al. Global priorities for addressing the burden of mental, neurological, and substance use disorders. In (eds V Patel V, D Chisholm, T Dua, R Laxminarayan, ME Medina-Mora ME): 1–27. *Disease Control Priorities, Third Edition: Volume 4 Mental, Neurological, and Substance Use Disorders*. International Bank for Reconstruction and Development/The World Bank, 2016.
- Faruqi SJ, Shahbaz NN, Nisa Q, Umer SR, Ali SG, Aziz MY. Cost of investigating neurological disease: experience of a tertiary care center in Karachi, Pakistan. *Cureus* 2020; **12**(7): e9291.
- Bonilla-Escobar FJ, Lim HM. A call for action for mental health: medical students and physicians' roles. *International Journal of Medical Students*, 2015; **3**(3): 121–2.
- Whiteford HA, Ferrari AJ, Degenhardt L, Feigin V, Vos T. Global burden of mental, neurological, and substance use disorders: an analysis from the Global Burden of Disease Study 2010. *PLoS One* 2015; **10**(2): e0116820.
- Trautmann S, Rehm J, Wittchen HU. The economic costs of mental disorders: do our societies react appropriately to the burden of mental disorders? *EMBO Rep* 2016; **17**: 1245–9.
- World Health Organization. *Making the Investment Case for Mental Health: A WHO/UNDP Methodological Guidance Note*. World Health Organization, 2019 (<https://apps.who.int/iris/bitstream/handle/10665/325116/WHO-UHC-CD-NCD-19.97-eng.pdf>).
- United Nations. *The Sustainable Development Goals Report 2016*. United Nations, 2016.
- Reeves WC, Pratt LA, Thompson W, Ahluwalia IB, Dhingra SS, McKnight-Eily LR, et al. Mental illness surveillance among adults in the United States. *MMWR* 2011; **60**: 1–29.
- Ranjan JK, Asthana HS. Prevalence of mental disorders in India and other South Asian countries. *Asian J Epidemiol* 2017; **10**: 45–53.
- Hossain MM, Purohit N, Sultana A, Ma P, McKyer EL, Ahmed HU. Prevalence of mental disorders in South Asia: an umbrella review of systematic reviews and meta-analyses. *Asian J Psychiatry* 2020; **51**: 102041.
- Nisar M, Mohammad RM, Fatima S, Shaikh PR, Rehman M. Perceptions pertaining to clinical depression in Karachi, Pakistan. *Cureus* 2019; **11**(7): e5094.
- Khalily MT, ur Rehman A, Bhatti MM, Hallahan B, Ahmad I, Mehmood MI, et al. Stakeholders' perspective on mental health laws in Pakistan: a mixed method study. *Int J Law Psychiatry* 2021; **74**: 101647.
- Gopalkrishnan N. Cultural diversity and mental health: considerations for policy and practice. *Front Public Health* 2018; **6**: 179.
- Caddick H, Horne B, Mackenzie J, Tilley H. *Investing in Mental Health in Low-Income Countries (ODI Insights)*. Overseas Development Institute, 2016.
- Khan N, Amjad A, Farooq A, Kausar R. Gender differences in common mental disorders. *Pak J Clin Psychol* 2016; **15**: 21–32.
- Kazmi SM. How Does Socio-Economic Factors Force Children into Child Labour? A Case Study of Sahiwal District, Punjab, Pakistan (Working Paper 150). Sustainable Development Policy Institute, 2015.
- Shagufta S, Shams S. Prevalence, differences, and predictors of anxiety and depression among pregnant and Non-pregnant women in Peshawar Khyber Pakhtunkhwa Pakistan. *FWU J Soc Sci* 2019; **13**: 167–76.
- Husain MI, Umer M, Chaudhry IB, Husain MO, Rahman R, Shakoor S, et al. Relationship between childhood trauma, personality, social support and depression in women attending general medical clinics in a low and middle-income country. *J Affect Disord* 2021; **292**: 526–33.
- LeMasters K, Bates LM, Chung EO, Gallis JA, Hagaman A, Scherer E, et al. Adverse childhood experiences and depression among women in rural Pakistan. *BMC Public Health* 2021; **21**(1): 400.
- Husain N, Parveen A, Husain M, Saeed Q, Jafri F, Rahman R, et al. Prevalence and psychosocial correlates of perinatal depression: a cohort study from urban Pakistan. *Arch Womens Ment Health* 2011; **14**: 395–403.
- Husain N, Chaudhry N, Jafri F, Tomenson B, Surhand I, Mirza I, et al. Prevalence and risk factors for psychological distress and functional disability in urban Pakistan. *WHO S E Asia J Public Health* 2014; **3**: 144–53.
- Mamun MA. Suicide and suicidal behaviors in the context of COVID-19 pandemic in Bangladesh: a systematic review. *Psychol Res Behav Manag* 2021; **14**: 695–704.
- Khan MM. Economic burden of mental illnesses in Pakistan. *J Ment Health Policy Econ* 2016; **19**: 155–66.
- Institute for Health Metrics and Evaluation. *Global Burden of Disease Study 2019 (GBD 2019) Data Resources*. IHME, 2019 (<https://ghdx.healthdata.org/gbd-2019>).
- Sikandar S. A letter from... Pakistan. *Lancet Psychiatry* 2020; **10**: 845.
- Amer M. Mental health: making mental healthcare accessible. *TNS* 2023; 16 Apr (<https://www.thenews.com.pk/tns/detail/1061020-mental-health-making-mental-healthcare-accessible#:~:text=An%20estimated%2024%20million%20people,ranges%20between%20500%20and%20600>).
- World Health Organization. WHO Pakistan celebrates World Mental Health Day (Press Release). WHO, 2021 (<https://www.emro.who.int/pak/pakistan-news/who-pakistan-celebrates-world-mental-health-day.html>).
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders* (4th edn, text revision) (DSM-IV-TR). APA, 2000.
- World Health Organization. *Tenth Revision of the International Classification of Diseases and Related Health Problems (ICD-10)*. WHO, 2009.
- Colton CW, Manderscheid RW. Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Prev Chronic Dis* 2006; **3**(2): A42.
- Pant PR, Banstola A, Bhatta S, Mytton JA, Acharya D, Bhattarai S, et al. Burden of injuries in Nepal, 1990–2017: findings from the Global Burden of Disease Study 2017. *Injury Prev* 2020; **26**(suppl 2): i57–66.
- World Health Organization. *Depression and Other Common Mental Disorders: Global Health Estimates*. WHO, 2017.
- Salk RH, Hyde JS, Abramson LY. Gender differences in depression in representative national samples: meta-analyses of diagnoses and symptoms. *Psychol Bull* 2017; **143**: 783–822.
- Boerma T, Hosseinpoor AR, Verdes E, Chatterji S. A global assessment of the gender gap in self-reported health with survey data from 59 countries. *BMC Public Health* 2016; **16**: 675.
- Kuehner C. Why is depression more common among women than among men? *Lancet Psychiatry* 2017; **4**: 146–58.
- Soyannwo T, Adebayo AM, Sigbeku O. Mental health problems of reproductive age group women in a rural community of south west Nigeria. *J Ment Health* 2018; **29**: 45–51.
- De Bernis L, Kinney MV, Stones W, ten Hoop-Bender P, Vivio D, Leisher SH, et al. Stillbirths: ending preventable deaths by 2030. *Lancet* 2016; **387**: 703–16.
- Nemeroff CB. Paradise lost: the neurobiological and clinical consequences of child abuse and neglect. *Neuron* 2016; **89**: 892–909.
- Humphreys KL, LeMoult J, Wear JG, Piersiak HA, Lee A, Gotlib IH. Child maltreatment and depression: a meta-analysis of studies using the childhood trauma questionnaire. *Child Abuse Negl* 2020; **102**: 104361.
- Nelson J, Klumpparendt A, Doebler P, Ehring T. Childhood maltreatment and characteristics of adult depression: meta-analysis. *Br J Psychiatry* 2017; **210**: 96–104.
- Husain N, Kiran T, Shah S, Rahman A, Saeed Q, Naeem S, et al. Efficacy of learning through play plus intervention to reduce maternal depression in women with malnourished children: a randomized controlled trial from Pakistan. *J Affect Disord* 2021; **278**: 78–84.

- 42 Lee J. Pathways from childhood bullying victimization to young adult depressive and anxiety symptoms. *Child Psychiatry Hum Dev* 2021; **52**: 129–40.
- 43 Struck N, Krug A, Yuksel D, Stein F, Schmitt S, Meller T, et al. Childhood maltreatment and adult mental disorders – the prevalence of different types of maltreatment and associations with age of onset and severity of symptoms. *Psychiatry Res* 2020; **293**: 113398.
- 44 Husain N, Kiran T, Fatima B, Chaudhry IB, Saeed Q, Masood SN, et al. Development and assessment of a mobile phone-based intervention to reduce maternal depression and improve child health. *Eur Psychiatry* 2016; **33**(suppl 1): S608–9.
- 45 Arshad U, Gauntlett J, Husain N, Chaudhry N, Taylor PJ. A systematic review of the evidence supporting mobile- and internet-based psychological interventions for self-harm. *Suicide Life Threat Behav* 2020; **50**: 151–79.
- 46 Petrosky E, Blair JM, Betz CJ, Fowler KA, Jack SP, Lyons BH. Racial and ethnic differences in homicides of adult women and the role of intimate partner violence – United States, 2003–2014. *Morb Mortal Weekly Rep* 2017; **66**(28): 741–6.
- 47 Brown S, Seals J. Intimate partner problems and suicide: are we missing the violence? *J Inj Violence Res* 2019; **11**: 53–64.
- 48 World Health Organization. *Violence Against Women Prevalence Estimates, 2018: Global, Regional and National Prevalence Estimates for Intimate Partner Violence Against Women and Global and Regional Prevalence Estimates for non-Partner Sexual Violence Against Women*. WHO, 2021.
- 49 National Institute of Population Study (NIPS)[Pakistan], ICF International. *Pakistan: Demographic and Health Survey 2012–13*. NIPS/ICF International, 2013.
- 50 Yahya AS, Khawaja S, Chukwuma J. Association of COVID-19 with intimate partner violence. *Prim Care Comp CNS Disord* 2020; **22**(3): 27392.
- 51 Kyu HH, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2018; **392**: 1859–922.
- 52 Keynejad RC, Hanlon C, Howard LM. Psychological interventions for common mental disorders in women experiencing intimate partner violence in low-income and middle-income countries: a systematic review and meta-analysis. *Lancet Psychiatry* 2020; **7**: 173–90.
- 53 Latif M, Husain MI, Gul M, Naz S, Irfan M, Aslam M, et al. Culturally adapted trauma-focused CBT-based guided self-help (CatCBT GSH) for female victims of domestic violence in Pakistan: feasibility randomized controlled trial. *Behav Cogn Psychother* 2021; **49**: 50–61.
- 54 Pakistan Bureau of Statistics. *Pakistan Social & Living Standards Measurement Survey (PSLM) 2018–19: National/Provincial (Social Report)*. PBS, 2020.
- 55 National Institute of Population Study (NIPS)[Pakistan], ICF. *Pakistan: Demographic and Health Survey 2017–18*. NIPS/ICF, 2019.
- 56 Pakistan Bureau of Statistics. *Household Integrated Economic Survey (HIES) 2018–19*. PBS, 2020.
- 57 Khalily MT. Mental health problems in Pakistani society as a consequence of violence and trauma: a case for better integration of care. *Int J Integ Care* 2011; **11**: e128.
- 58 Riaz F. *Psychological Resilience Mechanisms and Mental Health Perspectives of Marginalized Pakistani Communities* (Doctoral Dissertation). Monash University, 2019 (<https://doi.org/10.4225/03/5b2c44401ecf5>).
- 59 Mirza Z, Rahman A. Mental health care in Pakistan boosted by the highest office. *Lancet* 2019; **394**: 2239–40.
- 60 Rahman A, Malik A, Sikander S, Roberts C, Creed F. Cognitive behaviour therapy-based intervention by community health workers for mothers with depression and their infants in rural Pakistan: a cluster-randomised controlled trial. *Lancet* 2008; **372**: 902–9.
- 61 Imran N, Rahman A, Chaudhry N, Asif A. World Health Organization “School Mental Health Manual”-based training for school teachers in urban Lahore, Pakistan: study protocol for a randomized controlled trial. *Trials* 2018; **19**(1): 290.
- 62 Shaikh BT, Hafeez A, Ali N. Pakistan’s health and population think tank: a policy platform for achieving sustainable development goal 3 national targets. *East Mediterr Health J* 2019; **25**: 754–9.
- 63 The Dayspring. Bridging the mental health treatment gap through evidence and policy. *Dayspring* 2022; 25 Nov (<https://www.thedayspring.com.pk/bridging-mental-health-treatment-gap-through-evidence-and-policy/>).

