



COMPOSITIO MATHEMATICA

On exact ∞ -categories and the Theorem of the Heart

Clark Barwick

Compositio Math. **151** (2015), 2160–2186.

[doi:10.1112/S0010437X15007447](https://doi.org/10.1112/S0010437X15007447)





On exact ∞ -categories and the Theorem of the Heart

Clark Barwick

ABSTRACT

The new homotopy theory of *exact ∞ -categories* is introduced and employed to prove a Theorem of the Heart for algebraic K -theory (in the sense of Waldhausen). This implies a new compatibility between Waldhausen K -theory and Neeman K -theory. Additionally, it provides a new proof of the Dévissage and Localization theorems of Blumberg–Mandell, new models for the G -theory of schemes, and a proof of the invariance of G -theory under derived nil-thickenings.

Contents

1	Preliminaries	2162
2	Additive ∞-categories	2164
3	Exact ∞-categories	2165
4	Exact functors between exact ∞-categories	2167
5	Theories and duality	2171
6	Theorem of the Heart	2178
7	Application: Abelian models for the algebraic G-theory of schemes	2181
8	Application: a theorem of Blumberg–Mandell	2182
9	Application: G-theory of spectral Deligne–Mumford stacks	2184
	Acknowledgements	2184
	References	2184

Introduction

In this paper, we prove (Theorem 6.1) that a stable homotopy theory whose triangulated homotopy category admits a bounded t-structure has the same algebraic K -theory (in the sense of Waldhausen) as that of its heart. This is the *Theorem of the Heart* of the title. This result, which apparently has been expected by some experts for some time, has nevertheless gone unproved.

This theorem does, however, have an important predecessor. For 20 years Amnon Neeman has advanced the *algebraic K -theory of triangulated categories* [Nee97a, Nee97b, Nee98a, Nee98b, Nee99, Nee00a, Nee00b, Nee01, Nee05] as a way of extracting K -theoretic data directly from the triangulated homotopy category of a stable homotopy theory. As an approximation to Waldhausen K -theory, this form of K -theory has well-documented limitations: a beautiful example of Schlichting [Sch02] shows that Waldhausen K -theory can distinguish stable homotopy theories with equivalent triangulated homotopy categories. Nevertheless, the most impressive advance in the algebraic K -theory of triangulated categories is Neeman’s *Theorem of the Heart* [Nee98b, Nee99, Nee01], which expresses an equivalence between the Neeman K -theory

Received 25 February 2013, accepted in final form 22 October 2014, published online 19 August 2015.

2010 Mathematics Subject Classification 19D99, 55U40 (primary).

Keywords: algebraic K -theory, stable ∞ -categories, t-structures.

This journal is © Foundation Compositio Mathematica 2015.

of a triangulated category \mathcal{T} equipped with a bounded t-structure and the Quillen K -theory of its heart \mathcal{T}^\heartsuit .

Neeman's proof of his Theorem of the Heart is lengthy and difficult to read, so much so that it even generated a small controversy (see Neeman's discussion in [Nee97a, pp. 347–353]). The proof of our Theorem of the Heart, by contrast, is mercifully short, conceptually appealing, and logically independent of Neeman's. Consequently we regard Theorem 6.1 and its proof as a conclusive answer to Problem 76 of his survey [Nee05].

To prove our result, we introduce a natural homotopy-theoretic generalization of Quillen's notion of an exact category, which we call an *exact ∞ -category* (Definition 3.1). Because this notion involves a compatibility between certain homotopy limits and certain homotopy colimits, it is virtually impossible (or at the very least hideously inconvenient) to express in the more classical language of categories-with-weak-equivalences. Therefore we have to employ concepts from higher category theory; in particular, the theory of *Waldhausen ∞ -categories*, whose theory we studied in pitiless detail in [Bar13].

The key idea from [Bar13] is that algebraic K -theory is a homology theory for ∞ -categories. In fact, algebraic K -theory is the analogue of stable homotopy theory in this context. The behavior of these categorified homology theories under duality is the key phenomenon that makes our proof of the Theorem of the Heart work. More precisely, when algebraic K -theory is restricted to exact ∞ -categories, it enjoys a *self-duality* (Corollary 5.16.1). This self-duality is then used in conjunction with our ∞ -categorical Fibration theorem [Bar13, Proposition 10.12] to prove the following theorem.

THEOREM HEART. *If \mathcal{A} is a stable ∞ -category equipped with a bounded t-structure, then the inclusion of the heart $\mathcal{A}^\heartsuit \subset \mathcal{A}$ induces a K -theory weak equivalence*

$$K(\mathcal{A}^\heartsuit) \simeq K(\mathcal{A}).$$

This result is one of the very few general statements in algebraic K -theory that is capable of providing K -theory equivalences that do not arise from equivalences of the ∞ -categories themselves. (The only other example of such a general result we know of this kind is Quillen's Dévissage theorem.)

The full strength of the conceptual apparatus constructed here and in [Bar13] is necessary for this proof to work. In view of Schlichting's 'no-go theorem' [Sch02, Proposition 2.2], our use of the Fibration theorem makes it impossible for a proof at all similar to the one presented here to be adapted to the context of triangulated categories. On the other hand, there are other proposed versions of algebraic K -theory for ∞ -categories (most notably that of Blumberg *et al.* [BGT13]) that restrict attention to stable ∞ -categories or the like. These versions of K -theory just won't help for this problem: in fact, it isn't possible even to *express* the relevant cases of self-duality with this or any other form of K -theory that splits arbitrary cofiber sequences.

Let us underscore that this is not a new proof of an old theorem. Schlichting's example shows that there is no *a priori* reason to expect Neeman's Theorem of the Heart to say anything about Waldhausen K -theory. Nevertheless, our main result does yield a comparison between Neeman's K -theory and Waldhausen K -theory. Indeed, the conjunction of Neeman's Theorem of the Heart and Theorem 6.1 implies that the Waldhausen K -theory of a stable ∞ -category \mathcal{A} agrees with the Neeman K -theory of its triangulated homotopy category $\mathcal{T} = h\mathcal{A}$ (the variant denoted $K({}^w\mathcal{T})$ in [Nee05]), whenever the latter admits a bounded t-structure (Corollary 6.4.1). This verifies a conjecture of Neeman [Nee99, Conjecture A.5] for such stable homotopy theories.

This paper ends with a discussion of some immediate corollaries of the main theorem, which we include mostly as proof of concept.

- We give new models for the G -theory of schemes in terms of the perverse coherent sheaves of Arinkin, Bezrukavnikov, and Deligne (Example 7.1).
- We also give a new, short proof of the Dévissage and Localization theorems of Blumberg and Mandell [BM08] (Proposition 8.3 and Theorem 8.7), which immediately yields a host of useful fiber sequences in the algebraic K -theory of ring spectra (Example 8.8). More interesting examples can be found in our paper with Lawson [BL14].
- Finally, we show that the G -theory of spectral Deligne–Mumford stacks (in the sense of Lurie) is invariant under derived thickenings (Proposition 9.2).

1. Preliminaries

We use higher categories systematically in this paper. In particular, we are interested in ∞ -categories whose i -morphisms for $i \geq 2$ are all invertible. These days, it is fashionable to call these $(\infty, 1)$ -categories or even just ∞ -categories. There are very many models for the homotopy theory of ∞ -categories in this sense, and they are all equivalent in an essentially unique fashion, up to orientation; see Toën [Toe05] or Lurie [Lur09b] or Barwick and Schommer-Pries [BS11].

In this paper, we employ the homotopy theory of *quasicategories* developed by Joyal [Joy08a, Joy08b] and then further by Lurie [Lur09a]. These are simplicial sets C in which any inner horn $x : \Lambda_k^m \rightarrow C$ ($m \geq 2$ and $1 \leq k \leq m - 1$) admits a filler $\bar{x} : \Delta^m \rightarrow C$. When we use the phrase ‘ ∞ -categories’ in this text, we will be referring to these.

One point that is perhaps not so obvious is the notion of a *subcategory* of an ∞ -category.

RECOLLECTION 1.1 [Lur09a, § 1.2.11]. A *subcategory* of an ∞ -category A is a simplicial subset $A' \subset A$ such that for some subcategory $(hA)'$ of the homotopy category hA , the square

$$\begin{array}{ccc} A' \hookrightarrow & A & \\ \downarrow & & \downarrow \\ N(hA)' \hookrightarrow & N(hA) & \end{array}$$

is a pullback diagram of simplicial sets. In particular, note that a subcategory of an ∞ -category is uniquely specified by specifying a subcategory of its homotopy category. Note also that any inclusion $A' \hookrightarrow A$ of a subcategory is an inner fibration [Lur09a, Definition 2.0.0.3, Proposition 2.3.1.5].

We will say that $A' \subset A$ is a *full subcategory* if $(hA)' \subset hA$ is a full subcategory. In this case, A' is uniquely determined by the set A'_0 of vertices of A' , and we say that A' is *spanned by* the set A'_0 .

We will say that A' is *stable under equivalences* if the subcategory $(hA)' \subset hA$ above can be chosen to be stable under isomorphisms. Note that any inclusion $A' \hookrightarrow A$ of a subcategory that is stable under equivalences is a categorical fibration, i.e., a fibration for the Joyal model structure [Lur09a, Corollary 2.4.6.5].

The natural inputs for algebraic K -theory are what we call Waldhausen ∞ -categories.

RECOLLECTION 1.2 [Bar13, Definition 2.7]. A *Waldhausen ∞ -category* $(\mathcal{C}, \mathcal{C}_\dagger)$ consists of an ∞ -category \mathcal{C} equipped with a subcategory $\mathcal{C}_\dagger \subset \mathcal{C}$ that contains all the equivalences. A morphism of \mathcal{C}_\dagger will be said to be *ingressive* or a *cofibration*. These data are then required to satisfy the following conditions.

(1.2.1) The ∞ -category \mathcal{C} contains a *zero object*; i.e., an object that is both initial and terminal [Lur09a, Definition 1.2.12.1 and Remark 1.2.12.6].

(1.2.2) For any zero object 0 , any morphism $0 \rightarrow X$ is *ingressive*.

(1.2.3) Pushouts [Lur09a, § 4.4.2] of cofibrations exist and are cofibrations.

A functor $\mathcal{C} \rightarrow \mathcal{D}$ between Waldhausen ∞ -categories is said to be *exact* if it carries cofibrations to cofibrations and preserves both zero objects and pushouts of cofibrations.

The examples one may have in mind here include the nerve of an ordinary exact category in the sense of Quillen (in which the ingressive morphisms are the admissible monomorphisms), the nerve of a category with cofibrations in the sense of Waldhausen [Wal85] (in which the ingressive morphisms are the cofibrations), any ∞ -category with a zero object and all finite colimits (in which any morphism is ingressive).

In a sense, the defining property of algebraic K -theory is that it splits cofiber sequences. We'll discuss this point in more detail later. For now, let us explain what cofiber sequences *are*.

DEFINITION 1.3. In a Waldhausen ∞ -category, a *cofiber sequence* is a pushout square

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & X'' \end{array}$$

in which $X' \rightarrow X$ is ingressive and 0 is a zero object. We call $X \rightarrow X''$ the *cofiber* of the cofibration $X' \rightarrow X$.

We also have the dual notion of a coWaldhausen ∞ -category.

RECOLLECTION 1.4 [Bar13, Definition 2.16]. A *coWaldhausen ∞ -category* $(\mathcal{C}, \mathcal{C}^\dagger)$ consists of an ∞ -category \mathcal{C} equipped with a subcategory $\mathcal{C}^\dagger \subset \mathcal{C}$ that contains all the equivalences. A morphism of \mathcal{C}^\dagger will be said to be *egressive* or a *fibration*. These data are then required to satisfy the following conditions.

(1.4.1) The ∞ -category \mathcal{C} contains a zero object.

(1.4.2) For any zero object 0 , any morphism $X \rightarrow 0$ is *egressive*.

(1.4.3) Pullbacks [Lur09a, § 4.4.2] of fibrations exist and are fibrations.

A functor $\mathcal{C} \rightarrow \mathcal{D}$ between coWaldhausen ∞ -categories is said to be *exact* if it carries fibrations to fibrations and preserves both zero objects and pullbacks of fibrations.

In other words, coWaldhausen ∞ -categories are precisely the opposites of Waldhausen ∞ -categories. In fact, Waldhausen and coWaldhausen ∞ -categories and exact functors organize themselves into ∞ -categories \mathbf{Wald}_∞ and \mathbf{coWald}_∞ , and the formation of the opposite ∞ -category restricts to an equivalence of ∞ -categories

$$\mathbf{Wald}_\infty \xrightarrow{\sim} \mathbf{coWald}_\infty$$

[Bar13, Notations 2.13 and 2.17, and Proposition 2.18].

DEFINITION 1.5. In a coWaldhausen ∞ -category, a *fiber sequence* is a pullback square

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & X'' \end{array}$$

in which $X \twoheadrightarrow X''$ is egressive and 0 is a zero object. We call $X' \rightarrow X$ the *fiber* of the fibration $X \twoheadrightarrow X''$.

2. Additive ∞ -categories

In effect, an exact ∞ -category will be a Waldhausen ∞ -category \mathcal{C} that is also a coWaldhausen ∞ -category, with two additional properties: first, fiber sequences and cofiber sequences in \mathcal{C} must coincide; and second, \mathcal{C} must be *additive*.

This notion of additivity is similar to the notion of additivity for ordinary categories. Recall from [Bar13, Definition 4.5] the following.

RECOLLECTION 2.1. Suppose C an ∞ -category. Then C is said to *admit finite direct sums* if the following conditions hold.

- (2.1.1) The ∞ -category C admits a zero object.
- (2.1.2) The ∞ -category C has all finite products and coproducts.
- (2.1.3) For any finite set I and any I -tuple $(X_i)_{i \in I}$ of objects of C , the map

$$\coprod X_i \rightarrow \prod X_i$$

in hC , given by the maps $\phi_{ij} : X_i \rightarrow X_j$, where ϕ_{ij} is zero unless $i = j$, in which case it is the identity, is an isomorphism.

If C admits finite direct sums, then for any finite set I and any I -tuple $(X_i)_{i \in I}$ of objects of C , we denote by $\bigoplus X_i$ the product (or, equivalently, the coproduct) of the X_i .

Suppose C an ∞ -category that admits direct sums. Then the homotopy category hC is easily seen to admit direct sums as well. Moreover, the mapping spaces in C admit the natural structure of a homotopy-commutative H -space: for any morphisms $f, g \in \text{Map}_C(X, Y)$, one may define $f + g \in \text{Map}_C(X, Y)$ as the composite

$$X \xrightarrow{\Delta} X \oplus X \xrightarrow{f \oplus g} Y \oplus Y \xrightarrow{\nabla} Y.$$

DEFINITION 2.2. An ∞ -category C that admits finite direct sums will be said to be *additive* if its homotopy category hC is additive; i.e., if for any two objects X and Y , the monoid $\pi_0 \text{Map}_C(X, Y)$ is a group.

Remark 2.3. An ∞ -category C with direct sums is additive just in case, for any objects X and Y , the *shear map*

$$\text{Map}_C(X, Y) \times \text{Map}_C(X, Y) \rightarrow \text{Map}_C(X, Y) \times \text{Map}_C(X, Y)$$

in the homotopy category of Kan simplicial sets that is given informally by the assignment $(f, g) \mapsto (f, f + g)$ is an isomorphism. Note in particular that additivity is a condition, not additional structure.

Example 2.4. Clearly the nerve of any ordinary additive category is an additive ∞ -category. Similarly, any stable ∞ -category is additive.

3. Exact ∞ -categories

Now we are ready to define exact ∞ -categories.

DEFINITION 3.1. Suppose \mathcal{C} an ∞ -category, and suppose \mathcal{C}_\dagger and \mathcal{C}^\dagger two subcategories of \mathcal{C} . Call a morphism of \mathcal{C}_\dagger *ingressive* or a *cofibration*, and call a morphism of \mathcal{C}^\dagger *egressive* or a *fibration*.

(3.1.1) A pullback square

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X' & \twoheadrightarrow & Y' \end{array}$$

is said to be *ambigressive* if $X' \twoheadrightarrow Y'$ is ingressive and $Y \twoheadrightarrow Y'$ is egressive. Dually, a pushout square

$$\begin{array}{ccc} X & \twoheadrightarrow & Y \\ \downarrow & & \downarrow \\ X' & \longrightarrow & Y' \end{array}$$

is said to be *ambigressive* if $X \twoheadrightarrow Y$ is ingressive and $X \twoheadrightarrow X'$ is egressive.

(3.1.2) The triple $(\mathcal{C}, \mathcal{C}_\dagger, \mathcal{C}^\dagger)$ is said to be an *exact ∞ -category* if it satisfies the following conditions.

- (3.1.2.1) The underlying ∞ -category \mathcal{C} is additive.
- (3.1.2.2) The pair $(\mathcal{C}, \mathcal{C}_\dagger)$ is a Waldhausen ∞ -category.
- (3.1.2.3) The pair $(\mathcal{C}, \mathcal{C}^\dagger)$ is a coWaldhausen ∞ -category.
- (3.1.2.4) A square in \mathcal{C} is an ambigressive pullback if and only if it is an ambigressive pushout.

(3.1.3) In an exact ∞ -category, an *exact sequence* is a cofiber/fiber sequence as below.

$$\begin{array}{ccc} X' & \twoheadrightarrow & X \\ \downarrow & & \downarrow \\ 0 & \twoheadrightarrow & X'' \end{array}$$

We will abuse notation by writing

$$X' \twoheadrightarrow X \twoheadrightarrow X''$$

for this square.

Remark 3.2. Note that in an exact ∞ -category, a morphism of an exact ∞ -category is egressive just in case it appears as the cofiber of an ingressive morphism, and, dually, a morphism of an exact ∞ -category is ingressive just in case it appears as the fiber of an egressive morphism. Indeed, any cofiber of an ingressive morphism is egressive, and any egressive morphism is equivalent to the cofiber of its fiber. This proves the first statement; the second is dual. Consequently, the class of cofibrations in an exact ∞ -category specifies the class of fibrations, and vice versa.

Example 3.3. (3.3.1) The nerve NC of an ordinary category C can be endowed with a triple structure yielding an exact ∞ -category if and only if C is an ordinary exact category, in

the sense of Quillen, wherein the admissible monomorphisms are exactly the cofibrations, and the admissible epimorphisms are exactly the fibrations. To prove this, one may observe that the ‘minimal’ axioms of Keller [Kel90, Appendix A] simply *are* the axioms listed above.

(3.3.2) At the other extreme, any stable ∞ -category is an exact ∞ -category in which all morphisms are both egressive and ingressive, and, conversely, any ∞ -category that can be regarded as an exact category with the *maximal* triple structure (in which any morphism is both ingressive and egressive) is a stable ∞ -category.

Example 3.4. We may interpolate between these two extremes. Suppose \mathcal{A} a stable ∞ -category equipped with a t-structure, and suppose $a, b \in \mathbf{Z}$.

(3.4.1) The ∞ -category $\mathcal{A}_{[a, +\infty)} := \mathcal{A}_{\geq a}$ admits an exact ∞ -category structure, in which every morphism is ingressive, but a morphism $Y \rightarrow Z$ is egressive just in case the induced morphism $\pi_a X \rightarrow \pi_a Y$ is an epimorphism of \mathcal{A}^\heartsuit .

(3.4.2) Dually, the ∞ -category $\mathcal{A}_{(-\infty, b]} := \mathcal{A}_{\leq b}$ admits an exact ∞ -category structure, in which every morphism is egressive, but a morphism $X \rightarrow Y$ is ingressive just in case the induced morphism $\pi_b X \rightarrow \pi_b Y$ is a monomorphism of \mathcal{A}^\heartsuit .

(3.4.3) We may intersect these subcategories to obtain the full subcategory

$$\mathcal{A}_{[a, b]} := \mathcal{A}_{\geq a} \cap \mathcal{A}_{\leq b} \subset \mathcal{A},$$

and we may intersect the subcategories of ingressive and egressive morphisms described to obtain the following exact ∞ -category structure on $\mathcal{A}_{[a, b]}$. A morphism $X \rightarrow Y$ is ingressive just in case the induced morphism $\pi_b X \rightarrow \pi_b Y$ is a monomorphism of the abelian category \mathcal{A}^\heartsuit . A morphism $Y \rightarrow Z$ is egressive just in case the induced morphism $\pi_a X \rightarrow \pi_a Y$ is an epimorphism of \mathcal{A}^\heartsuit .

Example 3.5. Yet more generally, suppose \mathcal{A} a stable ∞ -category, and suppose $\mathcal{C} \subset \mathcal{A}$ any full additive subcategory that is closed under extensions. Declare a morphism $X \rightarrow Y$ of \mathcal{C} to be ingressive just in case its cofiber in \mathcal{A} lies in \mathcal{C} . Dually, declare a morphism $Y \rightarrow Z$ of \mathcal{C} to be egressive just in case its fiber in \mathcal{A} lies in \mathcal{C} . Then \mathcal{C} is exact with this triple structure.

Remark 3.6. Thomason and Trobaugh called a triple $(C, C_\dagger, C^\dagger)$ of ordinary categories with direct sums whose nerves satisfy conditions (3.1.2.2–4) a *category with bifibrations* [TT90, Definition 1.2.2]. Include additivity, and this is precisely the notion of an ordinary exact category. (As an aside, we remark that the theory of exact ∞ -categories we delve into here really uses the additivity condition; without it, one would be unable to ensure that exact ∞ -categories form a full subcategory of Waldhausen ∞ -categories.)

When a class of weak equivalences is included, Thomason and Trobaugh used the term *biWaldhausen category* [TT90, Definition 1.2.4]. This notion still does not require additivity. However, suppose \mathbf{A} a *complcial biWaldhausen category* [TT90, Definition 1.2.11] that is closed under the canonical homotopy pullbacks and canonical homotopy pushouts of [TT90, Definition 1.1.2] in the sense of [TT90, Definition 1.3.5] such that the mapping cylinder and cocylinder functors of [TT90, Definition 1.3.4] satisfy [TT90, Definition 1.3.1.5] and its dual. Then the relative nerve [Bar13, Definition 1.5] of $(\mathbf{A}, w\mathbf{A})$ is clearly a stable ∞ -category. Consequently, every example of a category with cofibrations and weak equivalences that Thomason and Trobaugh study [TT90, §§ 2–11] is actually nothing more than a model for some stable ∞ -category.

4. Exact functors between exact ∞ -categories

DEFINITION 4.1. Suppose \mathcal{C} and \mathcal{D} two exact ∞ -categories. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ will be said to be *exact* if it preserves both cofibrations and fibrations and if F is exact both as a functor of Waldhausen ∞ -categories and as a functor of coWaldhausen ∞ -categories.

We denote by $\text{Fun}_{\mathbf{Exact}_\infty}(\mathcal{C}, \mathcal{D})$ the full subcategory of $\text{Fun}(\mathcal{C}, \mathcal{D})$ spanned by the exact functors $\mathcal{C} \rightarrow \mathcal{D}$.

This definition, when set against the definition of exact functor of Waldhausen categories (Recollection 1.2), appears to overburden the phrase ‘exact functor’ and to create the possibility for some ambiguity; however, in Proposition 4.8 we will see that in fact no ambiguity obtains.

For now, let us construct the ∞ -category of exact ∞ -categories.

Notation 4.2. Denote by $\mathbf{Exact}_\infty^\Delta$ the following simplicial category. The objects of $\mathbf{Exact}_\infty^\Delta$ are small exact ∞ -categories; for any two exact ∞ -categories \mathcal{C} and \mathcal{D} , let $\mathbf{Exact}_\infty^\Delta(\mathcal{C}, \mathcal{D})$ be the maximal Kan complex $\iota \text{Fun}_{\mathbf{Exact}_\infty}(\mathcal{C}, \mathcal{D})$ contained in $\text{Fun}_{\mathbf{Exact}_\infty}(\mathcal{C}, \mathcal{D})$. We write \mathbf{Exact}_∞ for the simplicial nerve [Lur09a, Definition 1.1.5.5] of $\mathbf{Exact}_\infty^\Delta$.

Remark 4.3. We could have equally well defined \mathbf{Exact}_∞ as the full subcategory of the pullback

$$\mathbf{Wald}_\infty \times_{\mathbf{Cat}_\infty} \mathbf{coWald}_\infty$$

spanned by the exact ∞ -categories.

We have already remarked that the formation of the opposite of a Waldhausen ∞ -category defines an equivalence

$$\mathbf{Wald}_\infty \xrightarrow{\sim} \mathbf{coWald}_\infty.$$

Since exact ∞ -categories are defined by fitting together the structure of a Waldhausen ∞ -category and a coWaldhausen ∞ -category in a self-dual manner, we obtain the following lemma.

LEMMA 4.4. *The formation of the opposite restricts to an autoequivalence*

$$\text{op} : \mathbf{Exact}_\infty \xrightarrow{\sim} \mathbf{Exact}_\infty.$$

This permits us to dualize virtually any assertion about exact ∞ -categories.

We now set about showing that the inclusion $\mathbf{Exact}_\infty \hookrightarrow \mathbf{Wald}_\infty$ is fully faithful. For this, we use in a nontrivial way the additivity condition for exact ∞ -categories. In particular, this additivity actually guarantees a greater compatibility between pullbacks and pushouts and between fibrations and cofibrations than one might at first expect.

LEMMA 4.5. *In an exact ∞ -category, a pushout square*

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ \downarrow & & \downarrow \\ X' & \longrightarrow & Y' \end{array}$$

in which the morphism $X \rightarrow Y$ is ingressive is also a pullback square. Dually, a pullback square

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X' & \longrightarrow & Y' \end{array}$$

in which the morphism $Y \rightarrow Y'$ is egressive is also a pushout square.

Proof. We prove the first statement; the second is dual. Since $X' \twoheadrightarrow Y'$ is ingressive, we may form the cofiber

$$\begin{array}{ccc} X' & \twoheadrightarrow & Y' \\ \downarrow & & \downarrow \\ 0 & \twoheadrightarrow & Z' \end{array}$$

which is an ambigressive square. Hence the square

$$\begin{array}{ccc} X & \twoheadrightarrow & Y \\ \downarrow & & \downarrow \\ 0 & \twoheadrightarrow & Z' \end{array}$$

is also ambigressive, whence we conclude that

$$\begin{array}{ccc} X & \twoheadrightarrow & Y \\ \downarrow & & \downarrow \\ X' & \twoheadrightarrow & Y' \end{array}$$

is a pullback square. □

The next pair of lemmas give a convenient way to replace pushout squares with exact sequences.

LEMMA 4.6. *For any exact sequence*

$$X' \xrightarrow{i} X \xrightarrow{p} \twoheadrightarrow X''$$

of an exact ∞ -category C , the object W formed as the pushout

$$\begin{array}{ccc} X' & \twoheadrightarrow & X \\ \downarrow & & \downarrow \\ X & \twoheadrightarrow & W \end{array}$$

is a direct sum $X \oplus X''$. Dually, the object V formed as the pullback

$$\begin{array}{ccc} V & \twoheadrightarrow & X \\ \downarrow & & \downarrow \\ X & \twoheadrightarrow & X'' \end{array}$$

is a direct sum $X' \oplus X$.

Proof. We prove the first assertion; the second is dual. Choose a fibrant simplicial category D whose nerve is equivalent to C . Now for any object T , the shear map

$$\mathrm{Map}_D(X, T) \times \mathrm{Map}_D(X, T) \xrightarrow{\sim} \mathrm{Map}_D(X, T) \times \mathrm{Map}_D(X, T)$$

induces an equivalence

$$\begin{aligned} & (\mathrm{Map}_D(X, T) \times \mathrm{Map}_D(X, T)) \times_{\mathrm{id} \times i^*, (\mathrm{Map}_D(X, T) \times \mathrm{Map}_D(X', T)), \mathrm{id} \times 0}^h (\mathrm{Map}_D(X, T) \times \Delta^0) \\ & \quad \downarrow \sim \\ & (\mathrm{Map}_D(X, T) \times \mathrm{Map}_D(X, T)) \times_{i^* \times i^*, (\mathrm{Map}_D(X', T) \times \mathrm{Map}_D(X', T)), \Delta}^h \mathrm{Map}_D(X', T) \end{aligned} \tag{4.6.1}$$

where:

- i^* denotes the map $\text{Map}_D(X, T) \rightarrow \text{Map}_D(Z, T)$ induced by $i : X' \rightarrow X$;
- 0 denotes a vertex $\Delta^0 \rightarrow \text{Map}_D(X', T)$ corresponding to a zero map; and
- Δ denotes the diagonal map.

The source of (4.6.1) is the product of $\text{Map}_D(X, T)$ with the space of squares of the form

$$\begin{array}{ccc} X' & \xrightarrow{i} & X \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & T \end{array}$$

in C , and the target is equivalent to the space of squares of the form

$$\begin{array}{ccc} X' & \xrightarrow{i} & X \\ \downarrow i & & \downarrow \\ X & \longrightarrow & T \end{array}$$

in C . Consequently, the map (4.6.1) specifies an equivalence

$$\text{Map}_D(X, T) \times \text{Map}_D(X'', T) \xrightarrow{\sim} \text{Map}(W, T).$$

This equivalence is clearly functorial in T , so it specifies an equivalence $W \xrightarrow{\sim} X \oplus X''$. □

LEMMA 4.7. *In an exact ∞ -category, suppose that*

$$\begin{array}{ccc} X & \xrightarrow{i} & Y \\ p \downarrow & & \downarrow q \\ X' & \xrightarrow{i'} & Y' \end{array}$$

is either a pushout square in which $X \rightarrow Y$ is ingressive or a pullback square in which $Y \rightarrow Y'$ is egressive. Then the morphism

$$\begin{pmatrix} -p \\ i \end{pmatrix} : X \rightarrow X' \oplus Y$$

is ingressive, the morphism

$$(i' \quad q) : X' \oplus Y \rightarrow Y'$$

is egressive, and these maps fit into an exact sequence

$$X \twoheadrightarrow X' \oplus Y \twoheadrightarrow Y'.$$

Proof. We prove the assertion for pushout squares; the other assertion is dual. We form a diagram

$$\begin{array}{ccccc} X & \longrightarrow & X' & \twoheadrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ V & \longrightarrow & V' & \twoheadrightarrow & Y' \\ \downarrow & & \downarrow & & \downarrow \\ Y & \longrightarrow & Y' & \twoheadrightarrow & Z \end{array}$$

in which every square is a pullback square. By the previous lemma, V' is a direct sum $X' \oplus Y'$, and V is a direct sum $X \oplus Y'$. The desired exact sequence is the top rectangle. □

PROPOSITION 4.8. *The following are equivalent for a functor $\psi : \mathcal{C} \rightarrow \mathcal{D}$ between two ∞ -categories with exact ∞ -category structures.*

(4.8.1) *The functor ψ carries cofibrations to cofibrations, it carries fibrations to fibrations, and as a functor of exact ∞ -categories, ψ is exact.*

(4.8.2) *The functor ψ carries cofibrations to cofibrations, and as a functor of Waldhausen ∞ -categories, ψ is exact.*

(4.8.3) *The functor ψ carries fibrations to fibrations, and as a functor of coWaldhausen ∞ -categories, ψ is exact.*

Proof. It is clear that the first condition implies the other two. We shall show that the second implies the first; the proof that the third condition implies the first is dual. So suppose ψ preserves cofibrations and is exact as a functor of Waldhausen ∞ -categories. Because a morphism is egressive just in case it can be exhibited as a cofiber, ψ preserves fibrations as well. A pullback square

$$\begin{array}{ccc} X & \xrightarrow{i} & Y \\ p \downarrow & & \downarrow q \\ X' & \xrightarrow{i'} & Y' \end{array}$$

in which $p : X \twoheadrightarrow X'$ and $q : Y \twoheadrightarrow Y'$ are egressive can be factored as

$$\begin{array}{ccccc} X & \xrightarrow{\begin{pmatrix} -p \\ i \end{pmatrix}} & X' \oplus Y & \xrightarrow{\text{pr}_2} & Y \\ p \downarrow & & \downarrow \text{id} \oplus q & & \downarrow q \\ X' & \xrightarrow{\begin{pmatrix} -\text{id} \\ i' \end{pmatrix}} & X' \oplus Y' & \xrightarrow{\text{pr}_2} & Y' \end{array}$$

in which all three rectangles are pullbacks. By the previous lemma, the left-hand square is an ambigressive pullback/pushout, so when we apply ψ , we obtain a diagram

$$\begin{array}{ccccc} \psi X & \xrightarrow{\begin{pmatrix} -\psi p \\ \psi i \end{pmatrix}} & \psi X' \oplus \psi Y & \xrightarrow{\text{pr}_2} & \psi Y \\ \psi p \downarrow & & \downarrow \text{id} \oplus \psi q & & \downarrow \psi q \\ \psi X' & \xrightarrow{\begin{pmatrix} -\text{id} \\ \psi i' \end{pmatrix}} & \psi X' \oplus \psi Y' & \xrightarrow{\text{pr}_2} & \psi Y' \end{array}$$

in which the right-hand square is easily seen to be a pullback, and the left-hand square, being an ambigressive pushout, is also an ambigressive pullback. \square

COROLLARY 4.8.1. *The forgetful functors*

$$\mathbf{Exact}_\infty \rightarrow \mathbf{Wald}_\infty \quad \text{and} \quad \mathbf{Exact}_\infty \rightarrow \mathbf{coWald}_\infty$$

are fully faithful.

In particular, we may say that a Waldhausen ∞ -category \mathcal{C} ‘is’ an exact ∞ -category if it lies in the essential image of the forgetful functor $\mathbf{Exact}_\infty \rightarrow \mathbf{Wald}_\infty$, and we will treat this forgetful functor as if it were an inclusion. Since this functor is fully faithful, this is not a significant abuse of terminology. We make sense of the assertion that a coWaldhausen ∞ -category ‘is’ an exact ∞ -category in a dual manner.

4.9. The essential image of the forgetful functor $\mathbf{Exact}_\infty \rightarrow \mathbf{Wald}_\infty$ is spanned by those Waldhausen ∞ -categories that satisfy the following three criteria.

(4.9.1) The underlying ∞ -category is additive.

(4.9.2) The class of morphisms that can be exhibited as the cofiber of some cofibration is closed under pullback.

(4.9.3) Every cofibration is the fiber of its cofiber.

The essential image of the forgetful functor $\mathbf{Exact}_\infty \rightarrow \mathbf{coWald}_\infty$ is described in a dual manner.

5. Theories and duality

Algebraic K -theory is a particular example of what we called an additive theory in [Bar13]. In effect, additive theories are the natural homology theories for Waldhausen ∞ -categories. To tell this story, it is necessary to recall some pleasant facts about the ∞ -categories \mathbf{Wald}_∞ and \mathbf{coWald}_∞ .

RECOLLECTION 5.1. In particular, the ∞ -category \mathbf{Wald}_∞ (and hence also the ∞ -category \mathbf{coWald}_∞) enjoys a number of excellent formal properties. We showed in [Bar13, Proposition 4.6] that it admits direct sums, and we also showed in [Bar13, Proposition 4.7] that it is compactly generated [Lur09a, Definition 5.5.7.1] in the sense that every Waldhausen ∞ -category is in fact the filtered union of its finitely presented Waldhausen subcategories (that is, of Waldhausen subcategories that are compact as objects of \mathbf{Wald}_∞).

Furthermore, suppose $\mathcal{X} : \Lambda \rightarrow \mathbf{Wald}_\infty$ a diagram of Waldhausen ∞ -categories. The limit $\lim \mathcal{X}$ is computed by forming the limit in \mathbf{Cat}_∞ [Lur09a, Corollary 3.3.3.2] and then declaring a morphism to be ingressive if its image in each \mathcal{X}_α is so [Bar13, Proposition 4.3]. Similarly, if Λ is filtered, then the colimit $\operatorname{colim} \mathcal{X}$ is computed by forming the colimit in \mathbf{Cat}_∞ [Lur09a, Corollary 3.3.4.3] and then declaring $(\operatorname{colim} \mathcal{X})_\dagger$ to be the union of the images of the ∞ -categories $\mathcal{X}_{\alpha,\dagger}$ [Bar13, Proposition 4.4].

Since the ∞ -categories \mathbf{Wald}_∞ and \mathbf{coWald}_∞ are equivalent, it is clear that all these properties are enjoyed by the latter as well as the former. We will denote by $\mathbf{Wald}_\infty^\omega$ (respectively, by $\mathbf{coWald}_\infty^\omega$) the full subcategory of \mathbf{Wald}_∞ (respectively, \mathbf{coWald}_∞) spanned by the finitely presented Waldhausen ∞ -categories (respectively, the finitely presented \mathbf{coWald} ∞ -categories).

These formal properties can be regarded as analogues of a few of the formal properties enjoyed by the ordinary category $\mathbf{V}(k)$ of vector spaces: vector spaces are of course additive, and any vector space is the union of its finite-dimensional subspaces. Furthermore, the underlying set of a limit or filtered colimit of vector spaces is the limit or filtered colimit of the underlying sets.

We will only be interested in functors on \mathbf{Wald}_∞ or \mathbf{coWald}_∞ that are (1) trivial on the zero Waldhausen ∞ -category, and (2) completely determined by their values on finitely presented Waldhausen ∞ -categories.

RECOLLECTION 5.2 [Bar13, Definition 7.1]. A *theory* $\phi : \mathbf{Wald}_\infty \rightarrow \mathbf{Kan}$ is a functor that preserves terminal objects and filtered colimits.

Similarly, a *theory* $\phi : \mathbf{coWald}_\infty \rightarrow \mathbf{Kan}$ is a functor that preserves terminal objects and filtered colimits.

Notation 5.3. Denote by \mathbf{Thy} (respectively, \mathbf{Thy}^\vee) the full subcategory of the ∞ -category $\mathbf{Fun}(\mathbf{coWald}_\infty, \mathbf{Kan})$ spanned by the theories.

Example 5.4. The most important example of a theory (in either sense) is the *moduli space of objects* functor $\mathcal{C} \mapsto \iota\mathcal{C}$. Here $\iota\mathcal{C}$ denotes the largest Kan complex contained in \mathcal{C} [Lur09a, Proposition 1.2.5.3].

Of course there is a canonical equivalence between theories on Waldhausen ∞ -categories and theories on coWaldhausen ∞ -categories.

DEFINITION 5.5. For any theory $\phi : \mathbf{Wald}_\infty \rightarrow \mathbf{Kan}$, the *dual theory* ϕ^\vee is the composite $\phi \circ \text{op} : \mathbf{coWald}_\infty \rightarrow \mathbf{Kan}$. This construction clearly yields an equivalence of ∞ -categories $\text{Thy} \xrightarrow{\sim} \text{Thy}^\vee$.

Now we are prepared to define the key notion of an *exact duality* on a theory ϕ .

DEFINITION 5.6. An *exact duality* on a theory ϕ is an equivalence

$$\eta : \phi|_{\mathbf{Exact}_\infty} \xrightarrow{\sim} \phi^\vee|_{\mathbf{Exact}_\infty}$$

of the ∞ -category $\text{Fun}(\mathbf{Exact}_\infty, \mathcal{E})$.

Example 5.7. Suppose $\rho : \mathbf{Cat}_\infty^* \rightarrow \mathbf{Kan}$ a functor from the ∞ -category \mathbf{Cat}_∞^* of pointed ∞ -categories to \mathbf{Kan} that preserves the terminal object and filtered colimits. Then an equivalence $\rho \xrightarrow{\sim} \rho \circ \text{op}$ induces an exact duality on the composite

$$\mathbf{Wald}_\infty \rightarrow \mathbf{Cat}_\infty^* \rightarrow \mathbf{Kan}.$$

For instance, the functor $\iota : \mathbf{Cat}_\infty^* \rightarrow \mathbf{Kan}$ admits an equivalence $\iota \xrightarrow{\sim} \iota \circ \text{op}$; consequently, the theory $\iota : \mathbf{Wald}_\infty \rightarrow \mathbf{Kan}$ admits an exact duality.

A general theory does not reflect much about the (co)Waldhausen structure. *Additive* theories are much more sensitive. To talk about them, we have to recall our construction of \mathcal{F} and \mathcal{S} .

RECOLLECTION 5.8. In [Bar13, §5] we defined, for any Waldhausen ∞ -category \mathcal{C} , an ∞ -category $\mathcal{F}(\mathcal{C})$ and a full subcategory $\mathcal{S}(\mathcal{C}) \subset \mathcal{F}(\mathcal{C})$. An object of $\mathcal{F}(\mathcal{C})$ is a pair (m, X) consisting of a natural number m and a sequence

$$X_0 \rightrightarrows \cdots \rightrightarrows X_m$$

of cofibrations of \mathcal{C} ; a morphism $(m, X) \rightarrow (n, Y)$ of $\mathcal{F}(\mathcal{C})$ consists of a morphism $\eta : \mathbf{n} \rightarrow \mathbf{m}$ of Δ and a commutative diagram

$$\begin{array}{ccc} X_{\eta(0)} & \rightrightarrows & \cdots \rightrightarrows & X_{\eta(n)} \\ \downarrow & & & \downarrow \\ Y_0 & \rightrightarrows & \cdots \rightrightarrows & Y_n \end{array}$$

of \mathcal{C} . The full subcategory $\mathcal{S}(\mathcal{C}) \subset \mathcal{F}(\mathcal{C})$ is the one spanned by those pairs (m, X) such that X_0 is a zero object of \mathcal{C} .

The assignment $(m, X) \mapsto \mathbf{m}$ defines functors

$$p_{\mathcal{C}} : \mathcal{F}(\mathcal{C}) \rightarrow N\Delta^{\text{op}} \quad \text{and} \quad q_{\mathcal{C}} : \mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}.$$

Let us write $\mathcal{F}_m(\mathcal{C}) := p_{\mathcal{C}}^{-1}(\mathbf{m})$ and $\mathcal{S}_m(\mathcal{C}) := q_{\mathcal{C}}^{-1}(\mathbf{m})$.

The first basic fact to understand about these functors is that they are *cocartesian fibrations* [Lur09a, Definition 2.4.2.1]; that is, there are functors

$$\mathbf{F}_*(\mathcal{C}) : N\Delta^{\text{op}} \longrightarrow \mathbf{Cat}_\infty \quad \text{and} \quad \mathbf{S}_*(\mathcal{C}) : N\Delta^{\text{op}} \longrightarrow \mathbf{Cat}_\infty$$

(the functors that classify $p_{\mathcal{C}}$ and $q_{\mathcal{C}}$, [Lur09a, Definition 3.3.2.2]) and equivalences $\mathbf{F}_m(\mathcal{C}) \simeq \mathcal{F}_m(\mathcal{C})$ and $\mathbf{S}_m(\mathcal{C}) \simeq \mathcal{S}_m(\mathcal{C})$ such that for any morphism $\eta : \mathbf{n} \rightarrow \mathbf{m}$ of Δ , the space of morphisms $(m, X) \rightarrow (n, Y)$ of $\mathcal{F}(\mathcal{C})$ (respectively, of $\mathcal{S}(\mathcal{C})$) that cover η is equivalent to the space of morphisms $\eta_! X \rightarrow Y$, where $\eta_!$ is shorthand notation for the image of η under $\mathbf{F}_*(\mathcal{C})$ (respectively, $\mathbf{S}_*(\mathcal{C})$). In particular, for any object X of $p_{\mathcal{C}}^{-1}(\mathbf{m})$ (respectively, $q_{\mathcal{C}}^{-1}(\mathbf{m})$), there exists a special edge, called a *cocartesian edge*, from X to $\eta_! X$.

The functor $\mathbf{F}_*(\mathcal{C})$ is easy to describe: it carries \mathbf{m} to the full subcategory of $\text{Fun}(\Delta^m, \mathcal{C})$ spanned by those functors $X : \Delta^m \rightarrow \mathcal{C}$ such that each morphism $X_i \rightarrow X_{i+1}$ is ingressive; the functoriality in \mathbf{m} is obvious here. The functor $\mathbf{S}_*(\mathcal{C})$ is a tad trickier to describe: morally, it carries \mathbf{m} to the full subcategory of $\mathbf{F}_m(\mathcal{C})$ spanned by those objects X such that X_0 is a zero object, and a map $\eta : \mathbf{n} \rightarrow \mathbf{m}$ of Δ induces a functor $\mathbf{S}_m(\mathcal{C}) \rightarrow \mathbf{S}_n(\mathcal{C})$ that carries an object X to the object

$$0 \rightrightarrows X_{\eta(1)}/X_{\eta(0)} \rightrightarrows \cdots \rightrightarrows X_{\eta(n)}/X_{\eta(0)}.$$

One can opt to make compatible choices of these quotients to rectify this into an actual functor of ∞ -categories, or, alternately, one can forget about $\mathbf{S}_*(\mathcal{C})$ and stick with the cocartesian fibration $\mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$. In [Wal85], Waldhausen opted for the former; in [Bar13], we opted for the latter. Which approach one chooses to take is largely a matter of taste or convenience; conceptually, there is no difference.

In [Bar13, Corollary 5.20.1], we show that the left adjoints $\mathcal{F}_m(\mathcal{C}) \rightarrow \mathcal{S}_m(\mathcal{C})$ to the inclusions $\mathcal{S}_m(\mathcal{C}) \hookrightarrow \mathcal{F}_m(\mathcal{C})$ fit together over $N\Delta^{\text{op}}$ to make a natural transformation $\mathbf{F}_*(\mathcal{C}) \rightarrow \mathbf{S}_*(\mathcal{C})$.

The next significant thing to understand about $\mathcal{F}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ and $\mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ is that they are what we have called *Waldhausen cocartesian fibrations* [Bar13, Definition 3.20]. That is, there are subcategories

$$\mathcal{F}(\mathcal{C})_{\dagger} \subset \mathcal{F}(\mathcal{C}) \quad \text{and} \quad \mathcal{S}(\mathcal{C})_{\dagger} \subset \mathcal{S}(\mathcal{C})$$

such that for any $m \geq 0$, the pairs

$$(\mathcal{F}_m(\mathcal{C}), \mathcal{F}_m(\mathcal{C}) \cap \mathcal{F}(\mathcal{C})_{\dagger}) \quad \text{and} \quad (\mathcal{S}_m(\mathcal{C}), \mathcal{S}_m(\mathcal{C}) \cap \mathcal{S}(\mathcal{C})_{\dagger})$$

are Waldhausen ∞ -categories, and for any morphism $\eta : \mathbf{n} \rightarrow \mathbf{m}$ of Δ , the functors

$$\eta : \mathcal{F}_m(\mathcal{C}) \simeq \mathbf{F}_m(\mathcal{C}) \rightarrow \mathbf{F}_n(\mathcal{C}) \simeq \mathcal{F}_n(\mathcal{C}) \quad \text{and} \quad \eta : \mathcal{S}_m(\mathcal{C}) \simeq \mathbf{S}_m(\mathcal{C}) \rightarrow \mathbf{S}_n(\mathcal{C}) \simeq \mathcal{S}_n(\mathcal{C})$$

are exact functors of Waldhausen ∞ -categories. In effect, a morphism $(m, X) \rightarrow (n, Y)$ of either $\mathcal{F}(\mathcal{C})$ or $\mathcal{S}(\mathcal{C})$ will be declared ingressive just in case the morphism $\mathbf{n} \rightarrow \mathbf{m}$ is an isomorphism and the diagram

$$\begin{array}{ccc} X_0 & \longrightarrow & \cdots \longrightarrow & X_m \\ \downarrow & & & \downarrow \\ Y_0 & \longrightarrow & \cdots \longrightarrow & Y_m \end{array}$$

has the property that for any integer $0 \leq i \leq j \leq m$, the natural morphism $Y_i \cup^{X_i} X_j \rightarrow Y_j$ (whose source is the pushout in the ∞ -categorical sense, of course) is ingressive. We show

that $\mathcal{F}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ and $\mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ are Waldhausen cocartesian fibrations in [Bar13, Proposition 5.11 and Theorem 5.20]. Consequently, the functors that classify these fibrations can be lifted to simplicial Waldhausen ∞ -categories

$$\mathbf{F}_*(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Wald}_\infty \quad \text{and} \quad \mathbf{S}_*(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Wald}_\infty.$$

Dually, for any coWaldhausen ∞ -category \mathcal{D} , we obtain *coWaldhausen cartesian fibrations*

$$\mathcal{F}^\vee(\mathcal{D}) := \mathcal{F}(\mathcal{D}^{\text{op}})^{\text{op}} \rightarrow N\Delta \quad \text{and} \quad \mathcal{S}^\vee(\mathcal{D}) := \mathcal{S}(\mathcal{D}^{\text{op}})^{\text{op}} \rightarrow N\Delta.$$

The functors that classify these fibrations are simplicial coWaldhausen ∞ -categories

$$\mathbf{F}_*^\vee(\mathcal{D}) : N\Delta^{\text{op}} \rightarrow \mathbf{coWald}_\infty \quad \text{and} \quad \mathbf{S}_*^\vee(\mathcal{D}) : N\Delta^{\text{op}} \rightarrow \mathbf{coWald}_\infty.$$

DEFINITION 5.9 [Bar13, Definition 7.5]. A theory $\phi : \mathbf{Wald}_\infty \rightarrow \mathbf{Kan}$ will be said to be *additive* if for any Waldhausen ∞ -category \mathcal{C} , the simplicial space

$$\phi \circ \mathbf{S}_*(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Kan}$$

is a group object in the sense of [Lur09a, Definition 7.2.2.1]. That is, for any $m \geq 0$, the map

$$\phi \circ \mathbf{S}_m(\mathcal{C}) \rightarrow \prod_{i=1}^m \phi \circ \mathbf{S}_{\{i-1, i\}}(\mathcal{C})$$

is an equivalence, and the monoid $\pi_0(\phi \circ \mathbf{S}_1(\mathcal{C}))$ is a group. Write **Add** for the full subcategory of **Thy** spanned by the additive theories.

Dually, a theory $\phi : \mathbf{coWald}_\infty \rightarrow \mathbf{Kan}$ will be said to be *additive* just in case it is the dual theory of an additive theory $\mathbf{Wald}_\infty \rightarrow \mathbf{Kan}$. That is, ϕ is additive just in case, for any coWaldhausen ∞ -category \mathcal{C} , the simplicial space

$$\phi \circ \mathbf{S}_*^\vee(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Kan}$$

is a group object. Write **Add**[∨] for the full subcategory of **Thy**[∨] spanned by the additive theories.

RECOLLECTION 5.10. The main theorems of [Bar13] are (1) that there exist what we call the *fissile derived ∞ -category* $\mathbf{D}_{\text{add}}(\mathbf{Wald}_\infty)$ and a functor

$$\mathbf{Wald}_\infty \rightarrow \mathbf{D}_{\text{add}}(\mathbf{Wald}_\infty)$$

an equivalence

$$\mathbf{Exc}(\mathbf{D}_{\text{add}}(\mathbf{Wald}_\infty), \mathbf{Kan}) \simeq \mathbf{Add},$$

where **Exc** denotes the ∞ -category of reduced excisive functors that preserve sifted colimits [Bar13, Theorems 7.4 and 7.6], and (2) that the suspension in $\mathbf{D}_{\text{add}}(\mathbf{Wald}_\infty)$ of a Waldhausen ∞ -category \mathcal{C} is given by a formal colimit $\text{colim } \mathbf{S}_*(\mathcal{C})$ [Bar13, Corollary 6.9.1]. As a consequence, we deduced one may use the Goodwillie differential to find the best additive approximation to a theory ϕ :

$$D(\phi)(\mathcal{C}) \simeq \text{colim}_{m \rightarrow \infty} \Omega^m | \phi \circ \mathbf{S}_* \cdots \mathbf{S}_*(\mathcal{C}) |_{(N\Delta^{\text{op}})^m},$$

where $|\cdot|_{(N\Delta^{\text{op}})^m}$ denotes the colimit of an m -simplicial space [Bar13, Theorem 7.8]. The functor D then defines a left adjoint to the inclusion $\mathbf{Thy} \hookrightarrow \mathbf{Add}$.

Dually, the inclusion $\mathbf{Add}^\vee \hookrightarrow \mathbf{Thy}^\vee$ admits a left adjoint D^\vee , given by the formula

$$D^\vee(\phi)(\mathcal{C}) \simeq \text{colim}_{m \rightarrow \infty} \Omega^m | \phi \circ \mathbf{S}_*^\vee \cdots \mathbf{S}_*^\vee(\mathcal{C}) |_{(N\Delta^{\text{op}})^m}.$$

The purpose of the remainder of this section is to describe a circumstance in which an exact duality on a theory ϕ descends to an exact duality on $D\phi$, and to show that these conditions obtain when $\phi = \iota$, giving a functorial equivalence $K(\mathcal{C}) \simeq K(\mathcal{C}^{\text{op}})$ for exact ∞ -categories \mathcal{C} .

So suppose ϕ a theory with an exact duality. Note that $(D(\phi))^{\vee}$ is by construction equivalent to $D^{\vee}(\phi^{\vee})$, so such a result can be thought of as giving an equivalence $D(\phi) \simeq D^{\vee}(\phi^{\vee})$ on exact ∞ -categories.

Consequently, for an exact ∞ -category \mathcal{C} , we aim to produce a kind of duality between the Waldhausen cocartesian fibration $\mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ and the coWaldhausen cartesian fibration $\mathcal{S}^{\vee}(\mathcal{C}) \rightarrow N\Delta$.

More precisely, we will construct a functor $\tilde{\mathbf{S}}_*(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Cat}_{\infty}$ such that on the one hand, $\tilde{\mathbf{S}}_*$ classifies the cocartesian fibration $\mathcal{S}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$, and on the other, the composite of $\tilde{\mathbf{S}}_*$ with the functor $\text{op} : N\Delta^{\text{op}} \rightarrow N\Delta^{\text{op}}$ given by $[\mathbf{n}] \mapsto [\mathbf{n}]^{\text{op}}$ is a straightening of the cartesian fibration $\mathcal{S}^{\vee}(\mathcal{C}) \rightarrow N\Delta$.

In order to do this, we introduce thickened versions $\tilde{\mathcal{F}}, \tilde{\mathcal{F}}^{\vee}$ of the constructions $\mathcal{S}, \mathcal{S}^{\vee}$.

Notation 5.11. Let $\tilde{\mathbf{M}}$ be the following ordinary category. The objects are triples (m, i, j) consisting of integers $0 \leq i \leq j \leq m$, and a morphism $(n, k, \ell) \rightarrow (m, i, j)$ is a morphism $\phi : [\mathbf{m}] \rightarrow [\mathbf{n}]$ of Δ such that $k \leq \phi(i)$ and $\ell \leq \phi(j)$.

We declare an edge $(n, k, \ell) \rightarrow (m, i, j)$ of $N\tilde{\mathbf{M}}$ to be *ingressive* if the underlying edge $\mathbf{m} \rightarrow \mathbf{n}$ of Δ is an isomorphism and if $\ell = j$. Dually, we declare an edge $(n, k, \ell) \rightarrow (m, i, j)$ of $N\tilde{\mathbf{M}}$ to be *egressive* if the underlying edge $\mathbf{m} \rightarrow \mathbf{n}$ of Δ is an isomorphism and if $i = k$. We write $N\tilde{\mathbf{M}}_{\dagger}$ for the subcategory of $N\tilde{\mathbf{M}}$ consisting of the ingressive morphisms, and we write $N\tilde{\mathbf{M}}^{\dagger}$ for the subcategory of $N\tilde{\mathbf{M}}$ consisting of the egressive morphisms.

The fiber of the functor $N\tilde{\mathbf{M}} \rightarrow N\Delta^{\text{op}}$ over a vertex $\mathbf{n} \in N\Delta^{\text{op}}$ is the arrow ∞ -category $\mathcal{O}(\Delta^n) := \text{Fun}(\Delta^1, \Delta^n)$.

Now the functor $N\tilde{\mathbf{M}} \rightarrow N\Delta^{\text{op}}$ is a cartesian fibration, and so its opposite $N\tilde{\mathbf{M}}^{\text{op}} \rightarrow N\Delta$ is a cocartesian fibration. Furthermore, one verifies easily that for any morphism $\eta : \mathbf{n} \rightarrow \mathbf{m}$, the induced functor

$$\eta! : \mathcal{O}(\Delta^n) \rightarrow \mathcal{O}(\Delta^m)$$

is the obvious one, and it preserves both ingressive and egressive morphisms.

CONSTRUCTION 5.12. If \mathcal{C} is a Waldhausen ∞ -category, write $\tilde{\mathcal{F}}(\mathcal{C})$ for the simplicial set over $N\Delta^{\text{op}}$ satisfying the following universal property, which follows the general pattern set in [Lur09a, Corollary 3.2.2.13]. We require, for any simplicial set K and any map $\sigma : K \rightarrow N\Delta^{\text{op}}$, a bijection

$$\text{Mor}_{/(N\Delta^{\text{op}})}(K, \tilde{\mathcal{F}}(\mathcal{C})) \cong \text{Mor}_{s\mathbf{Set}(2)}((K \times_{N\Delta^{\text{op}}} N\tilde{\mathbf{M}}, K \times_{N\Delta^{\text{op}}} (N\tilde{\mathbf{M}})_{\dagger}), (\mathcal{C}, \mathcal{C}_{\dagger})),$$

functorial in σ . Here, the category $s\mathbf{Set}(2)$ is the ordinary category of pairs (X, A) of simplicial sets X equipped with a simplicial subset $A \subset X$. (Note that since the functor of K on the right-hand side carries colimits to limits, the simplicial set $\tilde{\mathcal{F}}(\mathcal{C})$ does indeed exist.) By [Bar13, Proposition 3.18], the map $\tilde{\mathcal{F}}(\mathcal{C}) \rightarrow N\Delta^{\text{op}}$ is a cocartesian fibration.

Dually, if \mathcal{C} is a coWaldhausen ∞ -category, write $\tilde{\mathcal{F}}^{\vee}(\mathcal{C})$ for the simplicial set over $N\Delta$ satisfying the following universal property. We require, for any simplicial set K and any map $\sigma : K \rightarrow N\Delta$, a bijection

$$\text{Mor}_{/N\Delta}(K, \tilde{\mathcal{F}}^{\vee}(\mathcal{C})) \cong \text{Mor}_{s\mathbf{Set}(2)}((K \times_{N\Delta} N\tilde{\mathbf{M}}^{\text{op}}, K \times_{N\Delta} (N\tilde{\mathbf{M}}^{\dagger})^{\text{op}}), (\mathcal{C}, \mathcal{C}^{\dagger})),$$

functorial in σ . By the dual of [Bar13, Proposition 3.18], the map $\tilde{\mathcal{F}}^{\vee}(\mathcal{C}) \rightarrow N\Delta$ is a cartesian fibration, and it is clear that

$$\tilde{\mathcal{F}}^{\vee}(\mathcal{C}) \cong \tilde{\mathcal{F}}(\mathcal{C}^{\text{op}})^{\text{op}}.$$

The objects of the ∞ -category $\widetilde{\mathcal{F}}(\mathcal{C})$ may be described as pairs (m, X) consisting of a nonnegative integer m and a functor $X : \mathcal{O}(\Delta^m) \rightarrow \mathcal{C}$ that carries ingressive morphisms to ingressive morphisms. Dually, the objects of the ∞ -category $\widetilde{\mathcal{F}}^\vee(\mathcal{C})$ may be described as pairs (m, X) consisting of a nonnegative integer m and a functor $X : \mathcal{O}(\Delta^m)^{\text{op}} \rightarrow \mathcal{C}$ that carries egressive morphisms to egressive morphisms.

Now if \mathcal{C} is a Waldhausen ∞ -category, we let $\widetilde{\mathcal{F}}(\mathcal{C}) \subset \widetilde{\mathcal{F}}(\mathcal{C})$ denote the full subcategory spanned by those pairs (m, X) such that for any integer $0 \leq i \leq m$, the object $X(i, i)$ is a zero object of \mathcal{C} , and for any integers $0 \leq i \leq k \leq j \leq \ell \leq m$, the square

$$\begin{array}{ccc} X(i, j) & \twoheadrightarrow & X(i, \ell) \\ \downarrow & & \downarrow \\ X(k, j) & \twoheadrightarrow & X(k, \ell) \end{array}$$

is a pushout. Dually, if \mathcal{C} is a coWaldhausen ∞ -category, let $\widetilde{\mathcal{F}}^\vee(\mathcal{C}) \subset \widetilde{\mathcal{F}}^\vee(\mathcal{C})$ denote the full subcategory spanned by those pairs (m, X) such that for any integer $0 \leq i \leq m$, the object $X(i, i)$ is a zero object of \mathcal{C} , and for any integers $0 \leq i \leq k \leq j \leq \ell \leq m$, the square

$$\begin{array}{ccc} X(k, \ell) & \longrightarrow & X(k, j) \\ \downarrow & & \downarrow \\ X(i, \ell) & \longrightarrow & X(i, j) \end{array}$$

is a pullback. Since ambigressive pullbacks and ambigressive pushouts coincide, we deduce that

$$\widetilde{\mathcal{F}}^\vee(\mathcal{C}) \cong \widetilde{\mathcal{F}}(\mathcal{C}^{\text{op}})^{\text{op}}.$$

Notation 5.13. As in [Bar13], the constructions above yield functors

$$\widetilde{\mathcal{F}} : \mathbf{Wald}_\infty \rightarrow \mathbf{Cat}_{\infty, /N\Delta^{\text{op}}} \quad \text{and} \quad \widetilde{\mathcal{F}}^\vee : \mathbf{coWald}_\infty \rightarrow \mathbf{Cat}_{\infty, /N\Delta}.$$

The functor $M \rightarrow \widetilde{M}$ given by the assignment $(m, i) \mapsto (m, 0, i)$ induces a natural transformation $\widetilde{\mathcal{F}} \rightarrow \mathcal{F}$ over $N\Delta^{\text{op}}$ and a natural transformation $\widetilde{\mathcal{F}}^\vee \rightarrow \mathcal{F}^\vee$ over $N\Delta$.

Furthermore, we can pass to the functors that classify these fibrations to obtain functors

$$\widetilde{\mathbf{S}}_* : \mathbf{Wald}_\infty \rightarrow \mathbf{Fun}(N\Delta^{\text{op}}, \mathbf{Cat}_\infty) \quad \text{and} \quad \widetilde{\mathbf{S}}_*^\vee : \mathbf{coWald}_\infty \rightarrow \mathbf{Fun}(N\Delta^{\text{op}}, \mathbf{Cat}_\infty).$$

For any Waldhausen ∞ -category (respectively, any coWaldhausen ∞ -category) \mathcal{C} , the simplicial category $\widetilde{\mathbf{S}}_*(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Cat}_\infty$ (respectively, $\widetilde{\mathbf{S}}_*^\vee(\mathcal{C}) : N\Delta^{\text{op}} \rightarrow \mathbf{Cat}_\infty$) carries an object \mathbf{m} to the full subcategory

$$\widetilde{\mathbf{S}}_{\mathbf{m}}(\mathcal{C}) \subset \mathbf{Fun}(\mathcal{O}(\Delta^m), \mathcal{C}) \quad (\text{respectively, } \widetilde{\mathbf{S}}_{\mathbf{m}}^\vee(\mathcal{C}) \subset \mathbf{Fun}(\mathcal{O}(\Delta^m)^{\text{op}}, \mathcal{C}))$$

spanned by those diagrams X such that for any integer $0 \leq i \leq m$, the object $X(i, i)$ is a zero object of \mathcal{C} , and for any integers $0 \leq i \leq k \leq j \leq \ell \leq m$, the square

$$\begin{array}{ccc} X(i, j) & \twoheadrightarrow & X(i, \ell) \\ \downarrow & & \downarrow \\ X(k, j) & \twoheadrightarrow & X(k, \ell) \end{array}$$

is an ambigressive pushout (respectively, for any integers $0 \leq i \leq k \leq j \leq \ell \leq m$, the square

$$\begin{array}{ccc} X(k, \ell) & \longrightarrow & X(k, j) \\ \downarrow & & \downarrow \\ X(i, \ell) & \longrightarrow & X(i, j) \end{array}$$

is an ambigressive pullback).

In light of the uniqueness of limits and colimits in ∞ -categories [Lur09a, Proposition 1.2.12.9], one readily has the following proposition.

PROPOSITION 5.14. *Suppose \mathcal{C} a Waldhausen ∞ -category. Then the functor*

$$\widetilde{\mathcal{F}}(\mathcal{C}) \longrightarrow N\Delta^{\text{op}}$$

is a cocartesian fibration, and the map $\widetilde{\mathcal{F}}(\mathcal{C}) \rightarrow \mathcal{S}(\mathcal{C})$ defined above is a fiberwise equivalence over $N\Delta^{\text{op}}$. Dually, if \mathcal{C} is a coWaldhausen ∞ -category, then the functor

$$\widetilde{\mathcal{F}}^{\vee}(\mathcal{C}) \longrightarrow N\Delta$$

is a cartesian fibration, and the map $\widetilde{\mathcal{F}}^{\vee}(\mathcal{C}) \rightarrow \mathcal{S}^{\vee}(\mathcal{C})$ defined above is a fiberwise equivalence over $N\Delta$.

5.15. In particular, we can use the equivalence of the previous proposition to endow, for any Waldhausen ∞ -category (respectively, any coWaldhausen ∞ -category) \mathcal{C} , the ∞ -category $\widetilde{\mathcal{F}}(\mathcal{C})$ (respectively, $\widetilde{\mathcal{F}}^{\vee}(\mathcal{C})$) with the structure of a Waldhausen ∞ -category. That is, let us declare a morphism $X \rightarrow Y$ of $\widetilde{\mathcal{F}}(\mathcal{C})$ (respectively, of $\widetilde{\mathcal{F}}^{\vee}(\mathcal{C})$) ingressive (respectively, egressive) just in case its image in $\mathcal{S}(\mathcal{C})$ (respectively, in $\mathcal{S}^{\vee}(\mathcal{C})$) is so. Observe that under this definition, $X \rightarrow Y$ is ingressive (respectively, egressive) just in case each of the morphisms $X(i, j) \rightarrow Y(i, j)$ of \mathcal{C} is so.

Consequently, the constructions $\widetilde{\mathbf{S}}$ and $\widetilde{\mathbf{S}}^{\vee}$ yield functors

$$\widetilde{\mathbf{S}} : \mathbf{Wald}_{\infty} \longrightarrow \text{Fun}(N\Delta^{\text{op}}, \mathbf{Wald}_{\infty}) \quad \text{and} \quad \widetilde{\mathbf{S}}^{\vee} : \mathbf{coWald}_{\infty} \longrightarrow \text{Fun}(N\Delta^{\text{op}}, \mathbf{coWald}).$$

Now suppose \mathcal{C} an exact ∞ -category. Since ambigressive pushouts and ambigressive pullbacks coincide in \mathcal{C} , it follows that there is a canonical equivalence

$$(\widetilde{\mathbf{S}}_* \circ \text{op})|_{\mathbf{Exact}_{\infty}} \simeq (\widetilde{\mathbf{S}}_*^{\vee})|_{\mathbf{Exact}_{\infty}},$$

where $\text{op} : N\Delta^{\text{op}} \xrightarrow{\sim} N\Delta^{\text{op}}$ is the opposite automorphism of $N\Delta^{\text{op}}$. We therefore deduce the following theorem.

THEOREM 5.16. *Suppose \mathcal{E} an ∞ -topos. For any pointed functor $\rho : \mathbf{Cat}_{\infty}^* \rightarrow \mathcal{E}_*$ that preserves filtered colimits, an equivalence $\rho \xrightarrow{\sim} \rho \circ \text{op}$ induces a canonical exact duality on the Goodwillie additivization $D\rho$.*

Proof. The equivalence $\rho \xrightarrow{\sim} \rho \circ \text{op}$, combined with the equivalence $\widetilde{\mathbf{S}}_* \circ \text{op} \simeq \widetilde{\mathbf{S}}_*^{\vee}$, yields an equivalence

$$|\rho \circ \widetilde{\mathbf{S}}_*| \simeq |\rho \circ \widetilde{\mathbf{S}}_* \circ \text{op}| \xrightarrow{\sim} |\rho \circ \text{op} \circ \widetilde{\mathbf{S}}_* \circ \text{op}| \simeq |\rho^{\vee} \circ \widetilde{\mathbf{S}}_*^{\vee}|. \quad \square$$

Applying this result to the functor ι yields the following corollary.

COROLLARY 5.16.1. *Algebraic K-theory admits an exact duality.*

6. Theorem of the Heart

In this section, we show that the Waldhausen K -theory of a stable ∞ -category with a bounded t-structure agrees with the K -theory of its heart. Amnon Neeman has provided an analogous result for his K -theory of triangulated categories [Sch02]; given Neeman’s result, our result here may be alternatively summarized as saying that the Waldhausen K -theory of a stable ∞ -category \mathcal{A} with a bounded t-structure agrees with Neeman’s K -theory of the triangulated homotopy category $\mathcal{T} = h\mathcal{A}$ (denoted $K({}^w\mathcal{T})$ in [Nee05]); this verifies a conjecture posed by Neeman [Nee99, Appendix A] for such stable homotopy theories.

In this section, suppose E a small stable ∞ -category equipped with a bounded t-structure $(E_{\leq -1}, E_{\geq 0})$. Our objective is to prove the following

THEOREM 6.1 (Heart). *The inclusions $E^\heartsuit \hookrightarrow E_{\geq 0}$ and $E_{\geq 0} \hookrightarrow E$ induce equivalences*

$$K(E^\heartsuit) \xrightarrow{\sim} K(E_{\geq 0}) \quad \text{and} \quad K(E_{\geq 0}) \xrightarrow{\sim} K(E) .$$

The fact that the inclusions induce isomorphisms $K_0(E^\heartsuit) \cong K_0(E_{\geq 0}) \cong K_0(E)$ is well known and trivial. Consequently, appealing to the Cofinality theorem of [Bar13, Theorem 10.11] (which states that idempotent completions induce equivalences on the connected cover of K -theory), we may therefore assume that E is idempotent complete. We now set about proving that the higher K -groups of E^\heartsuit and E agree.

Our proof is quite straightforward. The first main tool is the following.

THEOREM 6.2 (Special Fibration theorem, [Bar13, Proposition 10.12]). *Suppose C a compactly generated ∞ -category containing a zero object, suppose $L : C \rightarrow D$ an accessible localization, and suppose the inclusion $D \hookrightarrow C$ preserves filtered colimits. Assume also that the class of all L -equivalences of C is generated (as a strongly saturated class) by the L -equivalences between compact objects. Then L induces a pullback square of spaces*

$$\begin{array}{ccc} K(E^\omega) & \longrightarrow & K(C^\omega) \\ \downarrow & & \downarrow \\ * & \longrightarrow & K(D^\omega) \end{array}$$

where C^ω and D^ω are equipped with the maximal pair structure, and $E^\omega \subset C^\omega$ is the full subcategory spanned by those objects X such that $LX \simeq 0$.

As a consequence of [Lur09a, Corollary 5.5.7.4, Proposition 5.5.7.8], we proved in [Bar13, Corollary 10.12.2] that the functor $\tau_{\leq -1}$ induces a pullback square

$$\begin{array}{ccc} K(E_{\geq 0}) & \longrightarrow & K(E) \\ \downarrow & & \downarrow \\ * & \longrightarrow & K(E_{\leq -1}^{\max}) \end{array}$$

of spaces, where the ∞ -categories that appear are equipped with their maximal Waldhausen structure; i.e., the one in which every morphism is ingressive. (Note that for $E_{\geq 0}$ and E , the Waldhausen structure described in Example 3.4 is the maximal Waldhausen structure, but this is very much not the case for $E_{\leq -1}$.)

We now claim that the K -theory of the maximal Waldhausen ∞ -category $E_{\leq -1}^{\max}$ vanishes. Indeed, we may apply [Bar13, Corollary 8.2.1] to the ∞ -category $E_{\leq -1}$; this will ensure that the functor

$$\Sigma_{E_{\leq -1}}^{\infty} : E_{\leq -1}^{\max} \longrightarrow \widetilde{\mathbf{Sp}}(E_{\leq -1})^{\max}$$

induces an equivalence on K -theory. Note that the suspension functor on $E_{\leq -1}$ is the composite $\tau_{\leq -1} \circ \Sigma_E$. Since the t -structure is bounded, it therefore follows that $\Sigma_{E_{\leq -1}}^{\infty}$ is equivalent to the constant functor at 0.

Warning 6.3. Note that this argument applies only to the maximal Waldhausen ∞ -category $E_{\leq -1}^{\max}$. The K -theory of the Waldhausen structure on $E_{\leq -1}$ of Example 3.4, in which a morphism is ingressive just in case it induces a monomorphism on π_{-1} , will turn out to agree with the K -theory of E .

Consequently, the inclusion $E_{\geq 0} \hookrightarrow E$ induces an equivalence

$$K(E_{\geq 0}) \xrightarrow{\sim} K(E),$$

and we are left with showing that the map $K(E^{\heartsuit}) \longrightarrow K(E_{\geq 0})$ is an equivalence.

For this, we take opposites. The opposite ∞ -category E^{op} is endowed with the dual t -structure, with $(E^{\text{op}})_{\leq -n} = (E_{\geq n})^{\text{op}}$. Consider the functor

$$\tau_{\leq -1} = (\tau_{\geq 1})^{\text{op}} : (E_{\geq 0})^{\text{op}} = (E^{\text{op}})_{\leq 0} \longrightarrow (E^{\text{op}})_{\leq -1} = (E_{\geq 1})^{\text{op}}.$$

Our claim is that it induces a pullback square

$$\begin{array}{ccc} K((E^{\text{op}})^{\heartsuit}) & \longrightarrow & K((E^{\text{op}})_{\leq 0}) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & K((E^{\text{op}})_{\leq -1}^{\max}) \end{array} \tag{*}$$

where the ∞ -categories that appear are equipped with the Waldhausen structure in which a morphism is ingressive just in case it induces a monomorphism (in E^{op}) on π_0 (and in particular, the Waldhausen structure on $(E^{\text{op}})_{\leq -1}$ is maximal).


Of course $K((E^{\text{op}})_{\leq -1}^{\max})$ vanishes just as $K(E_{\leq -1}^{\max})$ vanishes. Consequently, if we verify that (*) is a pullback, we will deduce that the map $K((E^{\text{op}})^{\heartsuit}) \longrightarrow K((E^{\text{op}})_{\leq 0})$ is an equivalence, and in light of Corollary 5.16.1, which allows us to pass between a Waldhausen ∞ -category and its opposite under K -theory, the proof of Theorem 6.1 will be complete.

To prove that (*) is a pullback, we cannot simply appeal to the Special Fibration theorem, because the Waldhausen ∞ -categories that appear have non-maximal pair structures. Instead, we must appeal to what we have called the Generic Fibration theorem II; this is the ∞ -categorical variant of Waldhausen’s celebrated Fibration theorem. Unfortunately, this means that we will have to check some technical hypotheses, but there is no escape.

THEOREM 6.4 (Generic Fibration theorem II, [Bar13, Theorem 9.24]). *Suppose C a Waldhausen ∞ -category, and suppose that wC is a subcategory of C that satisfies the following conditions.*

(6.4.1) *Every equivalence of C lies in wC .*

(6.4.2) The morphisms of wC satisfy (the ∞ -categorical analogue of) Waldhausen’s gluing axiom. That is, for any cofibrations $U \twoheadrightarrow V$ and $X \twoheadrightarrow Y$ and any cube



in which the top and bottom faces are pushout squares, if $U \twoheadrightarrow X$, $V \twoheadrightarrow Y$ and $U' \twoheadrightarrow X'$ all lie in wC , then so does $V' \twoheadrightarrow Y'$.

(6.4.3) There are enough cofibrations in the following sense. The three functors

$$wC \cap C_{\dagger} \hookrightarrow wC, \quad w\text{Fun}(\Delta^1, C) \cap \text{Fun}(\Delta^1, C)_{\dagger} \hookrightarrow w\text{Fun}(\Delta^1, C),$$

and

$$w\mathbf{F}_1(C) \cap \mathbf{F}_1(C)_{\dagger} \hookrightarrow w\mathbf{F}_1(C)$$

are all weak homotopy equivalences, where cofibrations of $\text{Fun}(\Delta^1, C)$ are defined objectwise, and $w\text{Fun}(\Delta^1, C)$ and $w\mathbf{F}_1(C)$ are also defined objectwise.

Now denote by $C^w \subset C$ the full subcategory spanned by those objects X such that the cofibration $0 \twoheadrightarrow X$ lies in wC , and declare a morphism therein to be ingressive just in case it is so in C . Denote by $\mathbf{B}_*(C, wC)$ the simplicial ∞ -category whose ∞ -category of m -simplices is the full subcategory $\mathbf{B}_m(C, wC) \subset \text{Fun}(\Delta^m, C)$ spanned by those sequences of edges

$$X_0 \twoheadrightarrow X_1 \twoheadrightarrow \dots \twoheadrightarrow X_m$$

with the property that each $X_i \twoheadrightarrow X_j$ lies in wC ; declare a morphism of $\mathbf{B}_m(C, wC)$ to be a cofibration just in case it is so objectwise. Then C^w is a Waldhausen ∞ -category, and $\mathbf{B}_*(C, wC)$ is a simplicial Waldhausen ∞ -category, and the obvious functors induce a fiber sequence of spaces.

$$\begin{array}{ccc}
 K(C^w) & \longrightarrow & K(C) \\
 \downarrow & & \downarrow \\
 * & \longrightarrow & |K(\mathbf{B}_*(C, wC))|
 \end{array}$$

Here of course $|\cdot|$ denotes the geometric realization.

We apply this theorem where C is the Waldhausen ∞ -category $(E^{\text{op}})_{\leq 0}$, and wC is the subcategory $w((E^{\text{op}})_{\leq 0})$ consisting of those morphisms $X \twoheadrightarrow Y$ such that the induced morphism $\tau_{\leq -1}X \twoheadrightarrow \tau_{\leq -1}Y$ is an equivalence. It is an easy matter to check the gluing axiom in this setting. To show that we have enough cofibrations, let us employ the following construction: factor any morphism $f : X \twoheadrightarrow Y$ of wC as

$$X \twoheadrightarrow X_f \twoheadrightarrow Y,$$

where X_f is the fiber of the natural map $Y \twoheadrightarrow \tau_{\leq 0}(Y/X)$, where Y/X is the cofiber of f . This construction defines a deformation retraction $\text{Fun}(\Delta^1, wC) \twoheadrightarrow \text{Fun}(\Delta^1, wC \cap C_{\dagger})$ of the

inclusion $\text{Fun}(\Delta^1, wC \cap C_+) \hookrightarrow \text{Fun}(\Delta^1, wC)$. It follows from the functoriality of this construction that it also defines a deformation retraction of the inclusions

$$\text{Fun}(\Delta^1, w \text{Fun}(\Delta^1, C) \cap \text{Fun}(\Delta^1, C)_+) \hookrightarrow \text{Fun}(\Delta^1, w \text{Fun}(\Delta^1, C)),$$

and

$$\text{Fun}(\Delta^1, w\mathbf{F}_1(C) \cap \mathbf{F}_1(C)_+) \hookrightarrow \text{Fun}(\Delta^1, w\mathbf{F}_1(C)).$$

Consequently, these maps are all homotopy equivalences.

So Theorem 6.4 now ensures that we have a pullback square

$$\begin{array}{ccc} K((E^{\text{op}})^{\heartsuit}) & \longrightarrow & K((E^{\text{op}})_{\leq 0}) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & |K(\mathbf{B}_*((E^{\text{op}})_{\leq 0}, w(E^{\text{op}})_{\leq 0}))| \end{array}$$

and it remains only to identify the cofiber term. Now it follows from [Bar13, Proposition 10.10] that, in the situation of Theorem 6.4, the geometric realization $|K(\mathbf{B}_*(C, wC))|$ can be identified with $\Omega|w\mathbf{S}_*(C)|$, where $w\mathbf{S}_m(C) \subset \mathbf{S}_m(C)$ is the subcategory whose morphisms are objectwise in wC . In our case, recall, these are morphisms that induce equivalences after applying $\tau_{\leq -1}$. Consequently, to identify $K((E^{\text{op}})_{\leq -1}^{\max})$ with the geometric realization

$$|K(\mathbf{B}_*((E^{\text{op}})_{\leq 0}, w(E^{\text{op}})_{\leq 0}))|,$$

it is enough to show that the functor

$$p : w\mathbf{S}_m((E^{\text{op}})_{\leq 0}) \longrightarrow \iota\mathbf{S}_m((E^{\text{op}})_{\leq -1}^{\max})$$

induced by $\tau_{\leq -1}$ is a weak homotopy equivalence. Let us show that the inclusion

$$i : \iota\mathbf{S}_m((E^{\text{op}})_{\leq -1}^{\max}) \hookrightarrow \iota\mathbf{S}_m((E^{\text{op}})_{\leq 0}) \subset w\mathbf{S}_m((E^{\text{op}})_{\leq 0})$$

defines a homotopy inverse. Since $\tau_{\leq -1}$ is a localization functor, we obtain for every $m \geq 0$ a natural equivalence $p \circ i \simeq \text{id}$. In the other direction, the natural transformation $\text{id} \rightarrow \tau_{\leq -1}$ induces a natural transformation $\text{id} \rightarrow i \circ p$ (which of course induces an equivalence after applying $\tau_{\leq -1}$ and thus is an objectwise morphism of $w((E^{\text{op}})_{\leq 0})$). Hence i and p are homotopy inverse. \square

Amnon Neeman’s Theorem of the Heart now implies the following, which verifies some cases of his conjecture [Nee99, Conjecture A.5].

COROLLARY 6.4.1. *For any idempotent-complete stable ∞ -category \mathcal{A} , if the triangulated homotopy category $\mathcal{T} = h\mathcal{A}$ admits a bounded t -structure, then we have canonical equivalences*

$$K(\mathcal{A}) \simeq K(\mathcal{T}^{\heartsuit}) \simeq K({}^w\mathcal{T}).$$

7. Application: Abelian models for the algebraic G -theory of schemes

A trivial application of the Theorem of the Heart applies to show that the K -theory of an abelian category A agrees with the K -theory of its bounded derived ∞ -category $D^b(A)$ [TT90, 1.11.7]. However, tilting theory provides other bounded t -structures on the ∞ -category $D^b(A)$. The K -theory of the heart of any of these t -structures will agree with the K -theory of A . Let us explore one class of examples now.

Example 7.1. Suppose X a noetherian scheme equipped with a dualizing complex $\omega_X \in D^b(\mathbf{Coh}(X))$. Then Arinkin and Bezrukavnikov [AB10], following Deligne, construct a family of t-structures on $D^b(\mathbf{Coh}(X))$ in the following manner. (Here we use *cohomological* grading conventions, to maintain compatibility with [AB10].) Write X^{top} for the underlying topological space of X , and define $\dim : X^{\text{top}} \rightarrow \mathbf{Z}$ a map such that $i_x^! \omega_X$ is concentrated in degree $-\dim(x)$. Suppose $p : X^{\text{top}} \rightarrow \mathbf{Z}$ a function, called a *perversity*, such that for any points $z, x \in X^{\text{top}}$ such that $z \in \overline{\{x\}}$, one has

$$p(x) \leq p(z) \leq p(x) + \dim(x) - \dim(z).$$

Let $D^{p \geq 0} \subset D^b(\mathbf{Coh}(X))$ be the full subcategory spanned by those complexes F such that for any point $x \in X^{\text{top}}$, one has $i_x^! F \in D^{\geq p(x)}(\mathcal{O}_{X,x})$; dually, let $D^{p \leq 0} \subset D^b(\mathbf{Coh}(X))$ be the full subcategory spanned by those complexes F such that for any point $x \in X^{\text{top}}$, one has $i_{x,*} F \in D^{\leq p(x)}(\mathcal{O}_{X,x})$. Then $(D^{p \geq 0}, D^{p \leq 0})$ define a bounded t-structure on $D^b(\mathbf{Coh}(X))$ [AB10, Theorem 3.10].

The algebraic K -theory of the heart $D^{p,\heartsuit}$ of this t-structure now coincides with the G -theory of X . Let us list two special cases of this.

(7.1.1) Suppose S a set of prime numbers. Let \mathcal{E}_S be the full subcategory of $D^b(\mathbf{Coh}(\mathbf{Z}))$ generated under extensions by the objects

$$\mathbf{Z}, \quad \{\mathbf{Z}/p \mid p \in S\}, \quad \{\mathbf{Z}/p[1] \mid p \notin S\}.$$

Then \mathcal{E}_S is an abelian category whose K -theory coincides with the K -theory of \mathbf{Z} .

(7.1.2) For any noetherian scheme with a dualizing complex $\omega_X \in D^b(\mathbf{Coh}(X))$, the K -theory of the abelian category of *Cohen–Macaulay complexes* (i.e., those complexes $F \in D^b(\mathbf{Coh}(X))$ such that the complex

$$\mathbf{D}F := \mathbf{R}\text{Mor}_{\mathcal{O}_X}(F, \omega_X)$$

is concentrated in degree 0, [YZ06, § 6]) agrees with the G -theory of X .

8. Application: a theorem of Blumberg–Mandell

In this section, we give a new proof of the theorem of Blumberg and Mandell [BM08] that establishes a localization sequence

$$K(\pi_0 E) \rightarrow K(e) \rightarrow K(E)$$

for any suitable even periodic E_1 ring spectrum E with $\pi_0 E$ regular noetherian, where e denotes the connective cover of E . In light of [Bar13, Proposition 13.16], the key point is the identification of the fiber term; this is the subject of this section.

Recall [Lur12, Proposition 8.2.5.16] that a connective E_1 ring Λ is said to be *left coherent* if $\pi_0 \Lambda$ is left coherent as an ordinary ring, and if for any $n \geq 1$, the left $\pi_0 \Lambda$ -module $\pi_n \Lambda$ is finitely presented.

A left module M over a left coherent E_1 ring Λ is *almost perfect* just in case $\pi_m M = 0$ for $m \ll 0$ and for any n , the left $\pi_0 \Lambda$ -module $\pi_n M$ is finitely presented [Lur12, Proposition 8.2.5.17].

DEFINITION 8.1. Suppose Λ a left coherent E_1 ring, and suppose M a left Λ -module. We say that M is *truncated* if $\pi_m M = 0$ for $m \gg 0$. We say that M is *coherent* if it is almost perfect and truncated. Write $\mathbf{Coh}_\Lambda^\ell \subset \mathbf{Mod}_\Lambda^\ell$ for the full subcategory spanned by the coherent modules, and write $G(\Lambda)$ for $K(\mathbf{Coh}_\Lambda^\ell)$.

Warning 8.2. In general, it is not the case that a perfect Λ -module is coherent; consequently, the usual Poincaré duality map $K \rightarrow G$ for discrete rings does not have an obvious analogue for E_1 rings.

It turns out that G -theory is not a new invariant of E_1 rings, since we have the following new proof of the Dévissage theorem of Blumberg and Mandell [BM08].

PROPOSITION 8.3. *For any coherent E_1 ring Λ , the inclusion $N\mathbf{Mod}_{\pi_0\Lambda}^{\ell, \text{fp}} \hookrightarrow \mathbf{Coh}_{\Lambda}^{\ell}$ of the nerve of the category of finitely presented $\pi_0\Lambda$ -modules induces an equivalence*

$$G(\pi_0\Lambda) \xrightarrow{\sim} G(\Lambda).$$

Proof. We note that $\mathbf{Coh}_{\Lambda}^{\ell}$ is the full subcategory of the ∞ -category of almost perfect Λ -modules spanned by those that are bounded for the t-structure given by [Lur12, Proposition 8.2.5.18]. Furthermore, [Lur12, Proposition 8.2.5.11(2)] applies to ensure that $\mathbf{Coh}_{\Lambda}^{\ell}$ is idempotent complete. Consequently, the Theorem of the Heart applies, and the proof is complete once one observes that the heart $\mathbf{Coh}_{\Lambda}^{\ell, \heartsuit}$ can be identified with $N\mathbf{Mod}_{\pi_0\Lambda}^{\ell, \text{fp}}$ [Lur12, Remark 8.2.5.19]. \square

Consequently, from the point of view of ‘brave new algebra’, G -theory is a relatively coarse invariant.

Now we hope to compare the G -theory of an E_1 ring to the K -theory of the ∞ -category of truncated perfect modules. This requires a weak regularity hypothesis, which we formulate now.

DEFINITION 8.4. Let us say that a coherent E_1 ring Λ is *almost regular* if any truncated, almost perfect Λ -module has finite Tor-dimension.

Example 8.5. If the graded ring $\pi_*\Lambda$ has finite Tor-dimension (e.g., if $\pi_*\Lambda$ is a regular noetherian ring), then the Tor spectral sequence ensures that Λ is almost regular.

The following result is now an immediate consequence of [Lur12, Proposition 8.2.5.23(4)].

PROPOSITION 8.6. *Suppose Λ a coherent E_1 ring that is almost regular. Then the ∞ -category $\mathbf{Perf}_{\Lambda}^{\ell, b}$ of perfect truncated left Λ -modules coincides with the ∞ -category $\mathbf{Coh}_{\Lambda}^{\ell}$ of coherent left Λ -modules.*

Assembling all this, we obtain the following formulation of the Localization theorem of [BM08].

THEOREM 8.7. *Suppose Λ a coherent E_1 ring spectrum that is almost regular, and suppose $L : \mathbf{Mod}^{\ell}(\Lambda) \rightarrow \mathbf{Mod}^{\ell}(\Lambda)$ smashing Bousfield localization with the property that a left Λ -module M is L -acyclic just in case it is truncated. (Note that in this case, L is automatically a finite localization functor.) Then there is a fiber sequence of spaces*

$$G(\pi_0\Lambda) \rightarrow K(\Lambda) \rightarrow K(L(\Lambda)).$$

Of course when $\pi_0\Lambda$ is regular, the fiber term can be identified with $K(\pi_0\Lambda)$.

Example 8.8. Here are some examples of fiber sequences resulting from this theorem.

(8.8.1) Consider the Adams summand L with its canonical E_{∞} structure; its connective cover ℓ admits a canonical E_{∞} as well [BR08]. The fiber sequence above becomes

$$K(\mathbf{Z}) \rightarrow K(\ell) \rightarrow K(L).$$

(8.8.2) Similarly, one can use the E_∞ structure on complex K -theory KU and on its connective cover to obtain

$$K(\mathbf{Z}) \longrightarrow K(ku) \longrightarrow K(KU).$$

(8.8.3) For any perfect field k of characteristic $p > 0$ and any formal group Γ of height n over k , consider the Lubin–Tate spectrum $E(k, \Gamma)$ with its canonical E_∞ structure and its connective cover $e(k, \Gamma)$ with its induced E_∞ structure [BR08]. In this case, the fiber sequence above becomes

$$K(\mathbf{W}(k)[[u_1, \dots, u_{n-1}]]) \longrightarrow K(e(k, \Gamma)) \longrightarrow K(E(k, \Gamma)).$$

(8.8.4) Given any E_1 structure on Morava K -theory $K(n)$ and a compatible one on its connective cover $k(n)$, one has a fiber sequence

$$K(\mathbf{F}_p) \longrightarrow K(k(n)) \longrightarrow K(K(n)).$$

9. Application: G -theory of spectral Deligne–Mumford stacks

The purpose of this final very brief section is simply to note that the G -theory of locally noetherian spectral Deligne–Mumford stacks is insensitive to derived structure.

DEFINITION 9.1. For any spectral Deligne–Mumford stack \mathcal{X} , we write $\mathbf{Coh}(\mathcal{X})$ for the stable ∞ -category of coherent sheaves on \mathcal{X} [Lur11, Definition 2.6.20], i.e., those quasicoherent sheaves that are almost perfect and locally truncated. Write $G(\mathcal{X})$ for the algebraic K -theory of $\mathbf{Coh}(\mathcal{X})$.

Now the Theorem of the Heart, combined with [Lur11, Remark 2.3.20], instantly yields the following proposition.

PROPOSITION 9.2. For any locally noetherian spectral Deligne–Mumford stack \mathcal{X} with underlying ordinary Deligne–Mumford stack \mathcal{X}_0 , the embedding

$$N\mathbf{Coh}(\mathcal{X}_0) \hookrightarrow \mathbf{Coh}(\mathcal{X})$$

induces an equivalence

$$G(\mathcal{X}_0) \xrightarrow{\sim} G(\mathcal{X}).$$

Roughly speaking, just as G -theory is invariant under ordinary nilpotent thickenings, it turns out that G -theory is invariant under *derived* nilpotent thickenings as well.

ACKNOWLEDGEMENTS

I thank Andrew Blumberg and Mike Mandell for very helpful conversations about this paper. I also thank Dustin Clausen for helpful remarks about a previous version of this paper. I am very grateful to Benjamin Antieau for noticing that overzealous revision botched the proof of the main theorem in an earlier version of this paper. Finally, I thank the anonymous referee for a genuinely helpful report.

REFERENCES

- AB10 D. Arinkin and R. Bezrukavnikov, *Perverse coherent sheaves*, Mosc. Math. J. **10** (2010), 3–29; 271; MR 2668828 (2011g:14040).
- BR08 A. Baker and B. Richter, *Uniqueness of E_∞ structures for connective covers*, Proc. Amer. Math. Soc. **136** (2008), 707–714; MR 2358512 (2008m:55017).

- Bar13 C. Barwick, *On the algebraic K-theory of higher categories*, J. Topol., to appear, [arXiv:1204.3607](https://arxiv.org/abs/1204.3607).
- BL14 C. Barwick and T. Lawson, *Regularity of structured ring spectra and localization in K-theory*, Preprint (2014), [arXiv:1402.6038](https://arxiv.org/abs/1402.6038).
- BS11 C. Barwick and C. Schommer-Pries, *On the unicity of the homotopy theory of higher categories*, Preprint (2011), [arXiv:1112.0040](https://arxiv.org/abs/1112.0040).
- BGT13 A. J. Blumberg, D. Gepner and G. Tabuada, *A universal characterization of higher algebraic K-theory*, Geom. Topol. **17** (2013), 733–838, doi:[10.2140/gt.2013.17.733](https://doi.org/10.2140/gt.2013.17.733).
- BM08 A. J. Blumberg and M. A. Mandell, *The localization sequence for the algebraic K-theory of topological K-theory*, Acta Math. **200** (2008), 155–179; [MR 2413133](https://doi.org/10.2307/2413133) (2009f:19003).
- Joy08a A. Joyal, *Notes on quasi-categories*, Preprint (2008).
- Joy08b A. Joyal, *The theory of quasi-categories and its applications*, *Advanced Course on Simplicial Methods in Higher Categories*, Vol. 2, Quaderns, vol. 45 (Centre de Recerca Matemàtica, Barcelona, 2008).
- Kel90 B. Keller, *Chain complexes and stable categories*, Manuscripta Math. **67** (1990), 379–417; [MR 1052551](https://doi.org/10.1007/BF01232551) (91h:18006).
- Lur09a J. Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170 (Princeton University Press, Princeton, NJ, 2009); [MR 2522659](https://doi.org/10.1215/00141801-2010-018) (2010j:18001).
- Lur09b J. Lurie, *$(\infty, 2)$ -categories and the Goodwillie calculus I*, Preprint (2009).
- Lur11 J. Lurie, *Derived algebraic geometry VIII. Quasi-coherent sheaves and Tannaka duality theorems*, Preprint (2011).
- Lur12 J. Lurie, *Higher algebra*, Preprint (2012).
- Nee97a A. Neeman, *K-theory for triangulated categories. I(A). Homological functors*, Asian J. Math. **1** (1997), 330–417; [MR 1491990](https://doi.org/10.1142/S172894529700088a) (99m:18008a).
- Nee97b A. Neeman, *K-theory for triangulated categories. I(B). Homological functors*, Asian J. Math. **1** (1997), 435–529; [MR 1604910](https://doi.org/10.1142/S172894529700088b) (99m:18008b).
- Nee98a A. Neeman, *K-theory for triangulated categories. II. The subtlety of the theory and potential pitfalls*, Asian J. Math. **2** (1998), 1–125; [MR 1656552](https://doi.org/10.1142/S17289452000019008) (2000m:19008).
- Nee98b A. Neeman, *K-theory for triangulated categories. III(A). The theorem of the heart*, Asian J. Math. **2** (1998), 495–589; [MR 1724625](https://doi.org/10.1142/S17289452980004625).
- Nee99 A. Neeman, *K-theory for triangulated categories. III(B). The theorem of the heart*, Asian J. Math. **3** (1999), 557–608; [MR 1793672](https://doi.org/10.1142/S1728945299000672).
- Nee00a A. Neeman, *K-theory for triangulated categories $\mathfrak{3}_{\frac{1}{2}}$. A. A detailed proof of the theorem of homological functors*, K-Theory **20** (2000), 97–174; special issues dedicated to Daniel Quillen on the occasion of his sixtieth birthday, Part II; [MR 1798824](https://doi.org/10.1007/s00202002b18014) (2002b:18014).
- Nee00b A. Neeman, *K-theory for triangulated categories $\mathfrak{3}_{\frac{1}{2}}$. B. A detailed proof of the theorem of homological functors*, K-Theory **20** (2000), 243–298; special issues dedicated to Daniel Quillen on the occasion of his sixtieth birthday, Part III; [MR 1798828](https://doi.org/10.1007/s00202002b18015) (2002b:18015).
- Nee01 A. Neeman, *K-theory for triangulated categories $\mathfrak{3}_{\frac{3}{4}}$: a direct proof of the theorem of the heart*, K-Theory **22** (2001), 1–144; [MR 1828612](https://doi.org/10.1007/s00202002f19004) (2002f:19004).
- Nee05 A. Neeman, *The K-theory of triangulated categories*, Handbook of K-Theory, vols 1, 2 (Springer, Berlin, 2005), 1011–1078; [MR 2181838](https://doi.org/10.1007/978-3-540-26032-9_33) (2006g:19004).
- Sch02 M. Schlichting, *A note on K-theory and triangulated categories*, Invent. Math. **150** (2002), 111–116; [MR 1930883](https://doi.org/10.1007/s00202003h18015) (2003h:18015).
- TT90 R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, in *The Grothendieck festschrift, Vol. III*, Progress in Mathematics, vol. 88 (Birkhäuser, Boston, MA, 1990), 247–435; [MR 92f:19001](https://doi.org/10.1007/978-1-4612-1170-1_11).

- Toe05 B. Toën, *Vers une axiomatisation de la théorie des catégories supérieures*, *K-Theory* **34** (2005), 233–263; [MR 2182378](#) (2006m:55041).
- Wal85 F. Waldhausen, *Algebraic K-theory of spaces*, in *Algebraic and geometric topology* (New Brunswick, NJ, 1983), *Lecture Notes in Mathematics*, vol. 1126 (Springer, Berlin, 1985), 318–419; [MR 86m:18011](#).
- YZ06 A. Yekutieli and J. J. Zhang, *Dualizing complexes and perverse sheaves on noncommutative ringed schemes*, *Selecta Math. (N.S.)* **12** (2006), 137–177; [MR 2244264](#) (2008d:14004).

Clark Barwick clarkbar@gmail.com

Department of Mathematics, Massachusetts Institute of Technology,
Building 2, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA