

University of Kentucky, Lexington, KY ; ³Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY and ⁴Department of Internal Medicine, College of Medicine, University of Kentucky, Lexington, KY

OBJECTIVES/GOALS: To date, no study has jointly profiled RNA isoforms and modifications associated with Alzheimer's disease neuropathological changes (ADNC) in a human cohort. Therefore, the overall objective of this study is to identify and characterize RNA isoforms and modifications that associate with ADNC by leveraging long-read direct RNA sequencing. **METHODS/STUDY POPULATION:** This translational study will analyze human prefrontal cortex tissue from a well-defined cohort of 12 individuals. The cohort consists of six individuals with severe Alzheimer's disease neuropathological changes and six age- and sex-matched non-demented controls. Leveraging long-read direct RNA sequencing, which uniquely quantifies both RNA isoform and modifications simultaneously, can provide novel insights into how severe Alzheimer's disease neuropathological changes drive changes in RNA isoform expression patterns and alter RNA modifications contributing to isoform dysregulation. **RESULTS/ANTICIPATED RESULTS:** My preliminary data from human prefrontal cortex (n=2) revealed pseudouridine sites, 1 of ~170 known RNA modifications, in Alzheimer's disease-risk genes (APP, CLU, SORT1, and BIN1), supporting the need to map RNA modifications across Alzheimer's disease-relevant isoforms. My central hypothesis is that severe Alzheimer's disease neuropathological changes (ADNCs) drive transcriptomic and epitranscriptomic dysregulation by altering RNA isoforms. Thus, my two working hypotheses are 1) severe ADNC drives transcriptomic dysregulation of RNA isoforms and 2) severe ADNC drives epitranscriptomic dysregulation of RNA modifications. **DISCUSSION/SIGNIFICANCE OF IMPACT:** This project will be the first to jointly profile RNA isoforms and modifications accounting for Alzheimer's disease neuropathological changes, offering candidate regulatory targets for therapeutic development. Importantly, this research will equip me with an expertise in RNA biology, neurodegenerative diseases, and bioinformatics.

315

When clinical decision support falls short: Lessons from piloting a patient-centered prescribing tool for post-operative opioids at discharge

Shraddha Pandey, Samuel O'Brien, Colleen P. Flanagan, Tasce Bongiovanni, Hannah Decker, Jory Purvis Zhonghui Guan and Elizabeth Wick
University Of California, San Francisco

OBJECTIVES/GOALS: A clinical decision support tool (CDS) with an individualized prescribing approach was piloted to determine impact on opioid overprescribing rates for surgical patients at discharge. As a follow-up, clinicians were interviewed to determine current challenges to opioid prescribing and required components of an effective CDS for discharge opioids. **METHODS/STUDY POPULATION:** In the EHR-based randomized control trial, patients admitted to a qualifying surgical service for 24+ hours

postoperatively were randomized to the individualized CDS tool with a taper based on inpatient opioid use or the standard discharge order set. Performance of the CDS tool was assessed by rate of opioid mismatch, that is, the difference between median inpatient and daily discharge dose in morphine milligram equivalents (MMEs). Pilot results were further investigated using a sequential explanatory design. Semi-structured interviews were held with discharging clinicians. A purposive sampling strategy with snowball sampling was used to identify providers. Data saturation was used to determine sample sufficiency. Interview transcripts were coded and thematically analyzed, with 40% group coded. **RESULTS/ANTICIPATED RESULTS:** Among 61 randomized patients, median discharge MME (30 [IQR 22.5, 45] remained four times the median inpatient MME (7.5 [IQR 0, 30]) after the pilot period. Nine nurse practitioners and six general surgery residents were interviewed (n=15). Perceived challenges to opioid prescribing centered on three themes: limited protocolization for standard cases; lack of workflow efficiency; and systemic barriers (patient insurance and pharmacy). Clinicians identified at least one useful component of the tool for their current practice. Crucial CDS tool components identified by prescribing clinicians included seamless integration with the electronic health record, adequate clinician training, effective communication of treatment plan with the patient, and clinician buy-in to overcome current prescribing culture. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Multiple patient- and system-related factors impede standardization of opioid prescribing for postoperative patients. CDS tools in this area are a well-received idea, and the tool is now being redesigned to promote effective integration into clinician workflow with multiple stages of intervention, prior to resuming the randomized control trial.

316

Frontiers Trial Finder: Implementing a mobile platform integrated with a Clinical Trial Management System to enhance accessibility and engagement

Hanlue Kuo, Dharani Buddana, Dinesh Pal Mudarantakam and Jeffrey A. Thompson
University of Kansas Medical Center

OBJECTIVES/GOALS: A key barrier to clinical trial (CT) access is knowing which trials are enrolling. Frontiers has developed a Trial Finder mobile app that displays streamlined, up-to-date trial availability that can be integrated to other institutions. We're gathering usability feedback and exploring interest from other institutions in adopting the tool. **METHODS/STUDY POPULATION:** The Frontiers Trial Finder app is a unique tool that extracts trial data and status directly from a Clinical Trial Management System (CTMS). It includes helpful links to ClinicalTrials.gov and provides ways to connect with the study team. The app updates automatically every night, storing the extracted data in a separate stream that can be accessed through the app or online. Its interface is custom-designed in-house and fully customizable, allowing the integration of additional resources such as FAQs, links to partner affiliates, and more. During onboarding, we identified two key efforts: first, organizing the trial directory by defining parent groups and subgroups; and