The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

Cochlea	HANS ENGSTRÖM
THE INNERVATION OF THE ORGAN OF CORTI	H. SPOENDLIN
THE HUMAN COCHLEA DURING DEVELOPMENT AND AGEING	Göran Bredberg
THE ULTRASTRUCTURE AND THE FUNCTIONS OF THE SACCUS ENDOLYMPHATICUS AND ITS DECOMPRESSION IN MÉNIÈRE'S DISEASE.	PETER ADLINGTON
A CONTROLLED STUDY EVALUATION OF ADENO- TONSILLECTOMY IN CHILDREN PETER ADLIN	STUART R. MAWSON, GTON and MAIR EVANS
BLOOD BLISTERS AND THE OESOPHAGEAL CAST	N. J. BADHAM
CLINICAL RECORDS—	
HAEMORRHAGE FOLLOWING TONSILLECTOMY AND ADENOIDECTOMY (A REVIEW OF 18,184 OPERATIONS)	R. G. WILLIAMS
Oesophageal Candidiasis and its Radio- logical Diagnosis	J. E. DELAHUNTY
Unsuspected Thrombocytopenia Causing Massive Haemorrhage at Operation .	G. J. CONRAD
A GIANT TONSILLOLITH	L. H. HIRANANDANI
A Case of Benign Tumour of Tonsil	S. KAMESWARAN

GENERAL NOTES

London

Headley Brothers

109 Kingsway WC2

The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

- 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.
- 4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to Headley Brothers, 109 Kingsway, London, WC2. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.
- 5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.
- 6. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2.
- 7. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.
- 8. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.
- 9. All subscriptions, advertising and business communications should be sent to the publishers, Headley Brothers, 109 Kingsway, London, WC2.

United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, Headley Brothers, 109 Kingsway, London, WC2, England.

© Journal of Laryngology and Otology, 1967

CONTENTS

	PAGE
THE ULTRASTRUCTURE OF THE SENSORY CELLS OF THE COCHLEA. Hans Engström (Gothenburg, Sweden)	687
The Innervation of the Organ of Corti. H. Spoendlin (Zürich)	717
THE HUMAN COCHLEA DURING DEVELOPMENT AND AGEING. Göran Bredberg (Göteborg, Sweden)	739
THE ULTRASTRUCTURE AND THE FUNCTIONS OF THE SACCUS ENDOLYM- PHATICUS AND ITS DECOMPRESSION IN MÉNIÈRE'S DISEASE.	737
Peter Adlington (London)	759
A CONTROLLED STUDY EVALUATION OF ADENO-TONSILLECTOMY IN CHILDREN. Stuart R. Mawson, Peter Adlington and Mair Evans (London).	777
BLOOD BLISTERS AND THE OESOPHAGEAL CAST. N. J. Badham (Leicester)	791
Clinical Records—	
HAEMORRHAGE FOLLOWING TONSILLECTOMY AND ADENOIDECTOMY	
(A REVIEW OF 18,184 OPERATIONS). R. G. Williams (Hull)	805
Oesophageal Candidiasis and its Radiological Diagnosis. J. E. Delahunty (Oxford)	809
Unsuspected Thrombocytopenia Causing Massive Haemorrhage	809
AT OPERATION. G. J. Conrad (London)	815
A GIANT TONSILLOLITH. L. H. Hiranandani (Bombay)	819
A Case of Benign Tumour of Tonsil. S. Kameswaran (Madurai,	
India)	823
General Notes	825



Instrumentation for Audiology and Psychoacoustics

istributors of the World's leading makes





Our Range includes:

Portable Audiometers
Clinical Audiometers
Lekesy Audiometers
Speech Audiometers
Acoustic Impedance Meters
Zwislocki Bridge
Psychogalvanometer
Auditory Trainers
Group Hearing Alds
Individual Hearing Alds for
School use
Silicon Based Aural Impression
Materials
Audiometric Examination
Rooms and many other items



GRASON-STADLER






Please let us know your needs

G. C. Johnson, Leyton Instrument Company Limited, 54 Albert Road, Gaversham, Reading, Berks Reading 72293

BÉKÉSY Accessory



For the very first time a fully automatic unit has been made available for a Clinic Audiometer, making the combined units into a true BÉKÉSY Audiometer.


This automatic accessory is being made for the PETERS SPD/5 CLINIC AUDIOMETER, which has a continuous frequency range and continuously variable attenuation. The result is therefore a true Békésy audiometer.

It can be attached to or detached from the audiometer in under one minute so that the two units together offer the facilities of an advanced clinic audiometer or Békésy audiometry at will.

It can be fitted to any existing SPD/5 Audiometer.

Please write for full details and demonstration to:

ALFRED PETERS & SONS LIMITED 51 GELL STREET, SHEFFIELD 3



THE DROP

THAT SOOTHES THE INFECTED EAR

'Otosporin' rapidly destroys virtually all bacteria commonly found in ear infections. reduces irritation, inflammation and pain. 'Otosporin' brand Drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration. Issued in bottles of 5 ml.

'OTOSPORIN'



Burroughs Wellcome & Co. London (The Wellcome Foundation Ltd.)

When the patient needs a hearing aid-

Whom do you recommend?

INGRAMS, of course, the firm with the different makes to choose from.

The usual trial periods are available for all patients.

INGRAMS

2 Shepherd Street, London, W.1. Tel: HYDe Park 9041.

BRANCHES AT

Aberdeen—461 Union Street. Tel: Aberdeen 29632

Birmingham—Five Ways Shopping Centre. Tel: Midland 4314

Brighton-5 Duke Street. Tel: Brighton 28699

Croydon—11 Essex House, George Street. Tel: Croydon 4663


Edinburgh-17 Stafford Street. Tel: Caledonian 1942

Glasgow-38 Bath Street. Tel: Douglas 7710

Manchester—Peter House, St. Peter Square, Oxford St.

Tel: Central 5711

FOR CLINICAL AUDIOMETRY



Audiometer Booths

prefabricated to ensure correct acoustic environment for threshold audiometry, research, etc. Available in standard and special sizes.

Manual Audiometers

with narrow band masking for highest accuracy and extended resolution. Battery-operated transistor models also available.



Automatic **Audiometers**


As exclusive U.K. distributors, we are able to offer Rudmose Self-Recording Audiometers from the most simple to the most comprehensive Diagnostic Békésy



Calibration Service, Screening, **Speech & Industrial Audiometers**

Please write for technical literature to meet your needs

HEARING ADVISORY SERVICE LTD 80 New Bond St, London, W1 Tel: 01-493 9888



WHEN A HEARING AID IS RECOMMENDED

ONLY AMPLIVOX OFFERS THIS HEARING AID SERVICE

The design and manufacture of a complete range with individual fitting facilities.

Best of other makes in addition to Amplivox aids—Amplivox hearing aids are ONLY available from Amplivox.

Highest degree of training of any hearing aid organisation.

Master Hearing Aid evaluations.

Expert follow-up and rehabilitation advice.

14 branch offices with while-you-wait service.

FOR YOUR PATIENTS' GREATEST SATISFACTION
SPECIFY AMPLIVOX

AMPLIVOX


World's most experienced and complete hearing service


AMPLIVOX HEARING ADVISORY SERVICE 80 NEW BOND STREET LONDON, W.1 TEL: 01-493 9888


NEW FOR OTITIS EXTERNA AND CHRONIC OTITIS MEDIA

FOR INFLAMMATION AND INFECTION IN THE EAR

ANTI-INFLAMMATORY ANTI-BACTERIAL ANTIFUNGAL







۵

Δ

LOCORTEN-VIOFORM EARDROPS

flumethasone pivalate 0.02% iodochlorhydroxyquinoline 1%

LOCORTEN-VIOFORM EARDROPS

7.5 ml, in controlled-drop dispenser

LOCORTEN-VIOFORM EARDROPS

CIBA

Basic NHS price 9s. 8d

L.B.D/f

ORBENIN and its success in Gram-Positive Infections

Otitis media Otitis externa

Boils Carbuncles Infected dermatoses

Pneumonia Lung abscess Acute bronchitis

Osteomyelitis Osteitis

Tonsillitis Pharyngitis

Post-operative wound infections Burns Skin graft protection

Septicaemia Acute endocarditis

Staphylococcal enterocolitis

Staphylococcal urinary tract infections

Staphylococcal meningitis

RBENIN (cloxacillin sodium B.P.) is penicillin active against virtually all Gram-positive organisms, including taphylococci, B-haemolytic streptococci nd pneumococci, irrespective of their esistance to other antibiotics.


t is encouraging that the widespread ise of Orbenin in hospitals has not esulted in the emergence of resistant trains.

More than 300 references in the World iterature testify to the excellent clinical esults obtained with Orbenin in a wide variety of Gram-positive infections.

Dosage:

Two capsules (500mg.) or one vial (250mg.) q.i.d.

Availability:







Detailed literature is available on request.



Orbenin (cloxacillin sodium B.P.) (regd.) is a product of British research at Beecham Research Laboratories

Brentford, England. Telephone: ISLeworth 4111



THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$18.00 per year Canada \$19.00 per year Foreign \$19.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D. EDITOR

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.




Please mention The Journal of Laryngology and Otology when replying to advertisements

54 Albert Rd., Caversham, Reading, Berks. Reading 72293

KAMPLEX AUDIOMETRIC EQUIPMENT

AUDIOMETERS



SCREENING PORTABLE TA16
DIAGNOSTIC PORTABLE TA15

CLINIC AUDIOMETER
DOUBLE CHANNEL

DA1 DA2


AUTOMATIC AUDIOMETER
AB1

CLINIC AUDIOMETER DA1

ACOUSTIC IMPEDANCE APPARATUS ZA2

PNEUMOMETER ZP1


SKIN RESISTANCE APPARATUS RA4



SOUND LEVEL METER GA6
VISIBLE SPEECH APPARATUS
AUDITORY TRAINERS

Write for full catalogue and detailed descriptions to:

ALFRED PETERS & SONS LTD
51 GELL STREET, SHEFFIELD 3



What makes this antibiotic so valuable in the ENT unit?

Spectrum?

Exceptionally broad. It covers most gram-positive organisms—including the penicillin-resistant staphylococci—and many gram-negative organisms.

Action?

Intensely bactericidal. It gives a rapid kill of bacteria at or near the minimum inhibitory concentration, especially in the case of gram-negative organisms.

Safety?

Wide clinical usage has confirmed the laboratory evidence that Ceporin has a remarkably low toxicity. It is usually well tolerated by patients allergic to penicillin.

Clinical record?

The latest analysis of clinical results obtained with Ceporin shows a total of 131 ENT infections treated: 109 (83%) responded satisfactorily. There were 28 cases of tonsillitis, of which 24 responded well. The majority of these infections was caused by beta-haemolytic streptococci, but there were several mixed infections involving streptococci, staphylococci and Corynebacterium diphtheriae.

Otitis also responded well. Of the 28 cases treated, 24 responded satisfactorily. Most of these infections were caused by staphylococci but infections involving coliforms, pneumococci, Streptococcus pyogenes, proteus and Haemophilus influenzae were also represented. Often rapid clinical response was obtained with Ceporin after other antibiotics had failed. Conditions which respond well to Ceporin include: mastoiditis, otitis media, sinusitis, quinsy, pharyngeal abscess, and intracranial infections secondary to infected sinuses and mastoiditis. Because of its three main attributes broad spectrum, bactericidal action and low toxicity—Ceporin has proved valuable for use in place of more toxic drugs or combinations of antibiotics. **Presentation Vials containing** 250 mg, 500 mg and 1 gram of cephaloridine, packed singly and in boxes of five.

Ceporin is a Glaxo trade mark

Detailed literature is available from Glaxo Laboratories Limited Greenford, Middlesex