Comment

The needs of severely mentally ill offenders cannot be met in the hospital wings of prisons. Gunn et al (1991) surveyed 5% of prisoners in England and Wales and from their data suggested that 37% had psychiatric disorders. They also estimated that there could be as many as 700 sentenced prisoners who are psychotic.

Hospital wings of prisons are not recognised as hospitals; therefore powers under the Mental Health Act of forcibly treating psychotic prisoners under this legislation is clearly inappropriate. Many psychotic prisoners therefore find themselves confined in 'strip' cells.

Remand prisoners are under particular stress and the mentally ill are over-represented among prison suicides. Taylor & Gunn (1984) found that at least 9% of men remanded to Brixton prison were psychotic.

We were particularly concerned about the number of prisoners suffering from schizophrenia within Cardiff Prison. It has been estimated in a ten year follow-up of patients with schizophrenia in the community that 30% had some contact with the police and 12% were convicted of offences which were usually minor. Many of the offences took place because the patients were unable to care for themselves or had behaved inappropriately in a public place (Johnstone, 1991).

General psychiatrists need to be actively involved in the diversion of the mentally ill from the criminal justice system at all levels, from arrest by police to the extrication of the seriously mentally ill convicted prisoner. General psychiatrists ask where the extra resources are to come from to look after the diverted offender patients. The answer is that the majority of the schizophrenic inmates in our series were already known to general psychiatric services. Mentally disordered offenders should not be denied treatment merely by being offenders.

Acknowledgements

The authors would like to thank the Governor and staff of Cardiff Prison for their help in setting up this service.

References

ALLDERIDGE, P. (1990) Hospitals, madhouses and asylums: cycles in the care for the insane. In *Lectures on the History of Psychiatry* (Eds. Murray, R.M. & Turner, T.H.). London: Gaskell. (Royal College of Psychiatrists).

AMERICAN PSYCHIATRIC ASSOCIATION (1987) Diagnostic and Statistical Manual of Mental Disorders (3rd edition, revised) (DSM-III-R) Washington, DC: APA.

DEPARTMENT OF HEALTH & HOME OFFICE (1992) Review of Health and Social Services for Mentally Disordered Offenders and Others Requiring Similar Services: final summary report (Reed Committee) (CM 2088) London: HMSO.

GUNN, J., MADEN, A. & SWINTON, M. (1991) Treatment needs of prisoners with psychiatric disorders. *British Medical Journal*, **303**, 338-341.

JOHNSTONE, E.C. (1991) Disabilities and circumstances of schizophrenic patients: a follow-up study. British Journal of Psychiatry, 150, Suppl 1, 13.

TAYLOR, P.J. & GUNN, J. (1984) Violence and psychosis – risk of violence among psychotic men. British Medical Journal, 288, 194–198.

WELSH OFFICE (1992) Report of the All-Wales Advisory Group on Forensic Psychiatry. Cardiff: HMSO.

P.L. Huckle, Senior Registrar in Forensic Psychiatry; and Tegwyn Williams, Consultant Forensic Psychiatrist, The South Wales Forensic Psychiatric Service, The Caswell Clinic, Glanrhyd Hospital, Bridgend, Mid Glamorgan CF31 4LN

Addendum

Since the submission of this article, Cardiff Prison have changed their policy regarding the use of the unfurnished cell (the 'strip' cell). It is used only to manage actively suicidal prisoners, not to contain those who are psychotic.

672 Huckle & Williams