

## SOME BANACH SPACE EMBEDDINGS OF CLASSICAL FUNCTION SPACES

N.L. CAROTHERS AND S.J. DILWORTH

Banach space embeddings of the Orlicz space  $L_p + L_q$  and the Lorentz space  $L_{p,q}$  into the Lebesgue–Bochner space  $L_r(l_s)$  are demonstrated for appropriate ranges of the parameters.

### 1. INTRODUCTION

The purpose of this paper is to explain how the existence of isomorphic embeddings of particular classical Banach function spaces into the Lebesgue–Bochner spaces  $L_r(l_s)$  may be deduced quite easily from some known stochastic and analytic facts. Our approach has at its foundation a moment inequality of Rosenthal [9] for sums of independent random variables. Indeed, our Theorem 1 below, which describes an embedding of the Orlicz space  $L_r + L_s$  into  $L_r(l_s)$  in the range  $0 < r < s < \infty$ , is in effect a disguised form of a particular version (Theorem A below) of Rosenthal's inequality. Theorem A is a special case of a recent, very general result of Johnson and Schechtman [7], although the idea of using Rosenthal's inequality to define Banach space embeddings dates back at least to [6] (see also [8]). From Theorem 1 just a few short steps are required to deduce the existence of an embedding of the Lorentz space  $L_{p,q}$  into  $L_q(l_s)$  in the range  $q < p < s$ . The latter implies the surprising theorem of Schütt, obtained by rather different finite-dimensional methods, on the existence of an embedding of  $L_{p,q}$  into  $L_q$  for  $1 \leq q < p < 2$ . Schütt's theorem motivated our work to a great extent, and it is to be hoped that our approach helps to explain his theorem by exhibiting it in a different light.

### 2. NOTATION AND PRELIMINARY RESULTS

For  $0 < p < \infty$ ,  $L_p$  denotes the usual Lebesgue space of real-valued functions on  $[0, \infty)$  with its usual norm, denoted by  $\|\cdot\|_p$ . For  $0 < r < s < \infty$  and  $0 < t < \infty$ ,

---

Received 12 January 1990

The first author was supported in part by BGSU Faculty Research Committee Grant and the second author in part by NSF DMS-8801731.

---

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/91 \$A2.00+0.00.

$L_r + t \cdot L_s$  denotes the Orlicz space  $L_r + L_s$  equipped with the norm  $\|f\|_{L_r+tL_s} = K(t, f; L_r, L_s)$ , where

$$(1) \quad K(t, f; L_r, L_s) = \inf\{\|g\|_r + t\|h\|_s : f = g + h\}$$

is the standard  $K$ -functional of interpolation theory. The Lorentz space  $L_{p,q}$  is the space of real-valued functions on  $[0, \infty)$  with the norm  $\|f\|_{p,q} = [\int_0^\infty f^*(t)^q d(t^{q/p})]^{1/q}$ , where  $f^*$  denotes the decreasing rearrangement of  $|f|$ . This form of the  $L_{p,q}$  norm is not particularly suitable for our purposes, however, and we shall instead use the following formula which is derived from the Lions-Peetre  $K$ -method of interpolation (for example [1]):

$$(2) \quad \|f\|_{p,q} \sim \left[ \sum_{n=-\infty}^{+\infty} 2^{-n\theta q} K(2^n, f; L_r, L_s)^q \right]^{1/q}$$

provided  $1/p = (1 - \theta)/r + \theta/s$ . (Throughout,  $A \sim B$  means that there is a constant  $c > 0$ , depending only on the parameters  $p, q$ , et cetera, such that  $(1/c)A \leq B \leq cA$ .)

The Lebesgue–Bochner space  $L_r(l_s)$ ,  $0 < r, s < \infty$ , is the space of sequences  $(f_n)_{n=1}^\infty$  of functions on  $[0, 1]$  equipped with the norm

$$\|(f_n)\|_{L_r(l_s)} = \left[ \int_0^1 \left( \sum_{n=1}^\infty |f_n(t)|^s \right)^{r/s} dt \right]^{1/r}.$$

Given a family of Banach spaces  $(X_n)_{n=-\infty}^\infty$  and  $0 < p < \infty$ ,  $\left( \sum_{-\infty}^\infty \oplus X_n \right)_p$  denotes the space of sequences  $(x_n)_{n=-\infty}^\infty$ ,  $x_n \in X_n$ , with the norm  $\left( \sum_{-\infty}^\infty \|x_n\|^p \right)^{1/p}$ . When all the  $X_n$ 's are the same space  $X$ , we write  $l_p(X)$ .

We turn now to state the moment inequality of Rosenthal [9] in a form which illuminates its linear character. First some notation: given a sequence of functions  $(f_n)_{n=1}^\infty$  on  $[0, 1]$ , let  $\sum_n \oplus f_n$  denote the function  $f$  on  $[0, \infty)$  defined by  $f(n - 1 + t) = f_n(t)$  for  $n \geq 1$  and  $0 < t \leq 1$ . Identify  $[0, 1]$  with the (measure-equivalent) product space  $\Omega = [0, 1]^{\mathbb{N}}$ , and denote a typical element of  $\Omega$  by  $(s_1, s_2, \dots)$ , so that  $s_1, s_2, \dots$  are independent coordinates. The version of Rosenthal's inequality which we shall use may be stated succinctly in the following form.

**THEOREM A.** *Let  $0 < p < 2$ . Then*

$$(3) \quad \|(f_n(s_n))_{n=1}^\infty\|_{L_p(l_2)} \sim \left\| \sum_{n=1}^\infty \oplus f_n \right\|_{L_p+L_2}.$$

This version of Rosenthal's inequality was first obtained by Johnson and Schechtman [7]. (See also [2] or [4] for an approach more in keeping with the present paper.)

3. THE MAIN RESULTS

**THEOREM 1.** (See [4, Theorem 3.1]) *Let  $0 < r < s < \infty$ . Then  $L_r + L_s$  is linearly isomorphic to a subspace of  $L_r(l_s)$ .*

**PROOF:** Since  $2r/s < 2$ , we can apply (3) with  $f_n$  replaced by  $|f_n|^{s/2}$  to obtain

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \oplus f_n \right\|_{L_r + L_s} &\sim \left\| \sum_{n=1}^{\infty} \oplus |f_n|^{s/2} \right\|_{L_{2r/s} + L_2}^{2/s} \\ &\sim \left\| \left( \sum_{n=1}^{\infty} |f_n(s_n)|^s \right)^{1/2} \right\|_{2r/s}^{2/s} \\ &= \|(f_n(s_n))_{n=1}^{\infty}\|_{L_r(l_s)}. \end{aligned}$$

Hence the mapping  $\sum_n \oplus f_n \rightarrow (f_n(s_n))_{n=1}^{\infty}$  defines a linear isomorphism from  $L_r + L_s$  into  $L_r(l_s)$ . □

For our results on  $L_{p,q}$  the following simple proposition is the key.

**PROPOSITION 1.** *Let  $0 < r < s < \infty$  and let  $0 < t < \infty$ . Then  $L_r + t \cdot L_s$  is linearly isometric to  $L_r + L_s$ .*

**PROOF:** By dilation of (1) by  $\lambda > 0$ , we have

$$K(t, f; L_r, L_s) = \inf\{\lambda^{1/r} \|g(\lambda x)\|_r + t\lambda^{1/s} \|h(\lambda x)\|_s : f = g + h\}.$$

Select  $\lambda = t^{rs/(s-r)}$ , so that  $t = \lambda^{(s-r)/rs}$ . Then

$$\begin{aligned} K(t, f; L_r, L_s) &= \lambda^{1/r} \inf\{\|g(\lambda x)\|_r + \|h(\lambda x)\|_s : f = g + h\} \\ &= \lambda^{1/r} K(1, f(\lambda x); L_r, L_s) \\ &= t^{s/(s-r)} K\left(1, f\left(t^{rs/(s-r)} x\right); L_r, L_s\right). \end{aligned}$$

Thus the mapping  $f(x) \rightarrow t^{s/(s-r)} f(t^{rs/(s-r)} x)$  defines a linear isometry from  $L_r + t \cdot L_s$  onto  $L_r + L_s$ . □

**REMARK.** The above dilation argument rests on the fact that the function spaces are defined not on  $[0, 1]$  but on  $[0, \infty)$ .

**COROLLARY 1.** *Let  $0 < r < p < s < \infty$  and let  $0 < q < \infty$ . Then  $L_{p,q}$  is linearly isomorphic to a subspace of  $l_q(L_r + L_s)$ .*

**PROOF:** (2) says that  $L_{p,q}$  is isomorphic to a subspace of  $\left[ \sum_{-\infty}^{\infty} \oplus (L_r + 2^n \cdot L_s) \right]_q$ .

By Proposition 1 each  $L_r + t \cdot L_s$  is linearly isometric to  $L_r + L_s$ , and so the desired result follows. □

**THEOREM 2.** *Let  $0 < q < p < s$ . Then  $L_{p,q}$  is linearly isomorphic to a subspace of  $L_q(l_s)$ .*

**PROOF:** Applying Corollary 1 with  $r = q$  shows that  $L_{p,q}$  embeds into  $l_q(L_q + L_s)$ . Hence, by Theorem 1,  $L_{p,q}$  embeds into  $l_q(L_q(l_s))$  which is of course isomorphic to  $L_q(l_s)$ .  $\square$

**COROLLARY 2.** [10] *Let  $0 < q < p < 2$ . Then  $L_{p,q}$  is linearly isomorphic to a subspace of  $L_q$ .*

**PROOF:** Applying Theorem 2 with  $s = 2$  shows that  $L_{p,q}$  embeds into  $L_q(l_2)$ . It is a well-known consequence of Khintchine's inequality that the mapping  $(f_n(s))_{n=1}^{\infty} \rightarrow \sum_n f_n(s)r_n(t)$ , where  $(r_n(t))_{n=1}^{\infty}$  are the Rademacher functions, defines an isomorphic embedding of  $L_q(l_2)$  into  $L_q([0, 1]^2)$  (which is isomorphic to  $L_q$ ), and the desired conclusion follows.  $\square$

**REMARK.** It is also proved in [10] that if  $1 < p < q < 2$  then  $L_{p,q}$  is not isomorphic to a subspace of  $L_1$ . We indicate here a short proof of this fact. Since  $L_{p,q}$  has "type"  $p$  in this range (see for example [8] for the definition of this notion), it follows from the main result of [5] that if  $L_{p,q}$  embeds into  $L_1$ , then  $l_p$  embeds into  $L_{p,q}$ . But by [3, Theorem 2.8],  $l_p$  does not embed into  $L_{p,q}$ .

## REFERENCES

- [1] J. Bergh and J. Löfstrom, *Interpolation spaces* (Springer-Verlag, New York, 1976).
- [2] N.L. Carothers and S.J. Dilworth, 'Inequalities for sums of independent random variables', *Proc. Amer. Math. Soc.* **104** (1988), 221–226.
- [3] N.L. Carothers and S.J. Dilworth, 'Subspaces of  $L_{p,q}$ ', *Proc. Amer. Math. Soc.* **104** (1988), 537–545.
- [4] S.J. Dilworth, 'A scale of linear spaces related to the  $L_p$  scale', *Illinois J. Math.* (to appear).
- [5] S. Guerre and M. Levy, 'Espaces  $l_p$  dans les sous-espaces de  $L_1$ ', *Trans. Amer. Math. Soc.* **279** (1983), 611–616.
- [6] W.B. Johnson, B. Maurey, G. Schechtman and L. Tzafriri, 'Symmetric structures in Banach spaces', *Mem. Amer. Math. Soc.* **217** (1979).
- [7] W. B. Johnson and G. Schechtman, 'Sums of independent random variables in rearrangement invariant function spaces', *Ann. Probab.* **17** (1989), 789–808.
- [8] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II. Function Spaces* (Springer-Verlag, New York, 1979).
- [9] H.P. Rosenthal, 'On the subspaces of  $L_p$  ( $p > 2$ ) spanned by sequences of independent random variables', *Israel J. Math.* **8** (1970), 273–303.
- [10] C. Schütt, 'Lorentz spaces that are isomorphic to subspaces of  $L_1$ ', *Trans. Amer. Math. Soc.* **314** (1989), 583–595.

Department of Mathematics and Statistics  
Bowling Green State University  
Bowling Green OH 43403-0221  
United States of America

Department of Mathematics  
University of South Carolina  
Columbia SC 29208  
United States of America