

Corrigendum

Trajectories of change and long-term outcomes
in a randomised controlled trial of internet-
based insomnia treatment to prevent
depression – CORRIGENDUM

Philip J. Batterham, Helen Christensen, Andrew J. Mackinnon, John A. Gosling, Frances P. Thorndike,
Lee M. Ritterband, Nick Glozier and Kathleen M. Griffiths

Copyright and usage

© The Author(s), 2025. Published by Cambridge University Press on behalf of Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

DOI: <https://doi.org/10.1192/bjpo.bp.117.005231>, Published online by Cambridge University Press: 02 January 2018.

The authors regret the inclusion of an error in the supplementary materials of the above article. In [Table DS2](#), the column headings

under PSF suicidality were inadvertently switched. The corrected version of the table is available below.

The authors apologise for this error.

Reference

Batterham PJ, Christensen H, Mackinnon AJ, Gosling JA, Thorndike FP, Ritterband LM, et al. Trajectories of change and long-term outcomes in a randomised controlled trial of internet-based insomnia treatment to prevent depression. *BJPsych Open*. 2017;3(5):228–35. doi: [10.1192/bjpo.bp.117.005231](https://doi.org/10.1192/bjpo.bp.117.005231).

Table DS2 Estimated marginal means (s.e.) and observed means (s.d.) and by intervention condition over time

	PHQ-9 NS depression		GAD-7 anxiety		ISI insomnia		PSF suicidality	
	HealthWatch mean (s.e.)	SHUTi mean (s.e.)	HealthWatch mean (s.e.)	SHUTi mean (s.e.)	HealthWatch mean (s.e.)	SHUTi mean (s.e.)	HealthWatch mean (s.e.)	SHUTi mean (s.e.)
Estimated marginal mean (s.e.)								
Baseline	5.64 (0.15)	5.84 (0.15)	5.77 (0.18)	5.83 (0.18)	16.23 (0.18)	15.92 (0.18)	0.78 (0.07)	0.74 (0.07)
Intervention week 2	6.23 (0.16)	5.93 (0.19)	6.05 (0.19)	5.36 (0.22)	14.57 (0.19)	13.22 (0.22)		
Intervention week 4	5.61 (0.17)	4.95 (0.20)	5.59 (0.20)	4.55 (0.23)	13.53 (0.22)	10.15 (0.26)		
Intervention week 6	5.75 (0.18)	3.98 (0.21)	5.87 (0.20)	3.80 (0.24)	13.27 (0.23)	8.35 (0.28)		
Intervention week 8	5.53 (0.18)	3.22 (0.21)	5.70 (0.20)	3.18 (0.23)	12.91 (0.25)	7.24 (0.30)		
Post-test	4.57 (0.16)	2.87 (0.19)	5.01 (0.19)	3.21 (0.21)	13.17 (0.25)	7.35 (0.29)	0.81 (0.08)	0.51 (0.09)
6 months	4.02 (0.19)	2.92 (0.21)	4.23 (0.19)	3.15 (0.21)	12.17 (0.27)	7.74 (0.31)	0.72 (0.09)	0.55 (0.10)
12 months	3.74 (0.19)	2.89 (0.22)	4.08 (0.22)	3.00 (0.25)	11.45 (0.30)	7.70 (0.35)	0.69 (0.09)	0.42 (0.11)
18 months	4.44 (0.27)	2.55 (0.30)	4.30 (0.27)	3.07 (0.30)	11.52 (0.40)	8.02 (0.45)	1.04 (0.15)	0.70 (0.17)
	HealthWatch mean (s.d.)	SHUTi mean (s.d.)	HealthWatch mean (s.d.)	SHUTi mean (s.d.)	HealthWatch mean (s.d.)	SHUTi mean (s.d.)	HealthWatch mean (s.d.)	SHUTi mean (s.d.)
Observed mean (s.d.)								
Baseline	5.64 (3.59)	5.84 (3.82)	5.77 (4.33)	5.83 (4.36)	16.23 (4.32)	15.92 (4.18)	0.78 (1.69)	0.74 (1.71)
Intervention week 2	6.09 (3.47)	5.70 (3.76)	5.92 (4.15)	5.04 (4.22)	14.58 (4.25)	12.98 (3.90)		
Intervention week 4	5.53 (3.54)	4.51 (3.45)	5.55 (4.55)	3.88 (3.64)	13.63 (4.72)	9.64 (4.24)		
Intervention week 6	5.84 (3.98)	3.35 (3.32)	6.01 (4.71)	3.18 (3.46)	13.45 (4.92)	7.94 (4.13)		
Intervention week 8	5.45 (3.88)	2.62 (2.89)	5.61 (4.62)	2.69 (3.07)	13.22 (5.04)	6.49 (3.96)		
Post-test	4.50 (3.66)	2.49 (2.94)	4.83 (4.39)	2.69 (3.18)	13.21 (5.45)	6.88 (4.75)	0.78 (1.80)	0.51 (1.53)
6 months	3.85 (3.63)	2.57 (3.32)	3.99 (3.77)	2.71 (3.44)	11.84 (5.13)	7.26 (4.87)	0.72 (1.71)	0.55 (1.77)
12 months	3.62 (3.46)	2.31 (2.64)	3.98 (4.15)	2.45 (3.31)	11.18 (5.54)	6.83 (4.61)	0.68 (1.80)	0.28 (0.98)
18 months	4.52 (4.30)	2.01 (2.37)	4.43 (4.20)	2.53 (2.95)	11.19 (5.78)	7.70 (5.05)	1.09 (2.41)	0.45 (1.49)

PHQ-9NS: Patient Health Questionnaire-9 without sleep item; GAD-7: Generalized Anxiety Disorder-7; ISI: Insomnia Severity Index; PSF: Psychiatric Symptom Frequency scale.