

THE CATENARY AND TAME DEGREES ON A NUMERICAL MONOID ARE EVENTUALLY PERIODIC

SCOTT T. CHAPMAN[✉], MARLY CORRALES, ANDREW MILLER,
CHRIS MILLER and DHIR PATEL

(Received 11 September 2013; accepted 8 July 2014; first published online 9 September 2014)

Communicated by M. Jackson

Abstract

Let M be a commutative cancellative monoid. For m a nonunit in M , the catenary degree of m , denoted $c(m)$, and the tame degree of m , denoted $t(m)$, are combinatorial constants that describe the relationships between differing irreducible factorizations of m . These constants have been studied carefully in the literature for various kinds of monoids, including Krull monoids and numerical monoids. In this paper, we show for a given numerical monoid S that the sequences $\{c(s)\}_{s \in S}$ and $\{t(s)\}_{s \in S}$ are both eventually periodic. We show similar behavior for several functions related to the catenary degree which have recently appeared in the literature. These results nicely complement the known result that the sequence $\{\Delta(s)\}_{s \in S}$ of delta sets of S also satisfies a similar periodicity condition.

2010 *Mathematics subject classification*: primary 20M13; secondary 20M14, 11D05.

Keywords and phrases: catenary degree, tame degree, numerical monoid.

1. Introduction

Over the past 20 years, problems involving nonunique factorizations of elements in integral domains and commutative cancellative monoids have been widely popular in the mathematical literature (see [15] and its citation list). Much of this literature focuses on various combinatorial constants which describe in some sense how far these systems vary from the classical notion of unique factorization. While early work in this area focused on Krull domains and monoids (see [3, 4, 11, 13, 14, 16, 19]), many papers have recently considered these properties on numerical monoids (which are additive submonoids of the natural numbers). In particular, their elastic properties (see [8]), their delta sets (see [2, 5, 9]) and their catenary and tame degrees (see [1, 4, 6, 7, 12, 17, 18]) have been examined in detail. We take particular interest in the main result of [9], where, for a given numerical monoid S , the authors show that the sequence of delta sets $\{\Delta(s)\}_{s \in S}$ is eventually periodic. In this note, we prove an

The authors were supported by National Science Foundation grants DMS-1035147 and DMS-1045082.
© 2014 Australian Mathematical Publishing Association Inc. 1446-7887/2014 \$16.00

analogue of this result by showing that the similar sequences defined by the catenary degree, the tame degree and the various related forms of the catenary degree recently introduced in the literature (see [16]) are also eventually periodic. Our argument differs from the one offered in [9], as problems involving the catenary and tame degrees rely on the complete set of factorizations of an element, while those involving the delta sets are merely concerned with factorizations of differing lengths. We open in Section 2 with the necessary notation and definitions, and present our main result, with proofs, in Section 3.

2. Definitions and preliminaries

A numerical monoid S is a cofinite additive submonoid of $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. Both [20] and [21] are good general references on the subject. It is easy to show using elementary number theory that every numerical monoid has a unique minimal generating set. If these generators are n_1, \dots, n_k with $n_1 < n_2 < \dots < n_k$, then we use the notation

$$S = \langle n_1, \dots, n_k \rangle = \{a_1n_1 + \dots + a_kn_k \mid a_1, \dots, a_k \in \mathbb{N}_0\}.$$

If $\gcd(n_1, \dots, n_k) \neq 1$, then $\mathbb{N}_0 \setminus S$ is not finite, so we must have $\gcd(n_1, \dots, n_k) = 1$. We call k the *embedding dimension* of s . Since $\mathbb{N}_0 \setminus S$ is finite, there is a largest number in the complement of S , denoted $\mathcal{F}(S)$, and called the *Frobenius number* of S .

Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid. For $s \in S$, let

$$Z(s) = \{(a_1, \dots, a_k) \mid a_1n_1 + \dots + a_kn_k = s \text{ with each } a_i \in \mathbb{N}_0\}$$

be the *set of factorizations* of s in S . We say that the *length* of $z = (a_1, \dots, a_k) \in Z(s)$ is

$$|z| = a_1 + \dots + a_k.$$

Set

$$\mathcal{L}(s) = \{|z| : z \in Z(s)\} = \{m_1, \dots, m_l\},$$

where we assume that $m_1 < m_2 < \dots < m_{l-1} < m_l$. The set $\mathcal{L}(s)$ is known as the *set of lengths* of s . The *delta set* of an element, denoted $\Delta(s)$, is the set containing the values of the difference of consecutive elements of $\mathcal{L}(s)$, that is,

$$\Delta(s) = \{m_{i+1} - m_i \mid 1 \leq i < l\}.$$

Let $z = (a_1, \dots, a_k)$ and $z' = (b_1, \dots, b_k) \in Z(s)$. We say that the greatest common divisor of z and z' is

$$\gcd(z, z') = (\min\{a_1, b_1\}, \dots, \min\{a_k, b_k\}),$$

and we define the *distance* between z and z' as

$$d(z, z') = \max\{|z - \gcd(z, z')|, |z' - \gcd(z, z')|\}.$$

The distance function satisfies many of the usual properties of a metric; the interested reader can find these summarized in [15, Proposition 1.2.5]. If $Z' \subseteq Z(s)$, then set

$$d(z, Z') = \min\{d(z, z') \mid z' \in Z'\}.$$

A sequence

$$z = z_0, z_1, \dots, z_{n-1}, z_n = z'$$

of factorizations in $Z(s)$ is an N -chain if $d(z_i, z_{i+1}) \leq N$ for each $1 \leq i \leq n - 1$. For $s \in S$, we define the *catenary degree* of s (denoted $c(s)$) to be the minimal N such that there is an N -chain between any two factorizations of s .

The *tame degree* of an element $t(s)$ is constructed as follows. For each $i \leq k$, let $Z^i(s) := \{(a_1, \dots, a_k) \in Z(s) \mid a_i \neq 0\}$. We further let

$$t_i(s) = \max_{z \in Z(s)} d(z, Z^i(s)) \quad \text{and} \quad t(s) = \max_{i \leq k} t_i(s).$$

Alternatively, we can say that $t(s)$ is the minimal number such that $d(z, Z^i(s)) \leq t(s)$ for all $z \in Z(s)$ and all $i \leq k$.

Three variations on the catenary degree have appeared in the literature (most recently in [16]; see also [19]). Their definitions are as follows.

- (1) The *monotone catenary degree* of an element $c_{\text{mon}}(s)$ is the minimal number such that for any $z, z' \in Z(s)$ with $|z| \leq |z'|$, there exists a $c_{\text{mon}}(s)$ -chain $z = z_1, z_2, \dots, z_k = z'$ with the added restriction that $|z_i| \leq |z_{i+1}|$.
- (2) The *equivalent catenary degree* $c_{\text{eq}}(s)$ of an element $s \in S$ is the minimal number such that given $z, z' \in Z(s)$ with $|z| = |z'|$, there exists a $c_{\text{eq}}(s)$ -chain $z = z_1, \dots, z_k = z'$ with the added restriction that $|z_i| = |z_{i+1}|$.
- (3) We say that $a, b \in \mathcal{L}(s)$ (with $a < b$) are *adjacent* if $[a, b] \cap \mathcal{L}(s) = \{a, b\}$. Let $Z_l(s) = \{z \in Z(s) \mid |z| = l\}$. The *adjacent catenary degree* $c_{\text{adj}}(s)$ of an element $s \in S$ is the minimal number such that $d(Z_a(s), Z_b(s)) \leq c_{\text{adj}}(s)$ for all adjacent a, b .

We close this section by noting that computing done in connection with these results was run on the GAP numerical semigroups package [10]. Also, any undefined notation or definitions will be consistent with those used in the monograph [15].

3. Periodicity

Given a numerical monoid $S = \langle n_1, \dots, n_k \rangle$, we define $L(S) = \text{lcm}\{n_1, \dots, n_k\}$. When there is no ambiguity, we shall simply write L . The remainder of this section will consist of a proof of our main result, which is as follows.

THEOREM 3.1. *If $S = \langle n_1, \dots, n_k \rangle$ is a numerical monoid, then the sequences*

$$\{c(s)\}_{s \in S}, \quad \{t(s)\}_{s \in S}, \quad \{c_{\text{mon}}(s)\}_{s \in S}, \quad \{c_{\text{eq}}(s)\}_{s \in S}, \quad \text{and} \quad \{c_{\text{adj}}(s)\}_{s \in S}$$

are all eventually periodic with fundamental period a divisor of L .

Let $S = \langle n_1, \dots, n_k \rangle$ and suppose that $k = 2$. Using techniques from [6], one can readily verify that $t(s) = c(s) = n_2$ for large s (see also [15, Example 3.1.6]). Moreover, it also follows for large s that $c_{\text{adj}}(s) = c(s)$. Since for $k = 2$ we also have for all s that $c_{\text{eq}}(s) = 0$ and $c_{\text{mon}}(s) = \max\{c_{\text{eq}}(s), c_{\text{adj}}(s)\} = c_{\text{adj}}(s) = c(s)$ (see [16, page 1003]), we can assume throughout the remainder of our paper that $k \geq 3$.

The proof of Theorem 3.1 will rely on the following basic sequencing lemma, whose proof is left to the reader.

LEMMA 3.2. *Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid and $f : S \rightarrow \mathbb{N}_0$ a function. If there exist positive integers N and M such that $s \in S$ and $s > M$ imply that $f(s - N) \geq f(s)$, then $\{f(s)\}_{s \in S}$ is eventually periodic with fundamental period a divisor of N .*

The following definition is critical to all of our remaining proofs.

DEFINITION 3.3. Let s be an element of a numerical monoid $S = \langle n_1, \dots, n_k \rangle$ such that $s - L \in S$. For each i , with $1 \leq i \leq k$, define a map

$$\phi_i : Z(s - L) \rightarrow Z(s)$$

by

$$\phi_i : z \rightarrow z + \left(0, \dots, 0, \frac{L}{n_i}, 0, \dots, 0\right).$$

For each i , it is easy to verify that ϕ_i is *distance preserving* (that is, $d(z, z') = d(\phi_i(z), \phi_i(z'))$ for all $z, z' \in Z(s - L)$). In the next proposition, we describe the set $Z(s)$ in terms of the images under ϕ_i of $s - L$.

PROPOSITION 3.4. *If $S = \langle n_1, \dots, n_k \rangle$ and $s \in S$ are as in Definition 3.3 with $s \geq L(kn_k)$, then*

$$Z(s) = \bigcup_{i \leq k} \phi_i(Z(s - L)).$$

PROOF. Let $(a_1, \dots, a_k) \in Z(s)$. Then $\sum_{i=1}^k a_i n_i = s$. Observe that

$$kn_k \cdot \max_{i \leq k} a_i \geq \sum_{i=1}^k a_i n_i = s \geq L(kn_k).$$

If we denote $a_j = \max_{i \leq k} a_i$ and simplify, then $a_j \geq L > L/n_j$. So, we write $(a_1, \dots, a_k) = (a_1, \dots, a_j - (L/n_j), a_{j+1}, \dots, a_k) + (0, \dots, 0, L/n_j, 0, \dots, 0)$. Hence, $(a_1, \dots, a_k) = \phi_j(a_1, \dots, a_j - (L/n_j), a_{j+1}, \dots, a_k)$ and we conclude that $Z(s) = \bigcup_{i \leq k} \phi_i(Z(s - L))$, where the reverse inclusion is obvious. \square

Proposition 3.4 leads to the following observations concerning the catenary and tame degrees of relatively large elements of a numerical monoid.

THEOREM 3.5. *Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid.*

- (a) *If $s \in S$ with $s \geq \max\{L(kn_k), \mathcal{F}(S) + 2L + 1\}$, then $c(s - L) \geq c(s)$.*
- (b) *If $s \in S$ with $s \geq \max\{L(kn_k), \mathcal{F}(S) + L + 1 + n_k\}$, then $t(s - L) \geq t(s)$.*

PROOF. (a) Since $s \geq L(kn_k)$, we have $Z(s) = \bigcup_{i \leq k} \phi_i(Z(s - L))$ by Proposition 3.4. Let $j < l \leq k$. Then write $s = (s - 2L) + (L/n_j)n_j + (L/n_l)n_l$. By hypothesis, we have $s - 2L \geq \mathcal{F}(S) + 1$. So, $Z(s - 2L)$ is nonempty. Pick $(a_1, \dots, a_k) \in Z(s - 2L)$. We then observe that

$$\left(a_1, \dots, a_j + \frac{L}{n_j}, a_{j+1}, \dots, a_k\right) + \left(0, \dots, \frac{L}{n_l}, 0, \dots, 0\right) \in \phi_l(Z(s - L))$$

and

$$\left(a_1, \dots, a_l + \frac{L}{n_l}, a_{l+1}, \dots, a_k\right) + \left(0, \dots, \frac{L}{n_j}, 0, \dots, 0\right) \in \phi_j(Z(s - L))$$

represent the same factorization. Since j, l were arbitrary, we know that the images $\phi_p(Z(s - L))$ have pairwise nontrivial intersection.

The ϕ_p are all distance-preserving maps, so they conserve catenary degree locally within their image. Pick $x, y \in Z(s)$. Then we have $x \in \phi_j(Z(s - L))$ for some $j \leq k$, and $y \in \phi_l(Z(s - L))$ for some $l \leq k$. Now we have two cases.

Case 1. $l = j$. There is nothing to do; there exists a path with sufficiently small catenary degree within $\phi_l(Z(s - L))$, by distance preservation.

Case 2. $l \neq j$. Then there exists some element $z \in \phi_l(Z(s - L)) \cap \phi_j(Z(s - L))$. We can move from x to z within $\phi_j(Z(s - L))$, and then from z to y within $\phi_l(Z(s - L))$. Each time we have sufficiently small catenary degree.

Thus, we have produced a $c(s - L)$ -chain connecting x and y . We conclude that $c(s - L) \geq c(s)$.

(b) Since $s \geq L(kn_k)$, we again have $Z(s) = \bigcup_{i \leq k} \phi_i(Z(s - L))$ by Proposition 3.4. Pick $z \in Z(s)$. By Proposition 3.4, we have $z = \phi_i(z') \in Z(s)$ for some $z' \in Z(s - L)$. For an arbitrary j , let $z'_j \in Z^j(s - L)$ be such that $d(z', z'_j) = d(z', Z^j(s - L))$. Let $z_j = \phi_i(z'_j) \in Z^j(s)$. Observe that

$$s - L - n_j \geq \mathcal{F}(S) + L + 1 + n_k - L - n_j \geq \mathcal{F}(S) + 1 + n_k - n_j \geq \mathcal{F}(S) + 1.$$

So, we have $Z^j(s - L) \neq \emptyset$, and thus z'_j exists. We have

$$d(z, Z^j(s)) \leq d(z, z_j) = d(\phi_i(z'), \phi_i(z'_j)) = d(z', z'_j) \leq t(s - L).$$

Since z and j were arbitrary, let

$$d(z, Z^j(s)) = \max_{i \leq k} \max_{z' \in Z(s)} d(z'', Z^i(s)) = t(s).$$

It follows that $t(s - L) \geq t(s)$. □

To approach periodicity for the related versions of the catenary degree, we will need some further results. Given a factorization (a_1, \dots, a_k) of x with length a and large values for all a_i , we will produce in Lemma 3.7 a new factorization of x with

length a . To begin with this process, pick some i, j, k satisfying $1 \leq i < j < l \leq k$ and observe that

$$(\dots a_j - n_l, \dots, a_l + n_j, \dots) \tag{3.1}$$

is a factorization of x with length $a + (n_j - n_l) = a - (n_l - n_j)$ and

$$(\dots a_i + n_j, \dots, a_j - n_i, \dots) \tag{3.2}$$

is a factorization of x with length $a + (n_j - n_i)$. If we apply the exchange (3.1) $n_j - n_i$ times, and exchange (3.2) $n_l - n_j$ times, then we will produce a new factorization with length a . But, we need a_j to be sufficiently large. For this reason, we need an additional definition.

DEFINITION 3.6. Let $S = \langle n_1, \dots, n_k \rangle$ and assume that $k \geq 3$ throughout. Define

$$\omega(S) := \frac{L}{n_1} + \left\lceil \frac{L}{n_1 n_2} \right\rceil n_{k-1} (n_k - n_1).$$

When there is no ambiguity, this value will simply be denoted by ω and we call ω the *toppling number* of S .

Given Definition 3.6, we proceed with the previously promised lemma.

LEMMA 3.7 (The toppling lemma). Let S be as in Definition 3.6 and suppose that $s \in S$. Let $z = (a_1, \dots, a_k) \in Z(s)$ with $a_j \geq \omega$ for some $j \neq 1, k$. For any $1 \leq i, l \leq k$ with $i, l \neq j$, there exists $z' \in Z(s)$, of the form

$$\left(\dots, a_i + \left\lceil \frac{L}{n_1 n_2} \right\rceil [(n_l - n_j)n_j], \dots, a_j - \left\lceil \frac{L}{n_1 n_2} \right\rceil [n_j n_l - n_j n_i], \dots, a_l + \left\lceil \frac{L}{n_1 n_2} \right\rceil [(n_j - n_i)n_j], \dots \right),$$

and $|z| = |z'|$.

We refer to the process of changing z into z' in Lemma 3.7 as *toppling a_j to a_i and a_l* .

PROOF. Observe that

$$\begin{aligned} |z'| &= |z| + \left\lceil \frac{L}{n_1 n_2} \right\rceil ((n_l - n_j)n_j - (n_j n_l - n_j n_i) + (n_j - n_i)n_j) \\ &= |z| + \left\lceil \frac{L}{n_1 n_2} \right\rceil (n_i n_j - n_j^2 - n_j n_l + n_j n_i + n_j^2 - n_i n_j) = |z| + \left\lceil \frac{L}{n_1 n_2} \right\rceil (0) = |z| \end{aligned}$$

and hence $|z| = |z'|$. Also,

$$\begin{aligned} \sum_{i=1}^k a'_i n_i &= \sum_{i=1}^k a_i n_i + \left\lceil \frac{L}{n_1 n_2} \right\rceil ((n_l - n_j)n_j n_i - (n_j n_l - n_j n_i)n_j + (n_j - n_i)n_j n_l) \\ &= \sum_{i=1}^k a_i n_i + \left\lceil \frac{L}{n_1 n_2} \right\rceil (n_l n_j n_i - n_j^2 n_i - n_j^2 n_l + n_j^2 n_i + n_j^2 n_l - n_i n_j n_l) \\ &= \sum_{i=1}^k a_i n_i + \left\lceil \frac{L}{n_1 n_2} \right\rceil (0) = \sum_{i=1}^k a_i n_i. \end{aligned}$$

So, z and z' are factorizations of the same element. Furthermore,

$$\begin{aligned} a_j - \left\lfloor \frac{L}{n_1 n_2} \right\rfloor [(n_j - n_i)n_l + (n_l - n_j)n_i] &\geq \omega - \left\lfloor \frac{L}{n_1 n_2} \right\rfloor n_j(n_l - n_i) \\ &\geq \omega - \left\lfloor \frac{L}{n_1 n_2} \right\rfloor n_{k-1}(n_k - n_1) \\ &= \frac{L}{n_1} + \left\lfloor \frac{L}{n_1 n_2} \right\rfloor n_{k-1}(n_k - n_1) - \left\lfloor \frac{L}{n_1 n_2} \right\rfloor n_{k-1}(n_k - n_1) = \frac{L}{n_1} > \frac{L}{n_j}. \end{aligned}$$

So, all of the coefficients are positive (that is, $z' \in Z(s)$). This completes the proof. \square

Note that z' as constructed in Lemma 3.7 is in the image of three maps. First, $z' \in \phi_j(Z(s - L))$ by the last calculation in the above proof. Moreover,

$$a_i + \left\lfloor \frac{L}{n_1 n_2} \right\rfloor [(n_l - n_j)n_j] \geq \left\lfloor \frac{L}{n_1 n_2} \right\rfloor [(1)n_2] \geq \frac{L}{n_1} \geq \frac{L}{n_i}$$

and so $z' \in \phi_i(Z(s - L))$. Similarly,

$$a_i + \left\lfloor \frac{L}{n_1 n_2} \right\rfloor [(n_j - n_i)n_j] \geq \left\lfloor \frac{L}{n_1 n_2} \right\rfloor [(1)n_2] \geq \frac{L}{n_1} \geq \frac{L}{n_i}$$

and so $z' \in \phi_l(Z(s - L))$. Thus,

$$z' \in \phi_j(Z(s - L)) \cap \phi_i(Z(s - L)) \cap \phi_l(Z(s - L)).$$

Lemma 3.7 now allows us to prove an analogue of Theorem 3.5 for the sequence $\{c_{\text{eq}}(s)\}_{s \in S}$.

THEOREM 3.8. *Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid and suppose that $s \in S$. If*

$$N = \left\lfloor \frac{(k - 1)\omega}{1 - (n_1/n_k)} \right\rfloor$$

and $s \geq Nkn_k$, then $c_{\text{eq}}(s - L) \geq c_{\text{eq}}(s)$.

PROOF. Pick any two factorizations $z = (a_1, \dots, a_k)$ and $z' = (b_1, \dots, b_k)$ of s of the same length. Observe for $a_j = \max_{i \leq k} a_i$ that

$$ka_j n_k \geq \sum_{i=1}^k a_i n_i = s \geq kNn_k$$

and so $a_j \geq N$. Hence, it is clear using the definitions that $a_j \geq N \geq \omega \geq L/n_j$. The same can be said for $b_l = \max_{i \leq k} b_i \geq L/n_l$.

Case 1: $j = l$. Then z and z' are factorizations in $\phi_{j=l}(Z(s - L))$, so there exists a $c_{\text{eq}}(s - L)$ -chain connecting them in $\phi_{j=l}(Z(s - L))$. Thus, $c_{\text{eq}}(s) \leq c_{\text{eq}}(s - L)$.

Case 2: $j \neq 1, k$ and $l \neq j$. Note that this case is symmetric to the case $l \neq 1, k$ and $j \neq l$. Observe that $z \in \phi_j(Z(s - L))$ and $z' \in \phi_l(Z(s - L))$.

Case 2a: $l < j$. Topple z to some z'' by toppling a_j to a_l and a_k .

Case 2b: $l > j$. Topple z to some z'' by toppling a_j to a_1 and a_l .

Note that $z'' \in \phi_l(Z(s - L)) \cap \phi_j(Z(s - L))$. We can construct a $c_{\text{eq}}(s - L)$ -chain from z to z'' in $\phi_j(Z(s - L))$, and then from z'' to z' in $\phi_l(Z(s - L))$. Combining these chains, we have a $c_{\text{eq}}(s - L)$ chain from z to z' , so $c_{\text{eq}}(s) \leq c_{\text{eq}}(s - L)$.

Case 3: $a_1 \geq N$ and $b_k \geq N$ or $b_1 \geq N$ and $a_k \geq N$. Without loss of generality, let $a_1 \geq N$ and $b_k \geq N$. We have

$$\begin{aligned} \sum_{i=1}^k a_i = \sum_{i=1}^k b_i &\implies \sum_{i=1}^k a_i - b_k = \sum_{i=1}^{k-1} b_i \\ &\implies \sum_{i=1}^k a_i - b_k \leq (k - 1)b_m \implies \frac{\sum_{i=1}^k a_i - b_k}{k - 1} \leq b_m, \end{aligned}$$

where $b_m = \max_{i \neq k} b_i$. We also have

$$\begin{aligned} \sum_{i=1}^k a_i n_i = \sum_{i=1}^k b_i n_i &\implies b_k = \frac{\sum_{i=1}^k a_i n_i - \sum_{i=1}^{k-1} b_i n_i}{n_k} \\ &\implies b_k = \sum_{i=1}^k a_i \frac{n_i}{n_k} - \sum_{i=1}^{k-1} b_i \frac{n_i}{n_k}. \end{aligned}$$

Combining these results,

$$\begin{aligned} b_m &\geq \frac{\sum_{i=1}^k a_i - (\sum_{i=1}^k a_i (n_i/n_k) - \sum_{i=1}^{k-1} b_i (n_i/n_k))}{k - 1} \\ &\geq \frac{\sum_{i=1}^k a_i (1 - (n_i/n_k)) + \sum_{i=1}^{k-1} b_i (n_i/n_k)}{k - 1} \\ &\geq \frac{\sum_{i=2}^{k-1} a_i (1 - (n_i/n_k)) + b_i (n_i/n_k)}{k - 1} + \frac{a_1 (1 - (n_1/n_k)) + b_1 (n_1/n_k)}{k - 1} \\ &\geq \frac{a_1 (1 - (n_1/n_k))}{k - 1} \geq \frac{N(1 - (n_1/n_k))}{k - 1} \geq \omega \end{aligned}$$

for some $1 \leq m < k$. If $m = 1$, then both a_1 and $b_1 > \omega$, and we are in Case 1. If $m \neq 1$, then $b_m > \omega$ for some $m \neq 1, k$, and we are in Case 2. So, regardless of which pair of equal-length factorizations we choose, we can construct a $c_{\text{eq}}(s)$ -chain connecting them. We conclude that $c_{\text{eq}}(s - L) \geq c_{\text{eq}}(s)$, which completes the proof. \square

We prove a version of Theorem 3.8 for the sequence $\{c_{\text{adj}}(s)\}_{s \in S}$.

THEOREM 3.9. *Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid and suppose that $s \in S$. Suppose further that*

$$N = \left\lceil \frac{(k - 1)\omega + \Delta_{\text{max}}}{1 - (n_1/n_k)} \right\rceil,$$

where $\Delta_{\text{max}} = \max \Delta(S)$. If $s \geq kN$, then $c_{\text{adj}}(s - L) \geq c_{\text{adj}}(s)$.

PROOF. We note that, by [5], Δ_{\max} is finite. Pick $a, b \in \mathcal{L}(s)$ that are adjacent. Then $a = b + \Delta$ for some $\Delta \in \Delta(s)$. It is sufficient to show that there exist $x, y \in Z(s)$ such that $|x| = a$, $|y| = b$ and $x, y \in \phi_i(Z(s - L))$ for some $i \leq k$. For this would imply that $Z_{a-(L/n_i)}(s - L)$ and $Z_{b-(L/n_i)}(s - L)$ are nonempty, so we can pick $p \in Z_{a-(L/n_i)}(s - L)$ and $q \in Z_{b-(L/n_i)}(s - L)$ such that $d(p, q) = d(Z_{a-(L/n_i)}(s - L), Z_{b-(L/n_i)}(s - L))$. Since $a - (L/n_i)$ and $b - (L/n_i)$ are adjacent, we get $d(Z_{a-(L/n_i)}(s - L), Z_{b-(L/n_i)}(s - L)) \leq c_{\text{adj}}(s - L)$. Hence,

$$\begin{aligned} d(Z_a(s), Z_b(s)) &\leq d(\phi_i(p), \phi_i(q)) = d(p, q) \\ &= d(Z_{a-(L/n_i)}(s - L), Z_{b-(L/n_i)}(s - L)) \leq c_{\text{adj}}(s - L). \end{aligned}$$

So, our goal is to show that there exist $x, y \in \phi_i(Z(s))$ with $x \in Z_a(s)$ and $y \in Z_b(s)$.

Pick $z_a = (a_1, \dots, a_k) \in Z_a(s)$ and $z_b = (b_1, \dots, b_k) \in Z_b(s)$. As before, $a_i \geq N$ and $b_j \geq N$ for some $i, j \leq k$. We break our argument into five cases.

Case 1: $i = j$. Then $z_a \in \phi_i(Z(s - L)) \cap Z_a(s)$ and $z_b \in \phi_i(Z(s - L)) \cap Z_b(s)$. This completes the argument for Case 1.

Case 2: $i \neq 1, k$. This case breaks into two subcases.

Case 2a: $j > i$. Topple a_i to produce a factorization (a'_1, \dots, a'_k) , where $a'_1 \geq L/n_1$, $a'_i \geq L/n_i$ and $a'_j \geq L/n_j$.

Case 2b: $j < i$. Topple a_i to produce a factorization (a'_1, \dots, a'_k) , where $a'_j \geq L/n_j$, $a'_i \geq L/n_i$ and $a'_k \geq L/n_k$.

We have $(a'_1, \dots, a'_k) \in \phi_j(Z(s - L)) \cap Z_a(s)$. Since $z_b \in \phi_j(Z(s - L)) \cap Z_b(s)$, we are done with Case 2.

Case 3: $b_j \neq b_1, b_k$. This case also breaks into two subcases.

Case 3a: $i > j$. Topple b_j to produce a factorization (b'_1, \dots, b'_k) , where $b'_1 \geq L/n_1$, $b'_j \geq L/n_j$ and $b'_i \geq L/n_i$.

Case 3b: $i < j$. Topple b_j to produce a factorization (b'_1, \dots, b'_k) , where $b'_i \geq (L/n_i)$, $b'_j \geq L/n_j$ and $b'_k \geq L/n_k$.

We have $(b'_1, \dots, b'_k) \in \phi_i(Z(s - L)) \cap Z_b(s)$. We also know that $(a_1, \dots, a_k) \in \phi_i(Z(s - L)) \cap Z_a(s)$ by hypothesis. This completes Case 3.

Case 4: $i = k$ and $j = 1$. Set $a_m = \max_{i \neq k} a_i$. If $m = 1$, then (a_1, \dots, a_k) and (b_1, \dots, b_k) are both in the image of ϕ_1 and we are done. Thus, we assume that $m > 1$. We have

$$\begin{aligned} \sum_{i=1}^k a_i = \sum_{i=1}^k b_i + \Delta &\implies a_m(k - 1) \geq \sum_{i=1}^{k-1} a_i = \sum_{i=1}^k b_i - a_k + \Delta \\ &\implies a_m \geq \frac{\sum_{i=1}^k b_i - a_k + \Delta}{k - 1}. \end{aligned}$$

We also get

$$\sum_{i=1}^k a_i n_i = \sum_{i=1}^k b_i n_i \quad \text{and} \quad a_k = \frac{\sum_{i=1}^k b_i n_i - \sum_{i=1}^{k-1} a_i n_i}{n_k} = \sum_{i=1}^k b_i \frac{n_i}{n_k} - \sum_{i=1}^{k-1} a_i \frac{n_i}{n_k}.$$

Combining the above two results, we get

$$\begin{aligned} a_m &\geq \frac{\sum_{i=1}^k b_i - (\sum_{i=1}^k b_i(n_i/n_k) - \sum_{i=1}^{k-1} a_i(n_i/n_k)) + \Delta}{k-1} \\ \implies a_m &\geq \frac{\sum_{i=1}^k b_i(1 - (n_i/n_k)) + \sum_{i=1}^{k-1} a_i(n_i/n_k) + \Delta}{k-1} \\ &= \frac{\sum_{i=2}^{k-1} b_i(1 - (n_i/n_k)) + a_i(n_i/n_k)}{k-1} + \frac{b_1(1 - (n_1/n_k)) + a_1(n_1/n_k) + \Delta}{k-1} \\ &\geq \frac{b_1(1 - (n_1/n_k)) + \Delta}{k-1} \\ &\geq \frac{N(1 - (n_1/n_k)) + \Delta}{k-1} = \frac{\omega(k-1) + \Delta_{\max} + \Delta}{k-1} \geq \omega. \end{aligned}$$

Topple a_m to produce a factorization (a'_1, \dots, a'_k) , with $a'_1 \geq L/n_1$, $a'_m \geq L/n_j$ and $a'_k \geq L/n_k$. We have $(a'_1, \dots, a'_k) \in \phi_1(Z(s-L)) \cap Z_a(s)$. We also have $(b_1, \dots, b_k) \in \phi_1(Z(s-L)) \cap Z_b(s)$ by hypothesis. This completes Case 4.

Case 5: $i = 1$ and $j = k$. We have

$$\sum_{i=1}^k a_i = \sum_{i=1}^k b_i + \Delta \implies \sum_{i=1}^k a_i - b_k - \Delta = \sum_{i=1}^{k-1} b_i \leq (k-1)b_m \implies b_m \geq \frac{\sum_{i=1}^k a_i - b_k - \Delta}{k-1},$$

where $b_m = \max_{i \neq k} b_i$. We also get

$$\sum_{i=1}^k a_i n_i = \sum_{i=1}^k b_i n_i \implies b_k = \frac{\sum_{i=1}^k a_i n_i - \sum_{i=1}^{k-1} b_i n_i}{n_k} = \sum_{i=1}^k a_i \frac{n_i}{n_k} - \sum_{i=1}^{k-1} b_i \frac{n_i}{n_k}.$$

Combining the above two results, we get

$$\begin{aligned} b_m &\geq \frac{\sum_{i=1}^k a_i - (\sum_{i=1}^k a_i(n_i/n_k) - \sum_{i=1}^{k-1} b_i(n_i/n_k)) - \Delta}{k-1} \\ \implies b_m &\geq \frac{\sum_{i=1}^k a_i(1 - (n_i/n_k)) + \sum_{i=1}^{k-1} b_i(n_i/n_k) - \Delta}{k-1} \\ &= \frac{\sum_{i=2}^{k-1} a_i(1 - (n_i/n_k)) + b_i(n_i/n_k)}{k-1} + \frac{a_1(1 - (n_1/n_k)) + b_1(n_1/n_k) - \Delta}{k-1} \\ &\geq \frac{a_1(1 - (n_1/n_k)) - \Delta}{k-1} \geq \frac{N(1 - (n_1/n_k)) - \Delta_{\max}}{k-1} \geq \omega. \end{aligned}$$

Topple b_m to produce a factorization (b'_1, \dots, b'_k) , where $b'_1 \geq L/n_1$ and $b'_m \geq L/n_j$. Then $(b'_1, \dots, b'_k) \in \phi_1(Z(s-L)) \cap Z_b(s)$. We also know that $(a_1, \dots, a_k) \in \phi_1(Z(s-L)) \cap Z_a(s)$ by hypothesis. This completes Case 5.

We have covered all of the possible pairings of i and j . We conclude that $d(Z_a(s), Z_b(s)) \leq c_{\text{adj}}(s - L)$. But, a, b were arbitrary adjacent elements of $\mathcal{L}(s)$. It follows that $c_{\text{adj}}(s) \leq c_{\text{adj}}(s - L)$. This completes the proof. \square

We previously noted that $c_{\text{mon}}(s) = \max \{c_{\text{eq}}(s), c_{\text{adj}}(s)\}$. From Theorems 3.8 and 3.9, we readily obtain the following result.

COROLLARY 3.10. *Let $S = \langle n_1, \dots, n_k \rangle$ be a numerical monoid and suppose that $s \in S$. If*

$$N = \frac{(k-1)\omega + \Delta_{\max}}{1 - (n_1/n_k)}$$

and $s \geq kN$, then $c_{\text{mon}}(s - L) \geq c_{\text{mon}}(s)$.

Combining Theorems 3.5, 3.8 and 3.9 and Corollary 3.10 with Lemma 3.2 yields a proof of Theorem 3.1.

Acknowledgement

It is a pleasure to thank the referee for valuable suggestions, which resulted in an improvement of the manuscript.

References

- [1] F. Aguiló-Gost and P. A. García-Sánchez, ‘Factorization and catenary degree in 3-generated numerical semigroups’, *Electron. Notes Discrete Math.* **34** (2009), 157–161.
- [2] J. Amos, S. T. Chapman, N. Hine and J. Paixao, ‘Sets of lengths and unions of sets of lengths do not characterize numerical monoids’, *Integers* **7** (2007), #A50.
- [3] P. Baginski, S. T. Chapman, R. Rodriguez, G. Schaeffer and Y. She, ‘On the delta set and catenary degree of Krull monoids with infinite cyclic divisor class group’, *J. Pure Appl. Algebra* **214** (2010), 1334–1339.
- [4] V. Blanco, P. A. García-Sánchez and A. Geroldinger, ‘Semigroup-theoretical characterizations of arithmetical invariants with applications to numerical monoids and Krull monoids’, *Illinois J. Math.* **55** (2011), 1385–1414.
- [5] C. Bowles, S. Chapman, N. Kaplan and D. Resier, ‘Delta sets of numerical monoids’, *J. Algebra Appl.* **5** (2006), 1–24.
- [6] S. T. Chapman, P. A. García-Sánchez and D. Llena, ‘The catenary and tame degree of numerical monoids’, *Forum Math.* **21** (2009), 117–129.
- [7] S. T. Chapman, P. A. García-Sánchez, D. Llena, V. Ponomarenko and J. C. Rosales, ‘The catenary and tame degree in finitely generated commutative cancellative monoids’, *Manuscripta Math.* **120** (2006), 253–264.
- [8] S. T. Chapman, M. Holden and T. Moore, ‘Full elasticity in atomic monoids and integral domains’, *Rocky Mountain J. Math.* **36** (2006), 1437–1455.
- [9] S. T. Chapman, R. Hoyer and N. Kaplan, ‘Delta sets of numerical monoids are eventually periodic’, *Aequationes Math.* **77** (2009), 273–279.
- [10] M. Delgado, P. A. García-Sánchez and J. Morais, ‘NumericalSgps, a GAP package for numerical semigroups’, current version number 0.97 (2011) available via <http://www.gap-system.org/>.
- [11] A. Foroutan, ‘Monotone chains of factorizations’, in: *Focus on Commutative Rings Research* (ed. A. Badawi) (Nova Science, New York, 2006), 107–130.
- [12] A. Foroutan and A. Geroldinger, *Monotone Chains of Factorizations in C-Monoids*, Lecture Notes in Pure and Applied Mathematics, 241 (Marcel Dekker, New York, 2005), 99–113.
- [13] A. Geroldinger, *The catenary degree and tameness of factorizations in weakly Krull domains*, Lecture Notes in Pure and Applied Mathematics, 189 (Marcel Dekker, New York, 1997), 113–154.

- [14] A. Geroldinger, D. J. Gryniewicz and W. A. Schmid, 'The catenary degree of Krull monoids I', *J. Théor. Nombres Bordeaux* **23** (2011), 137–169.
- [15] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations: Algebraic, Combinatorial, and Analytic Theory* (Chapman and Hall/CRC, Boca Raton, FL, 2006).
- [16] A. Geroldinger and P. Yuan, 'The monotone catenary degree of Krull monoids', *Results Math.* **63** (2013), 999–1031.
- [17] F. Halter-Koch, 'The tame degree and related invariants of non-unique factorizations', *Acta Math. Univ. Ostrav.* **16** (2008), 57–68.
- [18] M. Omidali, 'The catenary and tame degree of numerical monoids generated by generalized arithmetic sequences', *Forum Math.* **24** (2012), 627–640.
- [19] A. Philipp, 'A characterization of arithmetical invariants by the monoid of relations II: The monotone catenary degree and applications to semigroup rings', *Semigroup Forum* **81** (2010), 424–434.
- [20] J. C. Rosales and P. A. García-Sánchez, *Finitely Generated Commutative Monoids* (Nova Science, New York, 1999).
- [21] J. C. Rosales and P. A. García-Sánchez, *Numerical Semigroups*, Vol. 20 (Springer, New York, 2009).

SCOTT T. CHAPMAN, Sam Houston State University,
Department of Mathematics, Box 2206, Huntsville,
TX 77341, USA
e-mail: scott.chapman@shsu.edu

MARLY CORRALES, University of Southern California,
Department of Mathematics, 3620 S. Vermont Ave.,
KAP 104, Los Angeles,
CA 90089-2532, USA
e-mail: marly.corrales@usc.edu

ANDREW MILLER, Amherst College,
Department of Mathematics,
Box 2239 P.O. 5000, Amherst,
MA 01002-5000, USA
e-mail: admiller15@amherst.edu

CHRIS MILLER, The University of Wisconsin at Madison,
Department of Mathematics, 480 Lincoln Dr., Madison,
WI 53706-1325, USA
e-mail: crmiller2@wisc.edu

DHIR PATEL, Rutgers University, Department of Mathematics,
Hill Center for the Mathematical Sciences,
110 Frelinghuysen Rd., Piscataway,
NJ 08854-8019, USA
e-mail: dp553@eden.rutgers.edu