## INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is  $12.0 \times 20.0$  cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size  $12.0 \times 20.0$  cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of Schistosoma haematobium. Journal of Helminthology, 47, 49-59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Journal of Helminthology volume 60 • no. 1 • mar 1986

## Contents

## volume 60 • no. 1 • pages 1-78 • mar 1986

volume 60 • no. 1 • mar 1986

	Pages
Comparative histochemical studies of glycosidase activity in some helminths. T. FUJINO and Y. ISHII	1–13
Bioecology of the parasites of high altitude homeothermic host-parasite systems. I. Influence of season and temperature on infections by strobilocerci of three species of <i>Hydatigera</i> in Indian house rat. S. K. MALHOTRA	
Bioecology of the parasites of high altitude homeothermic host-parasite systems. II. Infections by strobilocerci of <i>Hydatigera balaniceps</i> (Hall), <i>H. laticollis</i> (Rudolphi) and <i>H. parva</i> (Baer) in relation to sex, size and weight of Indian house rat. S. K. MALHOTRA	
Demonstration of carboxyl and thiol protease activities in adult <i>Schistosoma mansoni</i> , <i>Dirofilaria immitis, Angiostrongylus cantonensis</i> and <i>Ascaris suum</i> . J. MAKI and T. YANAGISAWA	
The life-cycle of <i>Syphacia muris</i> Yamaguti (Nematoda: Oxyuroidea) in the laboratory rat. J. W. LEWIS and J. D'SILVA	39-46
Studies on naturally-occurring ovine fascioliasis in the Sudan. E. M. HAROUN, H. A/GADIR and A. A. GAMEEL.	47–53
Studies on heterologous resistance between <i>Schistosoma bovis</i> and <i>Fasciola gigantica</i> in Sudanese cattle. A. I. YAGI, S. A. YOUNIS, E. M. HAROUN, A. A. GAMEEL, H. O. BUSHARA and M. G. TAYLOR	
Thermotaxis in third and fourth-stage <i>Dirofilaria immitis</i> larvae. M. MOK, D. ABRAHAM, R. B. GRIEVE and C. B. THOMAS	61-64
Studies on the cryopreservation of <i>Dictyocaulus viviparus</i> (Nematoda) third-stage larvae. E. R. JAMES and R. PEACOCK	65–74
Techniques for locating isotopically labelled schistosomula of <i>Schistosoma mansoni</i> in host tissues for ultrastructural investigations. J. E. CRABTREE and R. A. WILSON	
Announcements 1	4, 30, 54, 60

Runted by Taylor & Ezancis de Batopublished Brink by cambridge Baniversity Press Hampshire RG24 OPR and published by the London School of Hygiene and Tropical Medicine