

ARTICLE

Friendships, loneliness and psychological wellbeing in older adults: a limit to the benefit of the number of friends

Alexandra Thompson* , Michael A. Smith, Andrew McNeill and Thomas V. Pollet

Department of Psychology, Northumbria University, Newcastle upon Tyne, UK

*Corresponding author. Email: a.beecham@northumbria.ac.uk

(Accepted 20 April 2022; first published online 29 July 2022)

Abstract

Loneliness is linked to many negative health outcomes and places strain on the economy and the National Health Service in the United Kingdom. To combat these issues, the determinants of loneliness need to be fully understood. Although friendships have been shown to be particularly important in relation to loneliness in older adults, this association has thus far not been explored more closely. Our exploratory study examines the relationship between number of friends and loneliness, depression, anxiety and stress in older adults. Data were obtained from 335 older adults via completion of an online survey. Measures included loneliness (UCLA Loneliness Scale version 3), depression, anxiety and stress (Depression Anxiety Stress Scales DASS-21). Participants also reported their number of close friends. Regression analyses revealed an inverse curvilinear relationship between number of friends and each of the measures tested. Breakpoint analyses demonstrated a threshold for the effect of number of friends on each of the measures (loneliness = 4, depression = 2, anxiety = 3, stress = 2). The results suggest that there is a limit to the benefit of increasing the number of friends in older adults for each of these measures. The elucidation of these optimal thresholds can inform the practice of those involved in loneliness interventions for older adults. These interventions can become more targeted; focusing on either establishing four close friendships, increasing the emotional closeness of existing friendships or concentrating resources on other determinants of loneliness in this population.

Keywords: loneliness; friendships; older adults; psychological wellbeing

Introduction

Loneliness has been defined as an unpleasant or distressing experience resulting from a perceived qualitative or quantitative deficiency in one's social relationships (Russell *et al.*, 1980; Peplau and Perlman, 1982). As such, loneliness can be emotional or social. Emotional loneliness is derived from a perception of inadequate

intimacy in relationships, whereas social loneliness is due to deficits in the quantity of social relationships (Hawkey and Cacioppo, 2010).

Transient experiences of loneliness are believed to be adaptive in that they provide motivation to form and maintain social connections in order to promote the survival of genes (Cacioppo *et al.*, 2006; Cacioppo and Hawkey, 2009; Hawkey and Cacioppo, 2010). However, sustained loneliness has been repeatedly linked to many negative psychological and physiological health outcomes across age groups. These outcomes include anxiety and depression (Barg *et al.*, 2006; Cacioppo *et al.*, 2010; Age UK South Lakeland, 2018; Lee *et al.*, 2021), suicidality (Stravynski and Boyer, 2001; Van Orden *et al.*, 2010), maladaptive stress responses (Steptoe *et al.*, 2004; Adam *et al.*, 2006), cognitive decline and Alzheimer's disease (Wilson *et al.*, 2007; Boss *et al.*, 2015; Donovan *et al.*, 2017), cardiovascular disease (Momtaz *et al.*, 2012; Valtorta *et al.*, 2016, 2018), malnutrition (Ramic *et al.*, 2011), sleep quality (Yu *et al.*, 2018), functional decline (Perissinotto *et al.*, 2012) as well as increased risk of mortality (Holt-Lunstad *et al.*, 2010, 2015).

These negative impacts have associated economic costs. The cost of loneliness to employers in the United Kingdom (UK) has been estimated as being £2.5 billion due to increased staff turnover, reduced productivity, the impact of caring responsibilities and sickness absence related to ill health (New Economics Foundation and Co-op, 2017). The monetised impact of severe loneliness has been estimated as £9,900 per person due to its impact on wellbeing, health and productivity (Peytrignet *et al.*, 2020).

Further, loneliness is becoming increasingly prevalent in later life with in excess of 1 million UK residents over the age of 50 reporting that they are chronically lonely (Abrahams, 2018). This figure is expected to increase to 2 million by 2025 (Abrahams, 2018). Importantly, as the worldwide population is ageing (United Nations Department of Economic and Social Affairs, Population Division, 2019), the negative health correlates of loneliness are linked to increased strain on the National Health Service in the UK. For example, loneliness in older adults was consistently and positively associated with the number of general practitioner visits in both cross-sectional and longitudinal analyses as well as with Emergency Department visits for women (Burns *et al.*, 2020).

Due to the multiple negative impacts demonstrated above, the UK government has implemented, or is planning to implement, several initiatives with the goal of reducing loneliness in this age group. This includes strengthening their 'Tackling Loneliness' charity network, the awarding of £31.3 million of funding to charities supporting people who experience loneliness and the announcement of an additional £7.5 million to tackle loneliness during winter (Department for Digital, Culture, Media & Sport, 2021).

Clearly loneliness has a negative impact on both individuals and society as a whole. Therefore, it is important to fully understand contributors to this experience in order to reduce its deleterious impact.

Friendships in later life

At its core, loneliness relates to a perceived lack of meaningful social connections and interactions (Reichmann, 1959; Townsend, 1968; Cacioppo and Patrick, 2008). Friendships are a type of social connection which appear to be particularly

important to older adults, with evidence indicating that people in this age group are more satisfied with their friendships than their younger counterparts (Nicolaisen and Thorsen, 2017). Additionally, research has shown that a preference for emotionally close social partners increases with age (Carstensen, 1992).

Socio-emotional selectivity theory (Carstensen *et al.*, 1999) provides a framework for this observation. This theory posits that the perception of time causes individuals to prioritise particular social goals which are in competition with one another. Younger individuals who perceive time to be expansive focus on the pursuit of future-oriented knowledge-based goals. In contrast, those who perceive time as limited, such as older adults, become more focused on present-oriented goals. These present-focused goals include the aim of emotional satisfaction. As such, selectivity of social partners in this age group is increased with a preference for high-quality relationships emerging. Older adults are thought to construct their social world to match these social goals. This leads to a reduction in social network size beginning in early adulthood and results in a network which excludes novel social partners and maintains those that are already emotionally close (English and Carstensen, 2014). Thus, highlighting the notion that existing close friendships become more important with age.

Friendships and loneliness in older adults

Studies have repeatedly shown that in older adults, friendships are a greater determinant of loneliness than relationships with family members (*e.g.* Shiovitz-Ezra and Leitsch, 2010). For example, it has been shown that interactions with friends reduce loneliness to a greater extent than interactions with close relatives, including children, grandchildren and neighbours (Lee and Ishii-Kuntz, 1987; Mullins *et al.*, 1987; Pinquart and Sörensen, 2001; Steed *et al.*, 2007). Having close friends who reside in close proximity is more important than having relatives that do (Eshbaugh, 2009) and those with networks composed mainly of kin appear to be more vulnerable to loneliness and negative psychological wellbeing (Silverstein *et al.*, 1996). Additionally, 50 per cent of those who report having no friends also report feeling lonely (Holmén *et al.*, 1992), suggesting that social connections which are not friendships, such as kinship, may be less important in terms of loneliness in some cases.

As pointed out by Pinquart and Sörensen (2001), an explanation for a stronger association between friendships and loneliness in comparison to familial relationships and loneliness may lie in the quality of those relationships. Friendships tend to be of higher quality than familial relationships, as the latter are more likely based on obligation and may also involve care-giving responsibilities (Bengtson *et al.*, 1985). Given the importance of shared experiences, interests, attitudes and lifestyle for friendships, these relationships may offer more insight into understanding loneliness than family relationships (Rawlins, 1995).

Further evidence for the importance of friendships in relation to loneliness has been demonstrated. For example, Eshbaugh (2009) found that in older women living alone, having close friends who reside within 50 miles was a significant negative predictor of loneliness. A similar finding was evident in work by Mullins and Dugan (1990) in older adults living in independent living facilities. Further, a lack of friends was reported as a reason for feeling lonely in Finnish older adults (Savikko *et al.*, 2005). Additionally, frequency of contact with friends has been

evidenced as having a negative relationship with loneliness in older adults across marital statuses (Pinquart, 2003).

Quantity of friends and loneliness

As outlined above, friendships are clearly important in terms of reducing loneliness. In this context, it is important to note that research has consistently demonstrated that the quantity of friends has a negative association with loneliness. This has been found in adolescents (Lodder *et al.*, 2017), children (Nangle *et al.*, 2003), sophomore high school students (Russell *et al.*, 2012) and older adults (Shiovitz-Ezra and Leitsch, 2010). The association between this objective parameter and subjective loneliness has repeatedly been found despite the consensus being that loneliness is a subjective experience. Despite this association, there has been limited investigation of the ways in which the number of friends contributes to this. One notable exception is research exploring the discrepancy between an individual's ideal and actual number of friends (Russell *et al.*, 2012) which finds a curvilinear relationship between this discrepancy and loneliness in college students. Here, loneliness decreases as the number of actual friendships rises towards the number of ideal friendships. Once the ideal number of friendships is passed, loneliness begins to increase again. The authors explain this non-linear relationship in relation to the cognitive-discrepancy model of loneliness (Thibaut and Kelley, 2017). Here, a loss or gain of a friend when the number of friendships is close to an individual's ideal number may be especially important for determining loneliness. However, a loss or gain may be less important when the number is far above or below this ideal number.

Studies, apart from that of Russell *et al.* (2012), have overlooked the possibility that the relationship between the quantity of friends and loneliness may be curvilinear. The presence of a non-linear relationship may be indicative of a limit to the effect of increasing friends on loneliness. Work by Brummett *et al.* (2001) suggests that the link between the number of social contacts and risk of mortality in coronary patients is non-linear. Here, the mortality rate was highest in those with three or fewer social contacts. When four social contacts were present, the risk was reduced by more than half and remained at a similar level with further additions of social contacts. It is possible that a similar relationship may be present between the number of friends and loneliness, given the previously established links between loneliness and coronary heart disease (Valtorta *et al.*, 2016). The presence of a limit makes theoretical sense as it has previously been established that there is a constraint on the number of emotionally close relationships an individual can maintain within their social network, with an increase in emotional closeness linked to a reduction in network size (Roberts *et al.*, 2009). Additionally, studies have shown that intimacy in friendships differentiates between those who are lonely and those who are not (Williams and Solano, 1983; Hamid, 1989; Drageset *et al.*, 2011), although not in all cases (Mullins and Mushel, 1992). It is therefore plausible that, as more friends are added to a person's network, the emotional closeness within the network is decreased, which in turn impacts on how lonely the individual feels; thus, creating a natural limit. It is also possible that individuals experiencing loneliness may surround themselves with acquaintances as a coping strategy which could suggest a curved association between number of friends and loneliness.

As mentioned above, there are a variety of negative psychological outcomes associated with loneliness. Similarly to loneliness, previous research has shown that friendships have been linked to improved mental health. Quantity of friends is related to reduced levels of depression in older adults (Potts, 1997). Friendship support has been linked to better affect balance in the same age group (Montpetit *et al.*, 2017). Subjective isolation from friends is linked to a greater increase in both depression and psychological distress than subjective isolation from family (Taylor *et al.*, 2018). An increase in the number of friends is linked to a reduction in stress (van der Horst and Coffé, 2012). Older adult friendships are linked to better psychological wellbeing in comparison to family relationships (Nussbaum, 1994). Similarly, friendship networks have been shown to have a stronger relationship with psychological wellbeing than kin-based networks (Cable *et al.*, 2013). Finally, a lower number of social relationships has been linked to the maintenance of low negative affect and high negative affect (Huxhold *et al.*, 2020).

Of note, friendship closeness has also been associated with reduced levels of depression in both older adults (Bishop, 2008) and adults (Taylor *et al.*, 2015). As outlined previously, increasing a person's number of friendships may reduce levels of this closeness within their social network. This reduction in closeness to network contacts may in turn lessen the impact of the number of friends on depression creating a natural limit. However, as with loneliness, there has been little exploration of whether the number of friends relates to psychological health outcomes and there has been no exploration of what an optimal number of friends may be in terms of these outcomes in this age group.

Quantity versus quality

It is important to note here that quality of friendships has often been indicated as being important in terms of loneliness. Higher-quality friendships are related to lower levels of loneliness. This association has been found in the oldest old (Long and Martin, 2000), during adolescence (Lodder *et al.*, 2017) and in children (Parker and Asher, 1993). This relationship has also been found for social relationships more generally in older adults; with higher-quality social relationships predicting lower levels of loneliness (Pinquart and Sörensen, 2001).

Quality within friendships is therefore clearly important with regards to loneliness. As it is known that relationship closeness is correlated with relationship quality (*e.g.* Crespo *et al.*, 2008), we argue that it is more appropriate to focus on the ways in which close and, therefore, high-quality friendships impact loneliness rather than more peripheral, low-quality friendships. This allows the exploration of the impact of these arguably more important friendships in relation to loneliness in older adults. Further, this encompasses both the theoretical quantitative social and qualitative emotional aspects of loneliness as highlighted above.

The present study

As illustrated, a closer inspection of the relationship between number of close friends and loneliness and its psychological health correlates is warranted. To this end, the present study aimed to extend previous findings in the following

key ways. First, the study aimed to determine the relationship between the number of close friends and loneliness along with three of its associated psychological health outcomes: depression, anxiety and stress. Crucially, the existence of a non-linear relationship was explored. A further aim was to determine the optimal number of close friends for each of these parameters. As outlined above, a negative relationship between number of close friends and loneliness and each of the associated psychological wellbeing measures has previously been established. Here, we expected to replicate those findings. However, no hypothesis was made in relation to a potential curvilinear relationship. Instead, we posed the research question:

- (1) Is there a curvilinear relationship between number of close friends and loneliness in older adults?

As the second aim was exploratory, no hypothesis was made in this regard either. Again, a research question was posed:

- (2) What is the optimal number of close friends in terms of loneliness in this age group?

Method

Design

The present study was an exploratory, cross-sectional and correlational investigation into the relationship between number of close friends and loneliness, depression, anxiety and stress in older adults.

Participants

A target sample size of 400 was pre-registered, which was primarily determined by cost. Due to time constraints, this number fell somewhat short of the target. However, this number is still sufficient to perform the analysis (one predictor per 25 cases, Harrell, 2015; *also* Schmidt, 1971; Roscoe, 1975; Tabachnik and Fidell, 2009; Austin and Steyerberg, 2015). A total of 350 UK residents aged 65 or over responded to an online questionnaire. This was advertised via the recruitment platform Prolific (Palan and Schitter, 2018), Facebook adverts and via word of mouth. The Facebook and Prolific adverts were targeted specifically to UK residents aged 65 plus. Those who completed the questionnaire via Prolific were paid £2 upon completion. To be eligible to take part in the study, participants were required to not have had a current clinical diagnosis of depression or anxiety. The initial sample consisted of 350 respondents (138 male, 211 female and one participant did not specify).

Measures

Key dependent variables

Loneliness. There were four dependent variables examined in this study. The first was loneliness as measured by the University of California, Los Angeles, Loneliness Scale version 3 (UCLA-3; Russell, 1996). The UCLA-3 is a 20-item self-

report scale designed to measure an individual's subjective feelings of loneliness. Participants are asked how often each statement is descriptive of them (e.g. How often do you feel that you lack companionship?). Items are on a scale ranging from 1 (never) to 4 (often) and nine items are reverse scored. All items are summed to give one loneliness score; a higher score represents greater levels of loneliness. Version 3 of this scale has demonstrated good internal reliability (Cronbach's $\alpha = 0.96$) (e.g. Russell, 1996), with the reliability for the present study being 0.94, and has demonstrated a uni-dimensional factor structure (Russell, 1996).

Psychological wellbeing. The three psychological wellbeing measures were all assessed via the short version of the Depression Anxiety Stress Scales (DASS-21; Lovibond and Lovibond, 1995). The DASS-21 consists of 21 items and consists of three seven-item subscales measuring depression, anxiety and stress separately. Participants are asked how much each statement has applied to them over the past week (e.g. I found it hard to wind down). Items are on a scale ranging from 0 (never) to 3 (almost always). Items are summed for each scale separately and multiplied by two to allow comparison to the original 42-item DASS. No items are reverse scored. A higher score in a subscale represents a greater level of that particular psychological state. The separate depression, anxiety and stress scales have all repeatedly demonstrated good internal reliability (all Cronbach's $\alpha > 0.80$) (e.g. Osman *et al.*, 2012). Reliability for the present study is as follows: anxiety (Cronbach's $\alpha = 0.80$), stress (Cronbach's $\alpha = 0.88$) and depression (Cronbach's $\alpha = 0.90$). The DASS-21 has repeatedly demonstrated a three-factor structure representing the three separate scales (e.g. Crawford and Henry, 2003; Norton, 2007; Sinclair *et al.*, 2012; Scholten *et al.*, 2017). Table 1 contains the descriptive statistics for each of the dependent variables.

Key independent variable

The main independent variable of interest was the number of friends participants reported having. This variable was elicited by asking participants to 'Please indicate how many close friends you currently have'. The question eliciting the number of close friends was modelled after Russell *et al.* (2012). However, instead of asking participants to indicate this number on a seven-point Likert scale ranging from 'none' to '11+', we asked participants to provide the actual number, as the number of friends is better operationalised as a count variable. The definition of a close friend was left to the interpretation of the participant based on previous findings that friendships are not easily defined in older adults (Adams *et al.*, 2000).

Covariates

Four demographic covariates were included in this study as they have previously been linked to one or more of the dependent variables. Age was included as a continuous variable. Gender was included as a categorical variable with female being the reference category. Marital status was included as a categorical variable with response options being 'In a relationship', 'Married/registered civil partnership', 'Separated but still married or in a registered civil partnership', 'Divorced' and 'Widowed'. The reference category for this variable was 'Single, never married or in a civil partnership'. Highest attained level of education was also included as a categorical variable with response options being 'Some secondary school', 'GCSE

Table 1. Sample characteristics

Characteristic	Mean (SD)	N (%)
Age	69.36 (4.29)	
Gender:		
Female		201 (60)
Male		134 (40)
Marital status:		
Single		19 (6)
In a relationship		26 (8)
Married/civil partnership		207 (62)
Separated		4 (1)
Divorced		50 (15)
Widowed		28 (8)
Undisclosed		1 (0)
Education:		
Primary school		2 (1)
Some secondary school		28 (8)
GCSEs or equivalent		66 (20)
A-level or equivalent		67 (20)
Undergraduate degree		116 (35)
Postgraduate degree		56 (17)
Number of friends	4.26 (4.11)	
Loneliness	40.22 (11.64)	
Depression	6.99 (8.01)	
Anxiety	3.68 (5.42)	
Stress	8.23 (8.12)	

Notes: N = 335. SD: standard deviation. GCSE: General Certificate of Secondary Education.


(General Certificate of Secondary Education) or equivalent', 'A-level or equivalent', 'Undergraduate degree' and 'Postgraduate degree'. The reference category for this variable was 'Primary school'.

Other measures were collected as part of a separate study but are not reported on here (the ideal number of friends, the Satisfaction With Life Scale (Diener *et al.*, 1985), as well as three other measures of loneliness – the short De Jong Gierveld Scale (de Jong-Gierveld and van Tilburg, 1999), the short Social and Emotional Loneliness Scale for Adults (DiTommaso *et al.*, 2004) and a direct measure of loneliness). For a study containing this information, see https://osf.io/5f2ph/?view_only=6617265f58804de9b0145806375bb6a6.

Table 2. Zero-order correlations for study variables

	Number of friends	UCLA	DASS Depression	DASS Anxiety
UCLA	−0.50***			
DASS Depression	−0.26***	0.59***		
DASS Anxiety	−0.16**	0.37***	0.56***	
DASS Stress	−0.13*	0.44***	0.72***	0.62***

Notes: N = 335. UCLA: University of California, Los Angeles, Loneliness Scale. DASS: Depression Anxiety Stress Scales. Significance levels: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

**Figure 1.** Loneliness as a function of the number of close friendships.

Notes: Curvilinear fit with 95 per cent confidence intervals. Breakpoint is determined by segmented regression.

Procedure

Ethical approval was granted by the local ethics committee. Data were collected between May and July 2019. Once informed consent was obtained, participants completed all measures via an online questionnaire hosted by Qualtrics ([see https://www.qualtrics.com](https://www.qualtrics.com)). The demographic measures were completed first followed by each of the scales. The order these were presented in was randomised. The questions relating to the number of ideal and actual friends were completed last. The whole questionnaire took no longer than 20 minutes to complete.

Analytical approach

All analyses were performed using R version 3.6.1 (R Core Team, 2017). Observations which contained missing data for the independent variable and dependent variables were removed (13 cases), as were two outliers where the number of friends was listed as 105 (30 standard deviations (SD) away from mean) and age as 66,123 years. Finally, due to the presence of extreme values and asymmetry

Table 3. Hierarchical ordinary least squares regression analysis to predict loneliness

	Model 1	Model 2	Model 3	Model 4	Model 5
Number of friends	−4.591*** (0.499)	−4.725*** (0.496)	−4.757*** (0.498)	−4.705*** (0.501)	−4.783*** (0.500)
Number of friends ²	0.253*** (0.044)	0.268*** (0.044)	0.272*** (0.044)	0.266*** (0.045)	0.274*** (0.045)
Marital status (Ref. Single or undisclosed):					
In a relationship		−6.151* (2.858)	−6.235* (2.862)	−6.401* (2.867)	−5.243† (2.899)
Married/civil partnership		−6.946** (2.243)	−6.896** (2.245)	−7.139** (2.259)	−6.290** (2.285)
Separated or divorced		−3.958 (2.505)	−3.882 (2.508)	−3.738 (2.513)	−3.198 (2.520)
Widowed		−4.347 (2.814)	−3.953 (2.863)	−3.861 (2.865)	−3.022 (2.871)
Age			−0.096 (0.127)	−0.115 (0.128)	−0.090 (0.131)
Gender male (Ref. Female)				1.123 (1.138)	0.622 (1.158)
Education (Ref. Primary or some secondary school):					
GCSE or equivalent					−1.938 (2.133)
A-level or equivalent					1.734 (2.151)
Undergraduate degree					1.849 (2.026)
Postgraduate degree					0.636 (2.228)
Constant	52.212*** (1.156)	58.144*** (2.467)	64.784*** (9.088)	65.734*** (9.139)	62.786*** (9.917)
R ²	0.316	0.342	0.343	0.345	0.360
Adjusted R ²	0.312	0.330	0.329	0.329	0.336
Residual standard error	9.656 (df = 332)	9.528 (df = 328)	9.534 (df = 327)	9.535 (df = 326)	9.486 (df = 332)
F statistic	76.766*** (df = 2; 332)	28.443*** (df = 6; 328)	24.430*** (df = 7; 327)	21.497*** (df = 8; 326)	15.091*** (df = 12; 322)

Notes: N = 335. Standard errors are in parentheses. Ref.: reference category. GCSE: General Certificate of Secondary Education. df: degrees of freedom. Significance levels: † $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Figure 2. Depression as a function of the number of close friendships.

Notes: Curvilinear fit with 95 per cent confidence intervals. Breakpoint is determined by segmented regression.

(skewness = -1.37), the data were winsorised to 3 SD for the number of friends (12 friends) variable (12 cases). This resulted in a final sample of 335 participants; 290 of whom were recruited via Prolific and 45 via social media.

The data met the assumptions of non-multicollinearity, homoscedasticity and non-autocorrelation. The loneliness data met the assumption of normally distributed residuals. The three DASS variables all exhibited some skew in the distribution of residuals. However, this is to be expected given that most respondents were not currently experiencing high levels of depression, anxiety or stress. However, regression is relatively robust, therefore we used this technique (Berry, 1993) and we further assessed the robustness of these models with other techniques (*e.g.* segmented regression).

Prior to performing regression analyses, some demographic factor levels were grouped to result in a more equal N per cell. For marital status, single and undisclosed were combined into one level. As were separated and divorced. For education, primary and secondary school were combined. Based on a visual inspection of the data and a plot of residuals, the fit of polynomial regression models was assessed for all dependent variables. It was found that a quadratic model had the best fit to the data in each case. This was based on the model with a statistically significant reduction in residual sum of squares as indicated by a chi-squared difference test. A Davies Test was then employed to test for a non-constant regression parameter in the predictor. This was followed by segmented regression via the segmented package (Muggeo, 2008) and multivariate adaptive regression spline analysis ('mars') via the earth package (Milborrow *et al.*, 2011) to determine and confirm a breakpoint in the data. The segmented package utilises an algorithm which determines the breakpoint iteratively. Similarly, the earth package uses an algorithmic approach to examine breakpoints in the data. We also performed additional analyses including robustness checks (see https://osf.io/5f2ph/?view_only=6617265f58804de9b0145806375bb6a6).

Table 4. Hierarchical ordinary least squares regression analysis to predict depression

	Model 1	Model 2	Model 3	Model 4	Model 5
Number of friends	-2.096*** (0.393)	-2.066*** (0.397)	-2.016*** (0.397)	-2.136*** (0.395)	-2.162*** (0.39)
Number of friends ²	0.135*** (0.035)	0.135*** (0.035)	0.130*** (0.035)	0.143*** (0.035)	0.146*** (0.035)
Marital status (Ref. Single or undisclosed):					
In a relationship		1.973 (2.287)	2.105 (2.284)	2.490 (2.262)	2.809 (2.309)
Married/civil partnership		0.132 (1.794)	0.053 (1.792)	0.617 (1.782)	0.822 (1.820)
Separated or divorced		-0.195 (2.004)	-0.314 (2.002)	-0.649 (1.983)	-0.501 (2.008)
Widowed		1.164 (2.252)	0.543 (2.285)	0.328 (2.261)	0.600 (2.287)
Age			0.152 (0.101)	0.196† (0.101)	0.196† (0.104)
Gender male (Ref. Female)				-2.606** (0.898)	-2.721** (0.922)
Education (Ref. Primary or some secondary school):					
GCSE or equivalent					-1.142 (1.699)
A level or equivalent					-0.059 (1.713)
Undergraduate degree					0.088 (1.614)
Postgraduate degree					-0.357 (1.775)
Constant	11.982*** (0.910)	11.564*** (1.974)	1.083*** (7.252)	-1.120*** (7.211)	-0.940*** (7.900)
R ²	0.105	0.110	0.116	0.139	0.142
Adjusted R ²	0.100	0.094	0.097	0.117	0.110
Residual standard error	7.599 (df = 332)	7.623 (df = 328)	7.609 (df = 327)	7.524 (df = 326)	7.556 (df = 332)
F statistic	19.488*** (df = 2; 332)	6.776*** (df = 6; 328)	6.153*** (df = 7; 327)	6.559*** (df = 8; 326)	4.434*** (df = 12; 322)

Notes: N = 335. Standard errors are in parentheses. Ref.: reference category. GCSE: General Certificate of Secondary Education. df: degrees of freedom. Significance levels: † $p < 0.1$, ** $p < 0.01$, *** $p < 0.001$.




Figure 3. Anxiety as a function of the number of friendships.

Notes: Curvilinear fit with 95 per cent confidence intervals. Breakpoint is determined by segmented regression.

Results

Sample characteristics

Sample characteristics and descriptive statistics are presented in [Table 1](#). In the final sample, 60 per cent of participants were female and 40 per cent were male. The mean age of participants was 69 years ($SD = 4.29$). Most participants reported being married or in a civil partnership (62%) and 35 per cent reported having completed an undergraduate degree. The average number of close friends reported was four. All were residents of the UK.

Main analysis

Correlations

The bivariate correlations are presented in [Table 2](#). Number of friends was significantly negatively associated with each of the dependent variables. Loneliness was significantly positively associated with each of the psychological wellbeing variables.

Regressions

Hierarchical ordinary least squares regression analysis with the inclusion of a quadratic term was performed for each of the outcome measures to test for a curvilinear relationship.

Loneliness. After adjusting for all demographic information, the final model was significant ($F(12,322) = 15.09, p < 0.001$). In this model, both the linear term for number of close friends ($B = -4.78, p < 0.001$) and the quadratic term ($B = 0.27, p < 0.001$) remained significant. As quadratic relationships are difficult to interpret from coefficients alone, see [Figure 1](#).

Of the demographic covariates, only being married or in a civil partnership, in comparison to being single, was a significant predictor of loneliness ($B = -6.29, p =$

Table 5. Hierarchical ordinary least squares regression analysis to predict anxiety

	Model 1	Model 2	Model 3	Model 4	Model 5
Number of friends	−0.980*** (0.275)	−0.944*** (0.276)	−0.941*** (0.278)	−0.976*** (0.279)	−0.970*** (0.280)
Number of friends ²	0.066** (0.024)	0.064** (0.024)	0.064** (0.025)	0.068** (0.025)	0.067** (0.025)
Marital status (Ref. Single or undisclosed):		3.207* (1.591)	3.213* (1.595)	3.322* (1.597)	2.905† (1.620)
In a relationship					
Married/civil partnership		1.400 (1.249)	1.397 (1.251)	1.588 (1.258)	1.185 (1.277)
Separated or divorced		0.988 (1.394)	0.983 (1.398)	0.887 (1.399)	0.720 (1.408)
Widowed		0.749 (1.567)	0.723 (1.596)	0.662 (1.596)	0.444 (1.604)
Age			0.006 (0.071)	0.019 (0.071)	−0.015 (0.073)
Gender male (Ref. Female)				−7.45 (0.634)	−0.469 (0.647)
Education (Ref. Primary or some secondary school):					
GCSE or equivalent					−1.230 (1.192)
A-level or equivalent					−2.142† (1.201)
Undergraduate degree					−2.305* (1.132)
Postgraduate degree					−1.397 (1.245)
Constant	5.943*** (0.636)	4.508** (1.373)	4.069 (5.064)	3.440 (5.089)	7.716 (5.540)
R ²	0.046	0.061	0.061	0.065	0.080
Adjusted R ²	0.040	0.043	0.041	0.042	0.045
Residual standard error	5.313 (df = 332)	5.305 (df = 328)	5.313 (df = 327)	5.310 (df = 326)	5.299 (df = 332)
F statistic	8.026*** (df = 2; 332)	3.529** (df = 6; 328)	3.017** (df = 7; 327)	2.815** (df = 8; 326)	2.327*** (df = 12; 322)

Notes: N = 335. Standard errors are in parentheses. Ref.: reference category. GCSE: General Certificate of Secondary Education. df: degrees of freedom.

Significance levels: † $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Figure 4. Stress as a function of the number of close friendships.

Notes: Curvilinear fit with 95 per cent confidence intervals. Breakpoint is determined by segmented regression.

0.006). Those with this particular marital status reported less loneliness than those who were single. This final model explained 34 per cent of the variance in loneliness scores (Table 3).

As suggested by the plot, there appears to be a point at which the effect of the number of close friends on loneliness is greatly reduced. A Davies Test indicated that there was a significant difference between the slopes when segmenting the regression line at 3.90 friends ($p < 0.001$). Breakpoint analysis via the ‘segmented’ package further supported that a breakpoint could be elicited at this point ($\psi = 3.86$, 95% confidence interval (CI) = 2.77, 4.94). Multivariate adaptive regression spline analysis further suggested a knot at four close friends. This analysis suggests that the effect of additional close friends on loneliness is diminished once around four close friends are reached.

Depression. After adjusting for all demographic information, the final model was significant ($F(12,322) = 4.43$, $p < 0.001$). In this model, both the linear term for number of close friends ($B = -2.16$, $p < 0.001$) and the quadratic term ($B = 0.15$, $p < 0.001$) remained significant. The plot in Figure 2 demonstrates this relationship.

Table 4 summarises all models on depression. Of the demographic covariates, gender was a significant predictor of depression ($B = -2.72$, $p < 0.01$), with men reporting less depression than women. Educational attainment was also a significant predictor of depression (Table 4: all dummy variables $p < 0.01$). Those who reported that they had received an education past primary school level reported less depression than those that had not. This model explained 11 per cent of the variance in depression scores.

Similar to the effect for loneliness, there appeared to be a point at which the effect of number of close friends on depression is greatly reduced. A Davies Test indicated that there was a significant difference between the slopes when

Table 6. Hierarchical ordinary least squares regression analysis to predict stress

	Model 1	Model 2	Model 3	Model 4	Model 5
Number of friends	-1.462*** (0.412)	-1.396*** (0.413)	-1.374** (0.415)	-1.454*** (0.416)	-1.511*** (0.418)
Number of friends ²	0.104*** (0.037)	0.103*** (0.037)	0.100** (0.037)	0.109** (0.037)	0.112** (0.037)
Marital status (Ref. Single or undisclosed):					
In a relationship		5.440* (2.380)	5.499* (2.384)	5.754* (2.380)	6.148 (2.421)
Married/civil partnership		1.607 (1.868)	1.571 (1.871)	1.946 (1.875)	2.249 (1.908)
Separated or divorced		0.726 (2.086)	0.672 (2.089)	0.450 (2.086)	0.741 (2.104)
Widowed		1.760 (2.344)	1.482 (2.385)	1.339 (2.378)	1.563 (2.397)
Age			0.068 (0.105)	0.097 (0.106)	0.089 (0.109)
Gender male (Ref. Female):				-1.732† (0.945)	-1.797 (0.967)
Education (Ref. Primary or some secondary school):					
GCSE or equivalent					-1.582 (1.781)
A-level or equivalent					-0.132 (1.796)
Undergraduate degree					-0.909 (1.691)
Postgraduate degree					0.935 (1.860)
Constant	11.449*** (0.955)	9.555*** (2.055)	4.858 (7.571)	3.394 (7.586)	4.349 (8.280)
R ²	0.042	0.063	0.064	0.074	0.084
Adjusted R ²	0.036	0.046	0.044	0.051	0.049
Residual standard error	7.976 (df = 332)	7.936 (df = 328)	7.943 (df = 327)	7.914 (df = 326)	7.920 (df = 332)
F statistic	7.194*** (df = 2; 332)	3.661** (df = 6; 328)	3.192** (df = 7; 327)	3.233** (df = 8; 326)	2.448** (df = 12; 322)

Notes: N = 335. Standard errors are in parentheses. Ref.: reference category. GCSE: General Certificate of Secondary Education. df: degrees of freedom.

Significance levels: † $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

segmenting the regression line at 2.16 friends ($p < 0.001$). Breakpoint analysis via the 'segmented' package in R confirmed that a breakpoint could be elicited at this point ($\psi = 2.17$, standard error (SE) = 0.491, 95% CI = 1.19, 3.12). Multivariate adaptive regression spline analysis further suggested a knot at two close friends. This analysis suggests that the effect of additional close friends on depression is diminished once a cut-off of around two close friends is reached.

Anxiety. After adjusting for all demographic information, the final model was significant ($F(15,319) = 1.74$, $p = 0.043$). In the final model, both the linear term for number of close friends ($B = -0.74$, $p = 0.001$) and the quadratic term ($B = 0.04$, $p = 0.014$) remained significant. This quadratic relationship is illustrated in Figure 3. None of the demographic covariates were significant predictors of anxiety. This model explained 3.2 per cent of the variance in anxiety scores. For all coefficients, see Table 5.

Figure 3 suggests a breakpoint after which there is no additional reduction from adding close friends. A Davies Test indicated that there was a significant difference between the slopes when segmenting the regression line at 2.51 friends ($p < 0.01$). Breakpoint analysis confirmed that a breakpoint could be elicited at this point ($\psi = 2.51$, 95% CI = 1.02, 4.0). Multivariate adaptive regression spline analysis further suggested a knot at three close friends. This analysis suggests that the effect of additional close friends on anxiety is diminished once a threshold of approximately three close friends is reached.

Stress. After adjusting for all demographic information, the final model was significant ($F(12,322) = 2.448$, $p < 0.001$). In this model both the linear term for number of close friends ($B = -1.51$, $p < 0.01$) and the quadratic term ($B = 0.11$, $p < 0.05$) remained significant. This relationship is illustrated in Figure 4.

Of the demographic covariates, being in a relationship was a significant predictor of stress ($B = 6.15$, $p < 0.05$). Those who were in a relationship reported more stress than those who were single. This model explained 4.9 per cent of the variance in stress scores. For all coefficients, see Table 6.

Figure 4 suggests the presence of a threshold above which there is no further reduction in stress with a further increase of reported close friends. A Davies Test indicated that there was a significant difference between the slopes when segmenting the regression line at 2.05 friends ($p < 0.05$). Breakpoint analysis confirmed that a breakpoint could be elicited at this point ($\psi = 2.05$, SE = 0.64, 95% CI = 0.80, 3.30). Multivariate adaptive regression spline analysis further suggested a knot at two friends. This analysis suggests that the effect of additional close friends on stress is diminished once around two friends are reached.

Discussion

The purpose of this study was to explore the relationship between the reported number of close friends and loneliness, depression, anxiety and stress in older adults. Our findings demonstrated significant inverse curvilinear relationships between the number of close friends and each of these parameters. These findings support those of previous studies which demonstrated a negative relationship between the quantity of friends and loneliness (Mullins and Dugan, 1990;

Nangle *et al.*, 2003; Shiovitz-Ezra and Leitsch, 2010; Russell *et al.*, 2012; Lodder *et al.*, 2017). The present findings are also in line with previous work demonstrating that having a greater quantity of friends is associated with better self-reported mental health (Nussbaum, 1994; Potts, 1997; Bishop, 2008; van der Horst and Coffé, 2012; Cable *et al.*, 2013; Montpetit *et al.*, 2017; Taylor *et al.*, 2018; Huxhold *et al.*, 2020). Russell *et al.* (2012) have previously demonstrated that there is an inverse curvilinear relationship between the discrepancy between the number of actual and ideal friendships and loneliness in college students. However, we believe a curvilinear relationship for the effect of close friends on each outcome measure has gone untested previously.

Inspection of the plots for each of the outcome measures shows that there are relatively few individuals who report having many friends whilst simultaneously experiencing high levels of loneliness, depression, anxiety or stress. We have discussed the notion that having a large social network may reduce emotional closeness across the relationships within it (Roberts *et al.*, 2009) and, in turn, this lack of emotional closeness may increase feelings of loneliness (Williams and Solano, 1983; Hamid, 1989; Drageset *et al.*, 2011). It may be that whilst these individuals report having many close friends, these relationships are not actually providing the closeness, connection and understanding needed to stave off loneliness and its maladaptive correlates due to the size of their networks. However, further work, ideally of a longitudinal nature, is needed to corroborate this.

It is therefore possible that a small cluster of observations could be driving the quadratic fit. However, we believe that this is unlikely to be the case. Bootstrapping, which reduces the weight of extreme cases, further supported the curvilinear relationship (see https://osf.io/5f2ph/?view_only=6617265f58804de9b0145806375bb6a6). Further, we conducted segmented regression using both the 'segmented' and 'earth' packages to determine breakpoints in the data. These approaches reduce the effect of outliers and consistently detected breakpoints in the data (see https://osf.io/5f2ph/?view_only=6617265f58804de9b0145806375bb6a6).

Thus, each of our mental health-related variables demonstrated that there was a point past which the addition of more close friends no longer has a substantial beneficial effect. An increase of close friends was associated with a decrease in loneliness until four close friends were reached, for depression this number was found to be two close friends, for anxiety three close friends and for stress the threshold was two close friends. A slightly larger threshold value for loneliness in comparison to the other mental health variables is to be expected, as loneliness is more directly related to social connections and interactions than the other psychological well-being outcomes. The presence of a friendship threshold in terms of loneliness could be due to a reduction in emotional closeness to network members as network size is increased (Roberts *et al.*, 2009).

It is encouraging that the threshold for the number of close friends in each instance appears to be relatively low. Making and maintaining meaningful social connections takes time and effort (Lang *et al.*, 2013). However, given that on average a person's closest group of contacts, known as their support group, has been found to include around five members (Dunbar and Spoor, 1995), it is possible that many older adults will already have met this number, or be close to it. The elucidation of these thresholds for close friendships is important as they can allow

those involved in loneliness and mental health interventions to focus their limited resources on increasing social interaction opportunities for those with few or no close friends. This would be with the ultimate aim of reducing the negative physical and psychological impacts of loneliness as well as reducing the economic cost and strain on health services associated with loneliness. As research has previously shown that perceived quality of relationships has a protective effect against loneliness (Pinquart, 2003; Hawkley *et al.*, 2008), those with more friendships than a threshold could be encouraged to improve the quality and closeness of their friendships, rather than increase the number of friends they have.

A successful intervention to reduce loneliness could include elements of the previously successful 'Friendship Program' (Hamid, 1989). This programme encourages reflection on one's own aspirations for friendships as well as reflection on current friendships. It also incorporates education around attitudes and experiences in the process of building friendships. This includes actively utilising those already-established but lesser-used connections for support and by encouraging a proactive approach to maintaining and deepening friendships. The programme has been found to both improve the quality of current friendships as well to increase the number of friendships. Further benefits of improving the quality of these relationships include having increased resilience to adversity (Graber *et al.*, 2016) and a more adaptive stress response (Calhoun *et al.*, 2014).

Alternatively, these individuals could be included in interventions with the aim of improving the other contributing factors to loneliness and psychological wellbeing such as health status and mobility (Theeke, 2009). These could include cognitive-based interventions to improve mobility (Marusic *et al.*, 2018) and physical activity interventions to improve mobility (Yeom *et al.*, 2009) and health status (Hwang and Braun, 2015); as well as directly targeting psychological wellbeing with the aim of reducing loneliness in tandem. In this regard, a variety of interventions have been shown to be successful, including psychotherapy and behavioural interventions (Pinquart *et al.*, 2007). In particular, reminiscence therapy has repeatedly demonstrated a positive outcome in relation to depressive symptoms and subjective wellbeing (for a review, see Yen and Lin, 2018). This type of therapy involves recalling events from the past and sharing them with an observer or group who listen without making comment.

The present study has a number of strengths. To our knowledge, it is the first to explore the possibility of a curvilinear relationship between the quantity of close friends and loneliness, as well as its associated psychological wellbeing outcomes in an older adult population. It is also the first to assess the presence of breakpoints within this relationship. As mentioned previously, for both of these analyses robust techniques were employed to increase confidence in our findings.

We also focused explicitly on close friendships and did not include any peripheral friendships. This is because emotional closeness has repeatedly been shown to be more important than network size and frequency and contact with friends in terms of loneliness (Williams and Solano, 1983; Hamid, 1989; Drageset *et al.*, 2011). Close friends also appear to be more important than less-close friends in regards to subjective wellbeing (van der Horst and Coffé, 2012). We asked participants specifically to include only friends whom they thought of as close, although the interpretation of a close friend was left to the participant; this reduced any

confusion about who to include in this number. This clarity has not always been present in previous studies (e.g. Pinqart, 2003; Savikko *et al.*, 2005; Steed *et al.*, 2007).

However, the current study is not without its limitations. Firstly, the present research was an online study only available to those with access to the internet. Many older adults do not have this access (Yu *et al.*, 2016) and/or autonomy with internet use (Hargittai *et al.*, 2019) and so the current findings are not representative of the UK population. Second, the definition of a close friend was left to each participant's own interpretation. Respondents were able to decide what a close friend meant to them, and to allocate their personal contacts accordingly. Although in previous studies, similar to ours, no definition for friendships has been given (e.g. Mullins and Dugan, 1990; Russell *et al.*, 2012), individual interpretations may differ between participants and providing an explicit definition, such as in the study of Williams and Solano (1983), may have resulted in different findings. However, we believe that leaving this concept open to interpretation allows for more accurate inclusion of those to whom respondents feel close, rather than forcing them to exclude contacts based on a definition which may be incompatible with respondents' own views. This is something that has been cautioned against previously by Adams *et al.* (2000), who outline that it is not suitable to assign a definition of friendship to individuals as many do not share the same criteria for this type of relationship. Therefore, future research may benefit from repeating the study to include face-to-face data collection for those without access to the internet and potentially the incorporation of an established definition of a close friend and comparing if this differs from a respondent's definition.

As mentioned previously, we focused on only close friendships. Doing so allowed the determination of the number of high-quality and rewarding friendships that individuals should focus on to reduce loneliness and improve psychological wellbeing. This is important as these emotionally close friendships require bilateral effort, time and other resources to initiate and maintain and, as such, are more costly in comparison to less-emotionally close relationships (Roberts and Dunbar, 2011). However, despite the importance of these close relationships in terms of loneliness, more peripheral friendships could still be having an impact on our outcome measures. It may be beneficial for future work to take into account these friendships also.

Further, the present study did not explicitly take into account the impact of the quality of friendships on loneliness and mental health. Although we have mentioned the minority of participants who may have been experiencing low levels of closeness in the friendships they reported, as participants were asked *specifically* about close friends, it is likely that many of these contacts represent high-quality relationships. Regardless, as the quality of friendships has been linked to loneliness and mental health (e.g. Wheeler *et al.*, 1983; Mullins and Dugan, 1990), future studies would benefit from also including the quality or emotional closeness of these friendships explicitly as a covariate to determine whether the effect of the number of friends remains after adjusting for this.

Similarly, the present study did not account for the types of support that can be exchanged within friendships. Both emotional and instrumental support within social networks have been shown to predict levels of loneliness (e.g. Sanchez

et al., 2014). Including the type of support both given and received within close friendships in future analyses could assist in unpicking the aspects of these friendships which are most optimal in preventing loneliness in this age group.

It is important to note that the number of close friendships is not the only aspect of an individual's social network with implications for loneliness within this age group. Structural and compositional aspects of this network can also explain loneliness and its associated negative outcomes. For example, it has been demonstrated that social network size as well as brokerage and embeddedness can predict loneliness in older adults (Kim *et al.*, 2021). Compositionally, the presence of a romantic partner is protective against loneliness (Martina, 2021). This could potentially be due to the increased opportunity for accruing additional friends within a social network. The inclusion of such parameters in the above analyses would provide a deeper understanding of the effect of the number of close friendships on loneliness in older adults after controlling for these aspects.

In terms of additional future directions, as well as a replication, and taking into account the aforementioned issues, future research should establish who these close friends are. We know that these are emotionally close relationships but is there a difference in the impact on loneliness and psychological wellbeing in terms of the length of the relationship and whether these contacts belong to a dense network? Is it preferable to have contact with these friends in a certain way for a certain length of time or at a particular frequency? Or is the content of the interactions more important? Previous research suggests that network density is important in terms of happiness and subjective wellbeing (Huang *et al.*, 2019) and that frequency of contact has a differential effect on subjective wellbeing dependent on the mode utilised (van der Horst and Coffé, 2012). Therefore, these nuances are possible and should be explored in order to inform interventions further. Importantly, the current investigation was cross-sectional in nature. It may well be possible that there is a limit to the benefit of increasing the number of friendships over time. It also may be possible that having a larger number of friends leads to improved wellbeing. However, we cannot rule out the possibility of reverse causation. A longitudinal design would be necessary to determine this in future.

For now, we conclude that there is a nonlinear relationship between the reported number of friends and loneliness, depression, anxiety and stress in older adults. We further conclude that based on the present data there appears to be a limit to the beneficial effect of increasing the number of friends in this population. The findings are in need of corroboration incorporating a more representative sample of this population and future work is necessary to qualify further the nature of these friendships. However, these findings have important implications in the development and provision of loneliness and psychological wellbeing interventions.

Financial support. This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Conflict of interest. The authors declare no conflicts of interest.

Ethical standards. Ethical approval was obtained from Ethics Committee of the Faculty of Health and Life Sciences at Northumbria University (reference number 16881).

References

- Abrahams C (2018) *All the Lonely People: Loneliness in Later Life*. London: Age UK.
- Adam EK, Hawkey LC, Kudielka BM and Cacioppo JT (2006) Day-to-day dynamics of experience – cortisol associations in a population-based sample of older adults. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 17058–17063.
- Adams RG, Blieszner R and DeVries B (2000) Definitions of friendship in the third age: age, gender, and study location effects. *Journal of Ageing Studies* **14**, 117–133.
- Age UK South Lakeland (2018) *Loneliness, Depression and Anxiety: Exploring the Connection to Mental Health Through Testing Promising Approaches to Reducing Loneliness in South Lakeland*. London: Age UK.
- Austin PC and Steyerberg EW (2015) The number of subjects per variable required in linear regression analyses. *Journal of Clinical Epidemiology* **68**, 627–636.
- Barg FK, Huss-Ashmore R, Wittink MN, Murray GF, Bogner HR and Gallo JJ (2006) A mixed-methods approach to understanding loneliness and depression in older adults. *Journals of Gerontology: Series B* **61**, 329–339.
- Bengtson VL, Cutler NE, Mangen DJ and Marshall VW (1985) Generations, cohorts, and relations between age groups. In Binstock RH and Shanas E (eds), *Handbook of Aging and the Social Sciences*. New York, NY: Van Nostrand Reinhold, pp. 304–338.
- Berry WD (1993) *Understanding Regression Assumptions*. Los Angeles, CA: Sage.
- Bishop AJ (2008) Stress and depression among older residents in religious monasteries: do friends and God matter? *International Journal of Aging and Human Development* **67**, 1–23.
- Boss L, Kang DH and Branson S (2015) Loneliness and cognitive function in the older adult: a systematic review. *International Psychogeriatrics* **27**, 541–553.
- Brummett BH, Barefoot JC, Siegler IC, Clapp-Channing NE, Lytle BL, Bosworth HB, Williams RB and Mark DB (2001) Characteristics of socially isolated patients with coronary artery disease who are at elevated risk for mortality. *Psychosomatic Medicine* **63**, 267–272.
- Burns A, Leavey G, Ward M and O'Sullivan R (2020) The impact of loneliness on healthcare use in older people: evidence from a nationally representative cohort. *Journal of Public Health* 1–10.
- Cable N, Bartley M, Chandola T and Sacker A (2013) Friends are equally important to men and women, but family matters more for men's well-being. *Journal of Epidemiology and Community Health* **67**, 166–171.
- Cacioppo JT and Hawkey LC (2009) Perceived social isolation and cognition. *Trends in Cognitive Sciences* **13**, 447–454.
- Cacioppo JT, Hawkey LC, Ernst JM, Burleson M, Berntson GG, Nouriani B and Spiegel D (2006) Loneliness within a nomological net: an evolutionary perspective. *Journal of Research in Personality* **40**, 1054–1085.
- Cacioppo J, Hawkey L and Thisted R (2010) Perceived social isolation makes me sad. *Psychology and Aging* **25**, 453–463.
- Cacioppo JT and Patrick W (2008) *Loneliness: Human Nature and the Need for Social Connection*. New York, NY: W.W. Norton & Company.
- Calhoun CD, Helms SW, Heilbrin N, Rudolph KD, Hasting PD and Prinstein MJ (2014) Relational victimization, friendship, and adolescents' hypothalamic–pituitary–adrenal axis responses to an in vivo social stressor. *Developmental Psychopathology* **26**, 605–618.
- Carstensen LL (1992) Social and emotional patterns in adulthood: support for socioemotional selectivity theory. *Psychology and Aging* **7**, 331–338.
- Carstensen LL, Isaacowitz DM and Charles ST (1999) Taking time seriously: a theory of socioemotional selectivity. *American Psychologist* **54**, 165–181.
- Crawford JR and Henry JD (2003) The Depression Anxiety Stress Scales (DASS): normative data and latent structure in a large non-clinical sample. *British Journal of Clinical Psychology* **42**, 111–131.
- Crespo C, Davide IN, Costa ME and Fletcher GJO (2008) Family rituals in married couples: links with attachment, relationship quality, and closeness. *Personal Relationships* **15**, 191–203.
- de Jong-Gierveld J and van Tilburg T (1999) *Manual of the Loneliness Scale*. Amsterdam: Department of Social Research Methodology, Vrije Universiteit.
- Department for Digital, Culture, Media & Sport (2021) *Loneliness Annual Report*. London: Department for Digital, Culture, Media & Sport.

- Diener E, Emmons RA and Larsen RJ** (1985) The Satisfaction With Life Scale. *Journal of Personality Assessment* **49**, 71–75.
- DiTommaso E, Brannen C and Best LA** (2004) Measurement and validity characteristics of the short version of the Social and Emotional Loneliness Scale for Adults. *Educational and Psychological Measurement* **64**, 99–119.
- Donovan NJ, Wu Q, Rentz DM, Sperling RA, Marshall GA and Glymour MM** (2017) Loneliness, depression and cognitive function in older U.S. adults. *International Journal of Geriatric Psychiatry* **32**, 564–573.
- Drageset J, Kirkevold M and Espehaug B** (2011) Loneliness and social support among nursing home residents without cognitive impairment: a questionnaire survey. *International Journal of Nursing Studies* **48**, 611–619.
- Dunbar RIM and Spoors M** (1995) Social networks, support cliques, and kinship. *Human Nature* **6**, 273–290.
- English T and Carstensen LL** (2014) Selective narrowing of social networks across adulthood is associated with improved emotional experience in daily life. *International Journal of Behavioral Development* **38**, 195–202.
- Eshbaugh EM** (2009) The role of friends in predicting loneliness among older women living alone. *Journal of Gerontological Nursing* **35**, 13–16.
- Graber R, Turner R and Madill A** (2016) Best friends and better coping: facilitating psychological resilience through boys' and girls' closest friendships. *British Journal of Psychology* **107**, 338–358.
- Hamid P** (1989) Contact and intimacy patterns of lonely students. *New Zealand Journal of Psychology* **18**, 84–86.
- Hargittai E, Piper AM and Morris MR** (2019) From internet access to internet skills: digital inequality among older adults. *Universal Access in the Information Society* **18**, 881–890.
- Harrell FE** (2015) *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis*. Cham, Switzerland: Springer.
- Hawkey LC and Cacioppo JT** (2010) Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Annals of Behavioral Medicine* **40**, 218–227.
- Hawkey LC, Hughes ME, Waite LJ, Masi CM, Thisted RA and Cacioppo JT** (2008) From social structural factors to perceptions of relationship quality and loneliness: the Chicago Health, Aging, and Social Relations Study. *Journals of Gerontology: Series B* **63**, 375–384.
- Holmén K, Ericsson K, Andersson L and Winblad B** (1992) Loneliness among elderly people living in Stockholm: a population study. *Journal of Advanced Nursing* **17**, 43–51.
- Holt-Lunstad J, Smith TB, Baker M, Harris T and Stephenson D** (2015) Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspectives on Psychological Science* **10**, 227–237.
- Holt-Lunstad J, Smith TB and Layton JB** (2010) Social relationships and mortality risk: a meta-analytic review. *PLOS Medicine* **7**, e1000316.
- Huang X, Western M, Bian Y, Li Y, Côté R and Huang Y** (2019) Social networks and subjective wellbeing in Australia: new evidence from a national survey. *Sociology* **53**, 401–421.
- Huxhold O, Fiori KL, Webster NJ and Antonucci TC** (2020) The strength of weaker ties: an underexplored resource for maintaining emotional well-being in later life. *Journals of Gerontology: Series B* **75**, 1433–1442.
- Hwang PW-N and Braun KL** (2015) The effectiveness of dance interventions to improve older adults' health: a systematic literature review. *Alternative Therapies in Health and Medicine* **21**, 64–70.
- Kim H, Kwak S, Youm Y and Chey J** (2021) Social network characteristics predict loneliness in older adults. *Gerontology* **68**, 309–320.
- Lang FR, Wagner J, Wrzus C and Neyer FJ** (2013) Personal effort in social relationships across adulthood. *Psychology and Aging* **28**, 529–539.
- Lee GR and Ishii-Kuntz M** (1987) Social interaction, loneliness, and emotional well-being among the elderly. *Research on Aging* **9**, 459–482.
- Lee SL, Pearce E, Ajnakina O, Johnson S, Lewis G, Mann F, Pitman A, Solmi F, Sommerlad A, Steptoe A, Tymoszuk U and Lewis G** (2021) The association between loneliness and depressive symptoms among adults aged 50 years and older: a 12-year population-based cohort study. *The Lancet Psychiatry* **8**, 48–57.

- Lodder GMA, Scholte RHJ, Goossens L and Verhagen M** (2017) Loneliness in early adolescence: friendship quantity, friendship quality, and dyadic processes. *Journal of Clinical Child and Adolescent Psychology* **46**, 709–720.
- Long MV and Martin P** (2000) Personality, relationship closeness, and loneliness of oldest old adults and their children. *Journals of Gerontology: Series B* **55**, 311–319.
- Lovibond PF and Lovibond SH** (1995) *Manual for the Depression Anxiety Stress Scales*, 2nd Edn. Sydney: Psychology Foundation.
- Martina Š** (2021) Marital status, close social network and loneliness of older adults in the Czech Republic. *Ageing & Society* **41**, 671–685.
- Marusic U, Verghese J and Mahoney JR** (2018) Cognitive-based interventions to improve mobility: a systematic review and meta-analysis. *Journal of the American Medical Directors Association* **19**, 484–491.
- Milborrow S, Hastie T and Tibshirani R** (2011) *Earth: Multivariate Adaptive Regression Spline Models*. R Package. The Comprehensive R Archive Network.
- Montaz YA, Hamid TA, Yusoff S, Ibrahim R, Chai ST, Yahaya N and Abdullah SS** (2012) Loneliness as a risk factor for hypertension in later life. *Journal of Aging and Health* **24**, 696–710.
- Montpetit MA, Nelson NA and Tiberio SS** (2017) Daily interactions and affect in older adulthood: family, friends, and perceived support. *Journal of Happiness Studies* **18**, 373–388.
- Muggeo MRV** (2008) Segmented: an R package to fit regression models with broken-line relationships. *R News* **8**, 20–25.
- Mullins LC and Dugan E** (1990) The influence of depression, and family and friendship relations, on residents' loneliness in congregate housing. *The Gerontologist* **30**, 377–384.
- Mullins LC, Johnson P and Andersson L** (1987) Loneliness of the elderly: the impact of family and friends. *Journal of Social Behavior and Personality* **2**, 225–238.
- Mullins LC and Mushel M** (1992) The existence and emotional closeness of relationships with children, friends, and spouses: the effect on loneliness among older persons. *Research on Aging* **14**, 448–470.
- Nangle DW, Erdley CA, Newman JE, Mason CA and Carpenter EM** (2003) Popularity, friendship quantity, and friendship quality: interactive influences on children's loneliness and depression. *Journal of Clinical Child & Adolescent Psychology* **32**, 546–555.
- New Economics Foundation and Co-op** (2017) *The Cost of Loneliness to UK Employers*. New Economics Foundation.
- Nicolaisen M and Thorsen K** (2017) What are friends for? Friendships and loneliness over the lifespan – from 18 to 79 years. *International Journal of Aging and Human Development* **84**, 126–158.
- Norton PJ** (2007) Depression Anxiety and Stress Scales (DASS-21): psychometric analysis across four racial groups. *Anxiety, Stress and Coping* **20**, 253–265.
- Nussbaum JF** (1994) Friendship in older adulthood. In Hummert ML, Wiemann JM and Nussbaum JM (eds), *Interpersonal Communication in Older Adulthood*. Thousand Oaks, CA: Sage, pp. 209–225.
- Osman A, Wong JL, Bagge CL, Freedenthal S, Gutierrez PM and Lozano G** (2012) The Depression Anxiety Stress Scales-21 (DASS-21): further examination of dimensions, scale reliability, and correlates. *Journal of Clinical Psychology* **68**, 1322–1338.
- Palan S and Schitter C** (2018) Prolific.ac – a subject pool for online experiments. *Journal of Behavioral and Experimental Finance* **17**, 22–27.
- Parker JG and Asher SR** (1993) Friendship and friendship quality in middle childhood: links with peer group acceptance and feelings of loneliness and social dissatisfaction. *Developmental Psychology* **29**, 611–621.
- Peplau LA and Perlman D** (1982) *Loneliness: A Sourcebook of Current Theory, Research and Therapy*. New York, NY: Wiley.
- Perissinotto CM, Stijacic Cenzer I and Covinsky KE** (2012) Loneliness in older persons: a predictor of functional decline and death. *Archives of Internal Medicine* **172**, 1078–1083.
- Peytrignet S, Garforth-Bles S and Way R** (2020) *Loneliness Monetisation Report*. Simetrica: London.
- Pinquart M** (2003) Loneliness in married, widowed, divorced, and never-married older adults. *Journal of Social and Personal Relationships* **20**, 31–53.
- Pinquart M, Duberstein PR and Lyness JM** (2007) Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: a meta-analysis. *Ageing and Mental Health* **11**, 645–657.

- Pinquart M and Sörensen S** (2001) Influences on loneliness in older adults: a meta-analysis. *Basic and Applied Social Psychology* **23**, 245–266.
- Potts MK** (1997) Social support and depression among older adults living alone: the importance of friends within and outside of a retirement community. *Social Work* **42**, 348–362.
- Ramic E, Pranjic N, Batic-Mujanovic O, Karic E, Alibasic E and Alic A** (2011) The effect of loneliness on malnutrition in elderly population. *Medicinski Arhiv* **65**, 92–95.
- Rawlins WK** (1995) Friendships in later life. In Coupland J and Nussbaum JF (eds), *Handbook of Communication and Aging Research*. Mahwah, NJ: Routledge, pp. 227–257.
- R Core Team** (2017) *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Reichmann FF** (1959) Loneliness. *Psychiatry* **22**, 1–15.
- Roberts SGB and Dunbar RIM** (2011) Communication in social networks: effects of kinship, network size, and emotional closeness. *Personal Relationships* **18**, 439–452.
- Roberts SGB, Dunbar RIM, Pollet TV and Kuppens T** (2009) Exploring variation in active network size: constraints and ego characteristics. *Social Networks* **31**, 138–146.
- Roscoe JT** (1975) *Fundamental Research Statistics for the Behavioral Sciences*, 2nd Edn. Ann Arbor, MI: Holt, Rinehart and Winston.
- Russell D** (1996) UCLA Loneliness Scale version 3: reliability, validity and factor structure. *Journal of Personality Assessment* **66**, 3–4.
- Russell DW, Cutrona CE, McRae C and Gomez M** (2012) Is loneliness the same as being alone? *Journal of Psychology* **146**, 7–22.
- Russell D, Peplau LA and Cutrona CE** (1980) The revised UCLA Loneliness Scale: concurrent and discriminant validity evidence. *Journal of Personality and Social Psychology* **39**, 472–480.
- Sanchez M, Gierveld JDEJ and Delgado JBUZ** (2014) Loneliness and the exchange of social support among older adults in Spain and the Netherlands. *Ageing & Society* **34**, 330–354.
- Savikko N, Routasalo P, Tilvis RS, Strandberg TE and Pitkälä KH** (2005) Predictors and subjective causes of loneliness in an aged population. *Archives of Gerontology and Geriatrics* **41**, 223–233.
- Schmidt FL** (1971) The relative efficiency of regression and simple unit predictor weights in applied differential psychology. *Educational and Psychological Measurement* **31**, 699–714.
- Scholten S, Velten J, Bieda A, Chi Zhang X and Margraf J** (2017) Testing measurement invariance of the Depression, Anxiety, and Stress Scales (DASS-21) across four countries. *Psychological Assessment* **29**, 1376.
- Shiovitz-Ezra S and Leitsch SA** (2010) The role of social relationships in predicting loneliness: the National Social Life, Health, and Aging Project. *Social Work Research* **34**, 157–167.
- Silverstein M, Chen X and Heller K** (1996) Too much of a good thing? Intergenerational social support and the psychological well-being of older parents. *Journal of Marriage and the Family* **58**, 970.
- Sinclair SJ, Siefert CJ, Slavin-Mulford JM, Stein MB, Renna M and Blais MA** (2012) Psychometric evaluation and normative data for the Depression, Anxiety, and Stress Scales-21 (DASS-21) in a non-clinical sample of U.S. Adults. *Evaluation & the Health Professions* **35**, 259–279.
- Steed L, Boldy D, Grenade L and Iredell H** (2007) The demographics of loneliness among older people in Perth, Western Australia. *Australasian Journal on Ageing* **26**, 81–86.
- Step toe A, Owen N, Kunz-Ebrecht SR and Brydon L** (2004) Loneliness and neuroendocrine, cardiovascular, and inflammatory stress responses in middle-aged men and women. *Psychoneuroendocrinology* **29**, 593–611.
- Stravynski A and Boyer R** (2001) Loneliness in relation to suicide ideation and parasuicide: a population-wide study. *Suicide and Life-threatening Behavior* **31**, 32–40.
- Tabachnik BG and Fidell LS** (2009) *Using Multivariate Statistics*, 7th Edn. Boston, MA: Pearson.
- Taylor HO, Taylor RJ, Nguyen AW and Chatters L** (2018) Social isolation, depression, and psychological distress among older adults. *Journal of Aging and Health* **30**, 229–246.
- Taylor RJ, Chae DH, Lincoln KD and Chatters LM** (2015) Extended family and friendship support networks are both protective and risk factors for major depressive disorder and depressive symptoms among African-Americans and Black Caribbeans. *Journal of Nervous and Mental Disease* **203**, 132–140.
- Theeke LA** (2009) Predictors of loneliness in U.S. adults over age sixty-five. *Archives of Psychiatric Nursing* **23**, 387–396.
- Thibaut JW and Kelley HH** (2017) *The Social Psychology of Groups*. Abingdon, UK: Routledge.

- Townsend P** (1968) Isolation, desolation and loneliness. In Shanas E, Townsend P, Wedderburn D, Frus H, Milhaj P and Stehouwer J (eds), *Older People in Three Industrial Societies*. New York, NY: Atherton, pp. 258–287.
- United Nations Department of Economic and Social Affairs, Population Division** (2019) *World Population Ageing 2019*. New York, NY: United Nations Department of Economic and Social Affairs, Population Division.
- Valtorta NK, Kanaan M, Gilbody S and Hanratty B** (2018) Loneliness, social isolation and risk of cardiovascular disease in the English Longitudinal Study of Ageing. *European Journal of Preventive Cardiology* **25**, 1387–1396.
- Valtorta NK, Kanaan M, Gilbody S, Ronzi S and Hanratty B** (2016) Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. *Heart* **102**, 1009–1016.
- van der Horst M and Coffé H** (2012) How friendship network characteristics influence subjective well-being. *Social Indicators Research* **107**, 509–529.
- Van Orden KA, Witte TK, Cukrowicz KC, Braithwaite SR, Selby EA and Joiner TE** (2010) The interpersonal theory of suicide. *Psychological Review* **117**, 575–600.
- Wheeler L, Reis H and Nezlek JB** (1983) Loneliness, social interaction, and sex roles. *Journal of Personality and Social Psychology* **45**, 943–953.
- Williams JG and Solano CH** (1983) The social reality of feeling lonely: friendship and reciprocation. *Personality and Social Psychology Bulletin* **9**, 237–242.
- Wilson RS, Krueger KR, Arnold SE, Schneider JA, Kelly JF, Barnes LL, Tang Y and Bennett DA** (2007) Loneliness and risk of Alzheimer disease. *Archives of General Psychiatry* **64**, 234–240.
- Yen HY and Lin LJ** (2018) A systematic review of reminiscence therapy for older adults in Taiwan. *Journal of Nursing Research* **26**, 138–150.
- Yeom HA, Keller C and Fleury J** (2009) Interventions for promoting mobility in community-dwelling older adults. *Journal of the American Academy of Nurse Practitioners* **21**, 95–100.
- Yu B, Steptoe A, Niu K, Ku PW and Chen LJ** (2018) Prospective associations of social isolation and loneliness with poor sleep quality in older adults. *Quality of Life Research* **27**, 683–691.
- Yu RP, Ellison NB, McCammon RJ and Langa KM** (2016) Mapping the two levels of digital divide: internet access and social network site adoption among older adults in the USA. *Information Communication and Society* **19**, 1445–1464.

Cite this article: Thompson A, Smith MA, McNeill A, Pollet TV (2024). Friendships, loneliness and psychological wellbeing in older adults: a limit to the benefit of the number of friends. *Ageing & Society* **44**, 1090–1115. <https://doi.org/10.1017/S0144686X22000666>