

Physician aid-in dying and mental disorders

S0110

Pad, psychiatry and stigma

E. Kowalinski and C. Huber

Psychiatry, University of Basel, Basel, Switzerland

*Corresponding Author.

doi: 10.1192/j.eurpsy.2021.114

In 2018, the Swiss Academy of Medical Sciences (SAMW) published a new guideline on physician-assisted dying (PAD). In line with the SAMW guideline published in 2004, the patients' ability to judge, their self-determination, careful consideration and permanence of their wish to die as well as the lack of therapeutic options were set as necessary conditions. However, while the previous wording considered assisted suicide to be ethically justifiable if the patient's condition is terminal, the new guideline requires that it is unbearable. This difference has been the subject of intense discussion in Swiss health-care professionals and the population alike. This controversy is particularly important for those affected by mental illness who have a persistent desire to die. This is because mental disorders cannot usually be classified as terminal illnesses, but they can certainly lead to suffering that is perceived as unbearable. Furthermore, it is known that persons with mental illness are subject to stigmatization. It is therefore likely that there is a connection between the stigmatization of mentally ill people and the position on PAD for this group. This talk provides theoretical background on this discussion and proposes a study protocol to investigate the acceptance of PAD in relation to the type of illness as well as the factors of unbearable suffering and terminality. It will furthermore look into the criteria of the 2004 and 2018 guidelines and will explore if there is a connection between stigmatization and the assessment of whether a person should be granted access to assisted suicide.

Disclosure: No significant relationships.

Keywords: physician-assisted dying; Mental illness; suffering; terminality

S0107

Pad in forensic psychiatry

I. Franke

Forensic Psychiatry And Psychotherapy, Ulm University, Guenzburg, Germany

doi: 10.1192/j.eurpsy.2021.115

Introduction: A recent court decision in Germany defined assisted suicide as a basic human right. Consequently, the discussion regarding PAD needs to be extended to people who are in forensic/secure psychiatric hospitals or prisons, sometimes without any prospects of release. Several studies have shown that long-term hospitalization and detention are associated with feelings of hopelessness, depression and suicidal ideations. Moreover, the resources for adequate therapy are often rare. This results in complex moral challenges for mental health care.

Objectives: To review current practices in countries that allow PAD and to discuss ethical conflicts.

Methods: Literature review; international comparison of current regulations.

Results: A majority of the literature on PAD in detention refers to prisoners with terminal medical conditions. Single case reports of PAD-requests of mentally disordered offenders aroused great public interest. The resulting ethical conflicts are similar to those issues regarding PAD and mental disorder in general. However, in secure treatment settings and detention additional aspects such as adverse living conditions and inadequate access to mental health care need to be taken into account.

Conclusions: If unbearable pain is not a precondition for assisted suicide, then mentally disordered and healthy offenders have a right to request PAD, provided they have medical decision-making capacity. Considering the common insufficient mental health care for people in detention, policy and law makers need to ensure that access to PAD will not replace therapy. Professionals involved in PAD evaluations need support by specific guidelines.

Disclosure: No significant relationships.

Keywords: PAD; prison psychiatry; ethics; decision-making

S0108

Ethics of pad in mental disorders

M. Trachsel^{1,2,3*} and R. Jox⁴

¹Clinical Ethics Unit, University Hospital of Basel (USB), Basel, Switzerland; ²Clinical Ethics Unit, University Psychiatric Clinics (UPK) Basel, Basel, Switzerland; ³Institute Of Biomedical Ethics And History Of Medicine, University of Zurich (UZH), Zurich, Switzerland and ⁴Institute Of Humanities In Medicine, Lausanne University Hospital, Lausanne, Switzerland

*Corresponding Author.

doi: 10.1192/j.eurpsy.2021.116

Involuntary psychiatric hospitalization for suicide prevention and physician assistance in dying (PAD) for patients with severe and persistent mental illness (SPMI) combine to create a moral tension. Switzerland has the longest history of non-medicalized assistance in dying, considered as a civil right even beyond pathological situations. The debate in Switzerland centers on the notion of suffering in the context of PAD. In 2018, the Swiss Academy of Medical Sciences revised their end-of-life policy stipulating intolerable suffering due to severe illness or functional limitations and acknowledged as such by the physician as a core criterion for PAD. However, we argue that suffering is a necessary but insufficient condition for PAD, the other criteria being decision-making capacity (DMC) and refractoriness of the suffering. Moreover, we hold that suffering is a subjective experience that can only be quantified by the patient and cannot be compared between two persons in an objective way. According to this concept, however, some patients with SPMI, refractory suffering, and preserved DMC will meet the criteria for PAD. Therefore, we call for palliative care approaches in psychiatry which includes relief of suffering as much as possible, but also accepting PAD after a conscientious assessment of the criteria.

Disclosure: No significant relationships.

Keywords: physician assistance in dying; decision-making capacity; ethics; Mental illness