

Short Communication

Highlighting the ratio of sodium to potassium in population-level dietary assessments: cross-sectional data from New York City, USA

Stella S Yi^{1,*†}, Christine J Curtis¹, Sonia Y Angell^{1,‡}, Cheryl AM Anderson², Molly Jung³ and Susan M Kansagra^{1,§}

¹New York City Department of Health and Mental Hygiene, Bureau of Chronic Disease Prevention and Tobacco Control, 2 Gotham Center, 42-09 28th Street, 9th Floor, CN-46, Long Island City, NY 11101, USA; ²University of California, San Diego School of Medicine, Department of Family and Preventive Medicine, La Jolla, CA, USA;

³Albert Einstein College of Medicine, Department of Epidemiology and Population Health, New York, NY, USA

Submitted 14 February 2014; Final revision received 19 May 2014; Accepted 20 May 2014; First published online 20 June 2014

Abstract

Objective: To contrast mean values of Na:K with Na and K mean intakes by demographic factors, and to calculate the prevalence of New York City (NYC) adults meeting the WHO guideline for optimal Na:K (<1 mmol/mmol, i.e. <0.59 mg/mg) using 24 h urinary values.

Design: Data were from the 2010 Community Health Survey Heart Follow-Up Study, a population-based, representative study including data from 24 h urine collections.

Setting: Participants were interviewed using a dual-frame sample design consisting of random-digit dial telephone exchanges that cover NYC. Data were weighted to be representative of NYC adults as a whole.

Subjects: The final sample of 1656 adults provided 24 h urine collections and self-reported health data.

Results: Mean Na:K in NYC adults was 1.7 mg/mg. Elevated Na:K was observed in young, minority, low-education and high-poverty adults. Only 5.2% of NYC adults had Na:K in the optimal range.

Conclusions: Na intake is high and K intake is low in NYC adults, leading to high Na:K. Na:K is a useful marker and its inclusion for nutrition surveillance in populations, in addition to Na and K intakes, is indicated.

Keywords
Sodium
Potassium
Surveillance

High Na intake and low K intake are independently associated with an increased risk of developing high blood pressure^(1,2). To achieve dietary recommendations, most adults would need to decrease their Na intake and increase their K intake^(3,4), resulting in a decreased Na:K. An elevated Na:K is associated with higher blood pressure⁽⁵⁾ and poor cardiovascular outcomes⁽⁶⁾, and is attractive as a potential surveillance tool of population-level dietary habits. Although the USA has no formal

recommendation for Na:K⁽⁷⁾, the WHO defined an optimal Na:K as <1.0 mmol/mmol (i.e. <0.6 mg/mg)⁽⁸⁾. The objectives of the current analysis were to present mean urinary Na:K values from a population-based sample and to report the percentage of New Yorkers meeting the WHO-defined optimal Na:K.

Methods

Data were obtained from the New York City (NYC) Heart Follow-Up Study (HFUS), a cross-sectional study conducted in 2010 to assess Na intake in a population-based, representative sample of NYC adults aged 18 years and older. Detailed study information has been described in

† Present affiliation: New York University School of Medicine, Department of Population Health, New York, NY, USA.

‡ Present affiliation: Centers for Disease Control and Prevention, Atlanta, GA, USA.

§ Present affiliation: New York City Department of Health and Mental Hygiene, Division of Health Promotion and Disease Prevention, New York, NY, USA.




Fig. 1 Heart Follow-Up Study (HFUS): participant flow and response rates

a comprehensive Methods Report⁽⁹⁾. Study participants meeting the inclusion criteria were recruited from the Community Health Survey (CHS), an annual health survey conducted by the NYC Health Department. The CHS is designed after the national Behavioral Risk Factor Surveillance Survey (BRFSS) and includes questions either from national surveys or questions that have been validated by pre-testing in the NYC adult population. In 2010, more than 10 000 New Yorkers were interviewed using a dual-frame, stratified random sample design consisting of random-digit dial of landline and cellular telephone exchanges that cover NYC⁽¹⁰⁾. This recruitment design produces a sample that may be weighted to produce estimates that are representative of the NYC adult non-institutionalized population as a whole. Throughout the recruitment process, participation rates were closely monitored through daily and weekly reports by NYC Health Department survey staff.

Following the CHS questionnaire, participants meeting the three inclusion criteria (not pregnant, not breast-feeding or lactating, and not on current or past dialysis) were invited to participate in the HFUS. HFUS participants performed a 24 h urine collection; prior protocols were used as a basis for the design of the 24 h urine collection⁽¹¹⁾ but were modified to accommodate the unique needs of NYC and our study. Following the collection period, a home visit was scheduled with a medical technician. The home visit did not occur unless a signed informed consent was collected at the start of the visit. At the home visit, which followed a strict clinical protocol⁽¹²⁾, an aliquot of the 24 h urine collection was collected. Two specimen tubes of aliquoted urine were included in case the first tube was damaged during shipment or the laboratory needed to redo the urinalysis. The urine samples were sent to a collaborating laboratory for the analysis of Na and K. The laboratory staff flagged any values

that appeared to be incongruous with the volume of urine collected. No samples were flagged in this process. Results were sent via hard copy (laboratory values per participant) and via spreadsheet to the NYC Health Department, and were checked using double data entry from the original hard copy results. The Institutional Review Board of the NYC Health Department approved the study.

The 2010 CHS response rates were 17% and 28% for landline and cellular telephone exchanges, respectively. Cooperation rates among those who were reached were 77% among landline contacts and 94% among cellular contacts. Of the 2010 CHS participants screened for HFUS participation, 5830 eligible adults were identified. A total of 2305 agreed to provide a 24 h urine sample and of these, 1775 (30%) provided a sample that could be analysed by the laboratory. The overall participant flow is displayed in Fig. 1. Those who agreed to participate were slightly more likely than those who did not to be Hispanic, <65 years of age, lower income and obese; however, there were no meaningful differences in self-reported high blood pressure or general health status between CHS and HFUS participants⁽⁹⁾. Incomplete samples were defined as those provided by participants who reported missing a collection and as those with a total urine volume <500 ml or urine creatinine <6.05 mmol creatinine in males and <3.78 mmol creatinine in females (both being biologically implausible; Elliott P & Stamler J, personal communication, 2011). The final analytic sample size was 1656.

Measurement and definition of covariates

Na and K contents of 24 h urine samples were determined using the ion-selective electrode potentiometric method on the Roche DPP Modular analyser (Hoffman-La Roche, Ltd, USA). Laboratory values were normalized to a 24 h period prior to analysis. Results are presented in mg or

mg/mg. BMI was calculated from measured weight in kilograms divided by the square of height in metres. Demographic (age, sex, race/ethnicity, poverty, education, nativity) and other characteristics (smoking, physical activity, fruit and vegetable intake, fast-food consumption, soda consumption) were either included in the CHS or in the questionnaire asked of only the subset of HFUS participants. Poverty was assessed as annual combined household income and was grouped according to federal poverty guidelines. Nativity (US-born *v.* foreign-born) was determined by asking state and country of birth. Because Na intake is related to diet, which in turn is heavily influenced by culture, those reporting being born in Puerto Rico or other US Territories were considered 'foreign-born'. Physical activity was assessed using the validated question, 'During the past 30 d, other than your regular job, did you participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise?'. Fruit and vegetable intake was reported as servings/d and categorized (none, 1 to 4, 5+). Fast-food consumption was assessed as a continuous variable with the question, 'How often do you eat something from a fast-food restaurant or chain such as: McDonalds, KFC, Taco Bell, Golden Krust or similar places?', and was categorized as times/week (none, <1 (more than 0), 1 to <2, 2 to <3, 3 to <6). Per-day soda consumption was assessed as a continuous variable with the question, 'How often do you drink sugar sweetened soda? Do not include diet soda or seltzer', and categorized as a binary variable (two or more sodas daily, yes or no).

Statistical analysis

Differences in mean Na:K, Na intake and K intake by subgroups were assessed using *t* tests. The *P* trend across ordinal variable categories was assessed using Wald χ^2 tests. Multivariable linear regression models were used to understand factors associated with higher Na:K after adjustment for age, sex, race/ethnicity, poverty, education, nativity (foreign-born *v.* US-born), BMI, smoking status and physical activity. Comparisons of those with Na:K < 0.6 *v.* Na:K \geq 0.6 by subgroup were assessed using χ^2 tests. Analyses were performed incorporating sampling weights to account for complex survey design and non-response. Data were analysed using the SUDAAN statistical software package version 10.0.

Results

The final sample of 1656 participants was weighted to be representative of the NYC population to estimate Na and K intakes in all NYC adults. Over 75% of participants were between the ages of 25 and 64 years, and about half were male; half were Black or Hispanic; about half were foreign-born; 48% had an income less than 200% and 15% had income between 200 and 399% of the federal poverty level. The majority had less than a college

education (69%) and was US-born (58%). The population was evenly distributed as normal weight, overweight and obese, and about 40% were current or former smokers, 26% were physically inactive over the past 30 d, 11% reported no servings of fruits and vegetables the previous day, 49% ate at a fast-food chain in the past week, and 10% drank two or more sodas daily.

Mean Na:K in NYC adults was 1.7 mg/mg, with higher Na:K observed in: younger *v.* older adults; Black and Hispanic race/ethnic subgroups *v.* Whites; those with a higher *v.* a lower poverty level; those with a lower *v.* a higher educational attainment; US-born *v.* foreign-born; and obese *v.* normal BMI (Table 1). A higher Na:K was also associated with self-reported less healthy behaviours (high physical inactivity; low fruit and vegetable intake; high fast-food intake; and consumption of two or more sodas daily) compared with healthier behaviours. The observed patterns did not differ substantially in multivariable-adjusted analyses, but differences observed for poverty level and physical activity were no longer statistically significant (results not shown).

Only 5.2% of New Yorkers had Na:K < 0.6 mg/mg. Being 45 years or older, White, engaging in healthier behaviours, having a higher income, having a college degree and being at a healthy weight were associated with Na:K < 0.6 mg/mg (data not shown).

Discussion

In adult NYC residents, the mean Na:K assessed by biomarker data was 1.7 mg/mg, due to both high Na intake and low K intake. Only 5.2% of the NYC population had an optimal Na:K, suggesting there are very few who meet dietary recommendations for Na and K intakes. These findings are consistent with national data from the National Health and Nutrition Examination Survey 2003–2008 using 24 h dietary recall⁽⁴⁾.

In our analysis, young adults (age 18–25 years) had the highest Na:K. Further, a less-than-optimal Na:K was associated with obesity and physical inactivity, both independent risk factors for CVD. The association with younger age raises concerns about poor dietary behaviours in early adulthood, which could increase the risk of premature CVD⁽¹³⁾.

Adhering to the daily recommended Na limit for most adults (2300 mg) and the daily recommended K intake (4700 mg) would result in an Na:K of < 0.5 mg/mg. Even fewer NYC adults would meet this lower ratio. The main dietary recommendation to increase K is to increase fruit and vegetable consumption; this was corroborated in our study, since a lower Na:K was associated with five or more daily servings of fruits and vegetables. A lower Na:K was also associated with lower fast-food and sugary drink consumption, suggesting that environmental changes to lower consumption of these foods, while increasing access to fruits and vegetables, could help shift the population Na:K.

Table 1 Mean sodium:potassium, sodium and potassium intakes; 2010 Community Health Survey Heart Follow-Up Study (HFUS), New York City, USA

| | Mean Na:K | | Mean Na (mg/d) | | Mean K (mg/d) | |
|--|---------------|----------|----------------|------------|---------------|------------|
| | Weighted mean | 95 % CI | Weighted mean | 95 % CI | Weighted mean | 95 % CI |
| Overall | 1.7 | 1.6, 1.8 | 3239 | 3119, 3359 | 2175 | 2093, 2257 |
| Age group | | | | | | |
| 18–24 years | 2.2* | 2.0, 2.4 | 3445 | 3067, 3823 | 1716* | 1532, 1899 |
| 25–44 years (Ref.)† | 1.7 | 1.6, 1.8 | 3248 | 3038, 3457 | 2190 | 2036, 2343 |
| 45–64 years | 1.7 | 1.6, 1.8 | 3467 | 3286, 3649 | 2356 | 2235, 2477 |
| ≥ 65 years | 1.3* | 1.2, 1.4 | 2630* | 2452, 2809 | 2201 | 2071, 2330 |
| <i>P</i> trend | < 0.01 | | < 0.01 | | < 0.01 | |
| Sex | | | | | | |
| Male | 1.7 | 1.6, 1.8 | 3583* | 3392, 3775 | 2387* | 2241, 2533 |
| Female (Ref.) | 1.7 | 1.6, 1.8 | 2944 | 2799, 3088 | 1993 | 1907, 2078 |
| Race | | | | | | |
| Non-Hispanic White (Ref.) | 1.4 | 1.3, 1.5 | 3066 | 2892, 3240 | 2524 | 2369, 2679 |
| Non-Hispanic Black | 2.1* | 2.0, 2.3 | 3477* | 3204, 3751 | 1847* | 1690, 2004 |
| Hispanic | 1.8* | 1.7, 1.9 | 3395* | 3153, 3637 | 2021* | 1897, 2144 |
| Non-Hispanic Asian | 1.6 | 1.4, 1.8 | 3068 | 2626, 3510 | 2034* | 1869, 2198 |
| Other | 1.7§ | 1.3, 2.1 | 2992§ | 2649, 3335 | 1975*,§ | 1619, 2331 |
| Poverty income ratio | | | | | | |
| < 200 % FPL | 1.8* | 1.7, 2.0 | 3306 | 3123, 3488 | 1987* | 1890, 2084 |
| 200–399 % FPL | 1.7* | 1.5, 1.9 | 3350 | 3076, 3624 | 2168* | 2015, 2322 |
| ≥ 400 % FPL (Ref.) | 1.4 | 1.3, 1.5 | 3200 | 2983, 3416 | 2553 | 2358, 2749 |
| <i>P</i> trend | < 0.01 | | 0.20 | | < 0.01 | |
| Education | | | | | | |
| Less than high school | 1.9* | 1.8, 2.1 | 3376 | 3084, 3669 | 1938* | 1778, 2097 |
| Grade 12 or GED | 1.9* | 1.7, 2.0 | 3365 | 3111, 3619 | 1988* | 1862, 2113 |
| Some college | 1.7* | 1.6, 1.9 | 3166 | 2936, 3396 | 2060* | 1925, 2196 |
| College graduate (Ref.) | 1.4 | 1.3, 1.5 | 3096 | 2901, 3290 | 2576 | 2394, 2758 |
| <i>P</i> trend | < 0.01 | | 0.25 | | < 0.01 | |
| Country of birth | | | | | | |
| US-born (Ref.) | 1.8 | 1.7, 1.9 | 3299 | 3135, 3462 | 2132 | 2016, 2249 |
| Foreign-born‡ | 1.6* | 1.5, 1.7 | 3176 | 2999, 3353 | 2227 | 2110, 2343 |
| BMI category | | | | | | |
| Underweight, < 18.5 kg/m ² | 1.9§ | 1.1, 2.7 | 3130§ | 2103, 4157 | 1852*,§ | 1514, 2190 |
| Normal, 18.5–< 25.0 kg/m ² (Ref.) | 1.5 | 1.4, 1.7 | 3040 | 2835, 3245 | 2228 | 2100, 2355 |
| Overweight, 25.0–< 30.0 kg/m ² | 1.6 | 1.5, 1.7 | 3114 | 2925, 3303 | 2151 | 1997, 2304 |
| Obese, ≥ 30.0 kg/m ² | 1.9* | 1.8, 2.0 | 3609* | 3381, 3837 | 2179 | 2021, 2337 |
| <i>P</i> trend | < 0.01 | | < 0.01 | | 0.24 | |
| Smoking | | | | | | |
| Never (Ref.) | 1.7 | 1.6, 1.8 | 3149 | 2999, 3299 | 2124 | 2015, 2234 |
| Current | 1.9 | 1.7, 2.0 | 3496* | 3195, 3796 | 2143 | 1934, 2351 |
| Former | 1.6 | 1.4, 1.7 | 3287 | 3025, 3549 | 2334* | 2182, 2486 |
| Performed physical activity in last 30 d | | | | | | |
| Yes | 1.6 | 1.6, 1.7 | 3152 | 3014, 3290 | 2197 | 2101, 2293 |
| No (Ref.) | 1.9* | 1.7, 2.0 | 3491* | 3256, 3727 | 2110 | 1950, 2271 |
| Fruit and vegetable intake | | | | | | |
| None (Ref.) | 1.9 | 1.8, 2.1 | 3296 | 2969, 3624 | 1842 | 1655, 2029 |
| 1 to 4 servings/d | 1.7* | 1.6, 1.8 | 3270 | 3127, 3414 | 2166* | 2069, 2263 |
| 5+ servings/d | 1.4* | 1.2, 1.5 | 3050 | 2756, 3345 | 2531* | 2300, 2762 |
| <i>P</i> trend | < 0.01 | | 0.39 | | < 0.01 | |
| Times per week eat fast food | | | | | | |
| None (Ref.) | 1.5 | 1.4, 1.6 | 3126 | 2969, 3283 | 2385 | 2258, 2512 |
| < 1 | 1.8* | 1.7, 2.0 | 3293 | 3045, 3542 | 1965* | 1837, 2093 |
| 1 to < 2 | 1.9* | 1.7, 2.0 | 3513 | 3125, 3902 | 2031* | 1819, 2243 |
| 2 to < 3 | 2.0* | 1.7, 2.2 | 3604 | 3036, 4172 | 2075 | 1766, 2383 |
| 3 to < 6 | 2.0* | 1.8, 2.3 | 3200 | 2745, 3655 | 1650* | 1371, 1929 |
| <i>P</i> trend | < 0.01 | | 0.23 | | < 0.01 | |
| Two or more sodas daily | | | | | | |
| Yes | 2.2* | 2.0, 2.5 | 3915* | 3416, 4413 | 1911 | 1630, 2192 |
| No (Ref.) | 1.6 | 1.6, 1.7 | 3166 | 3048, 3284 | 2202 | 2116, 2288 |

Ref., reference category; FPL, federal poverty level; GED, General Educational Development.

P value for trend is calculated for ordinal, categorical variables using the Wald χ^2 test.

*Statistical significance at *P* < 0.05, *P* value compares mean in subgroup to reference.

†Reference group is 25–44-year-olds due to smaller sample size in the 18–24-year-old group.

‡Because diet and culture heavily influence Na and K intakes, those reporting being born in Puerto Rico or other US Territories were considered 'foreign-born'.

§Estimate's relative standard error is greater than 30 % or the sample size is less than fifty, making the estimate potentially unreliable.

Although 24 h urinary Na excretion is the gold standard measure of Na and K intake, it reflects ~97% of Na consumed but only ~80–85% of K intake⁽¹⁴⁾. Thus, the urinary Na:K used in the current analysis may overestimate the true ratio. While the current study used the gold standard methodology of estimating Na intake through the measurement of rigorously collected 24 h urine, limitations to this methodology exist. The potential for incomplete collection of urine by the participant is a primary limitation of this methodology. From study design to implementation, to laboratory and data analysis, the HFUS investigators took this into consideration and has been previously described⁽¹⁵⁾. Another limitation is that one 24 h collection may not be representative of an individual's typical intake given the high intra-person variability of Na excretion, although this is less problematic when assessing estimates at the population level. As an additional note, the units of Na:K differ depending on the measurement method (mg *v.* mmol). This affects the magnitude of the effect of a one-unit change, and should be considered when interpreting results across studies. Lastly, the results presented are only representative of the non-institutionalized adult population. While participation of different subgroups in the HFUS was similar to the overall CHS sample, there may be a slight over-representation of younger, lower-income and/or Hispanic adults. Expansion of the random digit-dial methodology to include cellular telephone users broadened the reach of the survey to those who do not have landline access, and improved representativeness. Similar methods have since been implemented at the national level⁽¹⁶⁾.

Improvements in meeting both Na and K daily intake recommendations could decrease risk of CVD, the leading cause of death. Na:K is a potential surveillance tool that can assess dietary changes over time in a single marker and can identify populations at high risk for nutrition-related chronic disease.

Acknowledgements

Financial support: The study was made possible by funding from the Robert Wood Johnson Foundation, the New York State Health Foundation, the National Association of County & City Health Officials and the Centers for Disease Control and Prevention (grant number 5U38HM000449-02), the W.K. Kellogg Foundation, the US Department of Health and Human Services, and New York City tax levy dollars. This funding is administered by the Fund for Public Health in New York, a private non-profit organization that supports innovative initiatives of the NYC Department of Health and Mental Hygiene. The funders had no role in the design, analysis or writing of this article. **Conflict of interest:** None. **Authorship:** S.S.Y. formulated the study question, conducted the literature search and all analyses. S.S.Y., S.Y.A. and C.J.C. were

involved in the study design and implementation. All authors contributed to editing and finalizing the text of the manuscript. *Ethics of human subject participation:* The Institutional Review Board of the NYC Health Department approved the study.

References

1. Appel LJ, Brands MW, Daniels SR *et al.* (2006) Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension* **47**, 296–308.
2. Cook NR, Obarzanek E, Cutler JA *et al.* (2009) Joint effects of sodium and potassium intake on subsequent cardiovascular disease: the Trials of Hypertension Prevention follow-up study. *Arch Intern Med* **169**, 32–40.
3. Cogswell ME, Zhang Z, Carriquiry AL *et al.* (2012) Sodium and potassium intakes among US adults: NHANES 2003–2008. *Am J Clin Nutr* **96**, 647–657.
4. Drewnowski A, Maillot M & Rehm C (2012) Reducing the sodium–potassium ratio in the US diet: a challenge for public health. *Am J Clin Nutr* **96**, 439–444.
5. Hedayati SS, Minhajuddin AT, Ijaz A *et al.* (2012) Association of urinary sodium/potassium ratio with blood pressure: sex and racial differences. *Clin J Am Soc Nephrol* **7**, 315–322.
6. Yang Q, Liu T, Kuklina EV *et al.* (2011) Sodium and potassium intake and mortality among US adults: prospective data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* **171**, 1183–1191.
7. Institute of Medicine (2010) *Strategies to Reduce Sodium Intake in the United States*. Washington, DC: The National Academies Press.
8. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Disease. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series no. 916*. Geneva: WHO; available at http://whqlibdoc.who.int/trs/WHO_TRS_916.pdf
9. Sanderson M, Yi S, Bartley K *et al.* (2012) *The Community Health Survey, Heart Follow-Up Study Methodology Report*. New York: The New York City Department of Health and Mental Hygiene; available at <http://www.nyc.gov/html/doh/html/diseases/hfus.shtml>
10. The New York City Department of Health and Mental Hygiene (2010) The New York City Community Health Survey. <http://www.nyc.gov/html/doh/html/survey/survey.shtml> (accessed April 2012).
11. Elliott P & Stamler R (1988) Manual of operations for 'INTER-SALT', an international cooperative study on the relation of sodium and potassium to blood pressure. *Control Clin Trials* **9**, 2 Suppl., 1S–117S.
12. The New York City Department of Health and Mental Hygiene (2014) The Community Health Survey Heart Follow-Up Study (HFUS) Clinical Protocol. <http://www.nyc.gov/html/doh/html/diseases/hfus.shtml> (accessed March 2014).
13. Biro FM & Wien M (2010) Childhood obesity and adult morbidities. *Am J Clin Nutr* **91**, issue 5, 1499S–1505S.
14. Institute of Medicine (2004) *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride and Sulfate*. Washington, DC: The National Academies Press.
15. Angell SY, Yi S, Eisenhower D *et al.* (2014) Sodium intake in a cross-sectional, representative sample of New York City adults. *Am J Public Health* (Epublication ahead of print version).
16. Hu SS, Balluz L, Battaglia MP *et al.* (2011) Improving public health surveillance using a dual-frame survey of landline and cell phone numbers. *Am J Epidemiol* **173**, 703–711.