Index

abstract thinking. See cognition, abstract thinking. abulia, 6 abuse, 226, 227 acetylcholinesterase inhibitors, donepezil, 33-36, 43-44, 96, 112, 132-33, 179-81, 187 galantamine, 10, 220 rivastigmine, 10, 39-40, 42, 65, 125, 155, 164, 173 activities of daily living (ADL), 24-26, 173, 175, 178, 184-85, 186, 215-18 bathing, 32-34, 52-53, 59, 79, 81, 83, 133, 135 dressing, 24, 28, 30, 66, 71, 179, 183, 196 mobility, 7, 66, 68, 123, 179-81, 183, 185, 202, 203 toileting, 7, 9, 13, 120, 123, 126, 198, 202 See also incontinence. transferring, 66, 179, 202, 203 wayfinding, 91, 114, 115, 143-44, 159, 183 See also instrumental activities of daily living. adjustment disorder, 36, 46, 48, 73, 166, 239 advance care planning. See care planning. aggression, 29, 203-4, 213, 238 agitation, 10, 29, 229, 238 alcohol use, 7, 188, 189, 212, 242 alexia, 6 allopurinol, 51, 57, 124 Alzheimer's disease aetiology, 6 age of onset, 113 comorbidities. See comorbidities.

diagnosis, 8, 15, 32 cognitive tests, 46, 48, 55 See also cognitive function. differential diagnosis, 50, 112, 153, 182, 197, 213 early detection, 13 imaging, 100 symptom presentation, 25, 34, 43, 59, 106, 118-19 neuropathology, 6, 55 amyloid plaques, 6, 32, 54, 55, 69, 71, 112, 114 frontal lobe, 67, 110, 164, 171, 191 hippocampal sclerosis, 32, 94, 110 prevalence, 5 progression, 8, 9, 12-13, 63, 115 severe stage, 34, 36 risk factors, 98, 115 cardiovascular, 23, 114 cerebral, 42 Alzheimer's Disease International (ADI), 1, 10, 220, 237 ambulation. See mobility. amlodipine besylate, 57, 102, 108, 162, 164, 169, 173, 174 amyloid cascade hypothesis, 6 amyloid plaques, 6, 137, 144, 153, 170, 177 cerebral load, 48, 63, 69, 71, 73, 131, 137, 144 cognitive reserve against, 54, 55, 69 deposition, 170 hippocampal sclerosis and, 32, 94 imaging for, 114 anaemia, 84 antidepressants, 77, 164, 187, 194 antipsychotics, 30, 32, 34, 61, 63, 205, 237, 238 anxiety, 40, 42, 46, 90, 135, 136, 166, 187

apraxia, 81, 83, 127, 191, 209-14, 226, 234 aspirin, 98, 110, 138, 164, 169, 173, 194 asthma, 79-80, 106 atrial fibrillation, 45-46, 56, 94, 96-126, 130, 132, 162 attention deficit, 6, 24, 36, 75, 96-97, 113-14, 186, 201 - 2Barthel Index, 24, 25-26, 36-38, 43, 50-53, 68-70, 73-75, 81-83 basal ganglia, 6, 110, 173 bathing, 32-34, 52-53, 59, 79, 81, 83, 133, 135 bed/chair transfers, 66, 179, 202, 203 behavioural dysfunction, 66-67, 135, 189, 191, 196-98, 230, 231, 238-39 aggression, 29, 203-4, 213, 238 agitation, 10, 29, 229, 238 disinhibition, 6, 214, 238 hoarding, 7, 197, 239 behavioural dysfunction, overeating, 29, 66-67, 120, 178–79, 213, 214 biopsychosocial model, 11, 149, 218, 239 bladder incontinence. See urinary incontinence. blood assays complete blood picture, 45, 47, 92, 96, 108, 171, 173 ESR (erythrocyte sedimentation rate), 41, 60, 92, 96, 110, 138, 149, 216 TSH (thyroid-stimulating hormone), 45, 96, 149, 150, 152, 154, 156, 173 blood pressure, 72, 136, 138, 166 bowel control, 73, 159, 179, 196, 202

250

251

bradycardia, 94, 195-96 bradykinesia, 6, 50, 199, 213 brain imaging. See imaging. brain insulin resistance, 115 brain reserve, 170 See cognitive reserve. bronchiectasis, 31 Calcichew, 49, 108, 138 cancer, 104, 162, 174 care homes, 123, 226 care planning, 10-14, 55, 57, 83, 106, 119, 225-28 Certified Dementia Care Planner, 10, 14 integrated care, 10, 13 carers, 4, 23, 30, 108, 111, 112, 176, 229-31, 234-36, 246-47 assessment by, 71, 75, 123, 129 community care and, 4 delusions involving, 34, 110 domestic assistance, 34, 77, 170, 172-73, 180, 188, 194, 207-8 family members as, 50-52, 54, 77, 226, 230, 231-33, 238-41 preparation for deterioration, 44, 191 primary care and, 15, 23, 24, 86 prompting by, 34, 88 reports by, 23, 27, 71, 191, 236 skills and education, 30, 50, 52, 82-83, 86, 104, 111, 203 communication skills, 34, 36, 44, 117, 137, 172, 233 preparation for deterioration, 44, 191 specialist care and, 4, 13 well-being and support of, 2, 10-12, 199, 209 stress, 102, 170, 189, 204, 210-12, 225, 229-30, 231 care services and resources, 237, 240 community centres, 4, 38, 100, 132, 142, 162 day care, 11, 52, 133, 168, 198, 199

social support networks, 57, 59, 73, 166, 215, 229 social workers, 4-5, 13, 46, 79, 81, 117-19, 131, 186-88 technology location-tracking devices, 26, 30, 38, 48, 65, 79, 97, 100 carotid bruit, 84, 130, 132, 162, 171, 197 cataract, 25, 66, 68, 109, 110, 145, 149, 156 CBD. See cortobasical degeneration. CBP. See complete blood picture. CDR. See Clinical Dementia Rating. cerebral atrophy, 52, 82, 88, 106, 108, 110, 124, 127 cerebral infarction, 187 cerebrovascular accident, 40-42, 126, 169-70, 173, 179-80, 182, 202 cervical myelopathy, 124 chlordiazepoxide, 169 cholinesterase inhibitors, 23, 63, 77, 92-93, 94, 127, 156, 184 chronic illnesses, 64, 80, 236 circadian cycle. 26, 48 See also sleep. Clinical Dementia Rating (CDR), 8, 47–72, 75-76, 96-97, 139-41, 217, 219 mild dementia, 49, 118 moderate dementia, 53, 58, 60, 62, 118, 150, 188, 200 questionable dementia, 45 Clock Drawing Test, 24 clock copying section, 61, 63, 65, 70, 71–73, 161, 163, 165 not completed, 24, 65, 77, 79, 80, 148, 150 executive element impairment, 24, 25-26, 30-31, 34, 36, 61, 63, 65 no impairment, 24, 29, 47, 48, 139, 163, 172, 180 interlocking pentagon task and, 92, 100 sensitivity, 38

significant impairment, 28, 32, 42, 56, 58, 61, 81, 93 slight impairment, 24, 28, 50, 84, 123, 137, 139 time denotation, 36, 38, 40, 91, 93, 139-63, 165 clopidogrel, 134 clothing, 28, 30, 49, 51, 63-64, 66, 120-21, 126 See also dressing. cognitive function abstract thinking, 9, 63, 71, 145, 146, 183, 186, 191 ADL function and dissociation, 59 attention deficit, 6, 24, 36, 75, 96–97, 113–14, 186, 201 - 2cognitive frailty, 75, 112, 146, 153, 175, 192, 194-95, 201 deterioration, 34, 36, 55, 73, 78, 92, 201, 206 executive function, 6, 7, 22-24, 63-64, 108-9, 145-46, 166-68, 170 naming, 22, 145, 146, 183, 186, 191 screening tests, 7, 36, 46, 52, 55, 209, 212, 216-18 cognitive reserve, 54, 55, 114, 115, 119, 131, 210, 215 cognitive stimulation therapy (CST), 10, 160, 233, 237 - 38cognitive tests. See cognitive function, screening tests. colchicine, 57 community care, 75-77, 79, 81, 99, 101-2, 133, 240-42, 247 community centres, 4, 38, 100, 132, 142, 162 community engagement (of patient), 9, 49-50, 76, 101, 122, 141, 167, 168 comorbidities, 6, 12, 53-54, 178, 239 anaemia, 83, 84 cardiovascular conditions, 39 chronic illnesses, 80, 236 depression, 4, 46, 61, 89 urinary tract infection, 96

252

comorbidities (cont.) vascular dementia, 46, 57 See also mixed dementia. complete blood picture, 45, 47, 92, 96, 108, 171, 173 computed tomography. See CT. conduction dysphasia, 182 Cornell Scale for Depression in Dementia, 55 corticobasal degeneration, 126-27, 213-14 counselling, 14, 36, 46, 227, 232 CT (computed tomography imaging), 41-42, 59, 60, 88, 177, 182, 190-91, 215-16 daily living activities. See activities of daily living. day care, 48, 50, 52, 71, 72-73, 74, 132-33, 167-68 deafness. See hearing impairment. delayed recall, 34, 38, 42-44, 105-7, 135, 139-41, 149-50, 174-75 delirium, 147, 171, 192, 209, 213, 239 delusions, 31-35, 60-63, 89-91, 107-14, 116-18, 153-54, 169 - 72defined, 212 depression, 45-47, 58-170, 184 - 88assessments in case studies, 45, 46, 57, 76, 89-91 carer stress and, 142 IADL dependency and, 42 pharmacological interventions, 77, 102, 164, 187, 194 prevalence, 219 as pseudodementia, 7 as risk factor, 92 suicidal ideation and, 101 Describe, Investigate, Create, Evaluate (DICE) model, 238 diabetes, 90-92, 98, 115, 173, 174, 177, 178, 239 dialectic model, 11

diazoxide, 174 diet, 34, 35, 59, 61, 84, 86, 113, 177 - 79digoxin, 57, 94, 96 disinhibition, 6, 214, 238 disorientation, 24, 32, 35, 57, 61, 95, 199-200, 202 distressed behaviours, 40, 196, 201, 204, 229, 230, 238 DLB. See Lewy body dementia. donepezil, 33-36, 43-44, 96, 112, 132-33, 179-81, 187 dressing, 24, 28, 30, 66, 71, 179, 183, 196 drinking. 7 See alcohol use. dyslipidaemia, 98, 106, 124, 126, 162, 164, 169, 173 early dementia, 15, 50, 70, 81, 87, 108, 177, 213-14 early detection services, 13, 209, 215, 216, 218, 231, 234, 241 EDS (Hong Kong), 13-15 early intervention services, 1, 4-13, 83, 84, 209, 215, 220 end-of-life care, 209, 228 enduring power of attorney (EPA/EPOA), 13, 191, 227 escitalopram, 135, 187, 194 esomeprazole, 115 ESR (erythrocyte sedimentation rate), 41, 60, 92, 96, 110, 138, 149, 216 executive function, 6, 7, 22-24, 63-64, 108-9, 145-46, 166-68, 170 exercise (physical), 10, 27-28, 31-32, 34, 59, 86, 231, 238 family care. See carers. famotidine, 45, 88, 171 fasting lipids profile, 45, 47-98, 124, 127, 171, 173 fasting sugar profile, 94,

FAST. See Functional Assessment Staging. flupentixol, 169 forgetfulness. See memory impairment.

frailty, 75, 110, 125, 127, 153, 157, 194, 234-36 frontal lobe pathology, 67, 110, 164, 171, 191 frontotemporal dementia (FTD), 6-7, 8, 15, 179, 190-91, 204, 208, 213-14 Functional Assessment Staging (FAST), 8, 12, 129, 151, 209, 219, 226 gait abnormalities. 6, 123, 130, 132, 162, 171 See also ambulation; mobility. galantamine, 10, 220 gaze palsy, 50 GDS. See Global Deterioration Scale. General Practitioner Assessment of Cognition. See GPCOG. glaucoma, 145, 149 Global Action Plan on the Public Health Response to Dementia 2017-2025, 2, 220 Global Deterioration Scale, 8, 217, 219 goitre, 45 gout, 23, 29, 31, 56, 74, 78, 86, 124 GPCOG Informant Interview, 211-12 grocery shopping. See shopping. hallucinations, 6, 10, 123, 125, 172, 189, 199–201, 213 See also delusions. head injury, 98-99, 212, 216 hearing impairment, 47, 138-39, 150-51, 204, 215, 242 heart disease, 60, 98, 124, 138, 156, 169 hippocampal sclerosis, 32, 94, 110 hip replacement, 25, 33, 45, 202, 239 hoarding, 7, 197, 239 hobbies, 74, 97, 99, 103, 105, 122, 143-44 home care, 3, 11-13 home modifications, 12, 100, 166

home safety, 4, 129, 166, 225 household tasks, 26, 95, 107, 129, 136, 198, 201 housekeeping, 24-26, 51, 53, 58, 60, 73, 117 hygiene environmental, 65, 68, 126, 139 personal, 6, 120, 126-31, 139, 196-97, 214, 239 hypercholesterolaemia, 115, 116, 177, 178, 180, 182, 202, 204 hyperlipidaemia, 56, 90, 188 hyperorality, 7 hyperplasia, 31, 80, 118, 124, 174, 196-97, 199, 204 hypertension, 29, 31, 66, 72-150 hypotension, 6, 195-96 hysterectomy, 88, 173 ICD-11, 8 imaging, 3, 8, 46, 100, 191, 196, 208, 216 CT (computed tomography), 41–42, 59, 60, 88, 177, 182, 190-91, 215-16 See also magnetic resonance imaging(MRI) X-ray, 3 incontinence, 198, 214 bowel control, 73, 159, 179, 196, 202 urinary, 6, 47, 73, 89, 96-97, 126, 196, 202 independence, 69, 78-79, 83-85, 110, 129, 142, 144, 146 insomnia. See sleep. Instrumental Activities of Daily Living (IADL), 218 aids and support reminders, 26, 85, 127-29, 137, 153-55, 159, 163–64, 197–98 assessment tools, 218 grocery shopping, 24-26, 36, 38, 58, 60, 64, 66, 68 laundry, 24, 26, 51, 53, 58, 60, 73, 75 meal preparation, 24, 28, 47, 49, 58, 60, 87, 193–95

medication management, 47, 104, 125, 211, 234 integrated care, 10, 13 ipratropium bromide, 80, 124 irritability. See also aggression, 29, 61, 90, 91, 123, 164, 189-91, 207 isosorbide dinitrate, 96 isosorbide mononitrate, 124, 138 language impairment, 34, 38, 54-55, 81-83, 100-1, 121-25, 190-91, 207 - 8laundry, 24, 26, 51, 53, 58, 60, 73, 75 Lewy body dementia, 6, 8, 10, 50, 125, 199, 200, 213 life events, 11-12, 233 lifestyle factors, 52, 54, 179, 242 alcohol use, 7, 188, 189, 212, 242 smoking, 7, 215, 242 living tasks. See activities of daily living (ADL). location-tracking devices, 26, 30, 38, 48, 65, 79, 97, 100 low motivation, 52, 77, 87, 92, 109, 133, 163, 172 magnetic resonance imagine (MRI), 45-46, 47, 106, 113, 140, 169, 205-6, 215 - 16meal preparation, 24, 28, 47, 49, 58, 60, 87, 193-95 mecobalamin, 84, 163 medial lateral temporal atrophy, 57, 82, 140, 171 mediastinum, 138 memantine, 55, 59, 61, 179, 181, 182, 184, 237 memory clinics, 3–5, 13, 92 memory impairment, 47, 48, 71, 72-161 Alzheimer's disease early stage, 8 carer concerns, 72-129 delayed recall, 34, 38, 42-44, 105-7, 135, 139-41, 149-50, 174-75

deterioration, 79, 117, 125, 129, 166, 168 family reported, 22-24, 59-81, 89-91 IADL and, 51 Lewy body dementia, 6 masking, 110 misplacement of belongings, 121, 139, 141, 144 patient insight, 48 meningioma, 110, 206 mental health, 147, 196, 212, 229, 239-40 anxiety, 40, 42, 46, 90, 135, 136, 147, 166, 187 See also depression. suicidal ideation, 101, 102, 189, 227 metformin, 92, 98-99, 124, 126, 162-63, 173 methotrexate, 49 methyldopa, 185-86 metoprolol succinate, 169, 173 mild cognitive impairment (MCI), 7-19, 153, 162, 164, 168, 173, 175, 177 mild dementia, 41, 47, 49, 118, 145-46, 159-60, 176-77 Mini-Mental State Examination. See MMSE. mixed dementia, 61, 88, 102, 127, 182, 203, 237 MMSE (Mini-Mental State Examination), 168, 191, 216-17, 234 interlocking pentagon task, 34, 38, 90, 92, 99, 100, 158, 160 mobility, 66, 68, 123, 179-81, 183, 185, 202, 203 moderate Alzheimer's disease, 9, 29, 70–71, 76, 78, 83, 234, 237 moderate cognitive impairment, 123 moderate dementia, 4, 10, 28, 30, 35, 40, 66 Montreal Cognitive Assessment (MoCA), 145, 146, 147, 155–56, 161-62, 168, 170, 217-18

mood change. 46-47, 56-58, 76-77, 89-90, 92-95, 99-103, 107-10, 185-86 See also depression. morphine, 138 Multispecialty Interprofessional Teams, 4 naming, 22, 145, 146, 183, 186, 191 National Dementia Strategy, 3 neurofibrillary tangles, 6 neuroimaging. See imaging. neuroleptics, 90, 172, 237 Neuropsychiatric Interview Ouestionnaire, 55 NICE (National Institute for Health and Care Excellence), 5, 215, 220, 237, 240 non-Alzheimer's dementia, 29, 31, 35, 37, 43, 45, 49, 213 - 14non-pharmacological interventions, 10-11, 12, 23-24, 30, 32, 40, 112, 237

object agnosia, 6 obstructive pulmonary disease, 80, 96, 124 occupational therapists, 4-5, 10, 13, 144 old age, extreme, 32 osteoarthritis, 23, 88, 173, 195, 204 osteoporosis, 45, 138, 156 overeating, 29, 66, 120, 178-79, 213, 214 palliative care. See end-of-life care. pancytopenia, 31 pantoprazole, 49, 94, 98, 102, 110, 126, 156, 164 paracetamol, 126, 171 Parkinson disease dementia (PDD), 10, 50, 213 parkinsonism, 6, 50, 125, 189, 198-99, 213-14 atypical, 50 bradykinesia, 6, 50, 199, 213 pepcidine, 169

personal care. 9, 72, 74, 76, 101, 167 See also activities of daily living (ADL). personality change, 6, 11, 32, 100, 133, 206-8, 229, 231 person-centred care, 2, 5, 11, 15, 40, 226, 238 PET (positron emission tomography), 6, 177 physical exercise, 10, 238 power of attorney (EPA/ EPOA), 13, 191, 227 primary care, 1-18, 62-81, 84-110, 147-75 carers and, 236 diagnostic accuracy, 15 key considerations, 232, 237 models, 3, 11 with specialist outreach, 3 prescribing, 10 resources for, 212 roles, 11-12, 15 service integration, 5, 12 specialist care and, 3 primary progressive aphasia (PPA), 8, 96, 213-14 pseudobulbar palsy, 6 pseudodementia. See depression. psychiatric disorders. 31, 33, 35, 37, 43, 45, 49, 51 See also depression; mental health. psychoeducation, 36, 46, 102, 106, 229-30 psychomotor retardation, 6 psychotherapy, 46, 92, 188 public awareness, 59, 153, 241

Quality and Outcomes Framework, 3 quetiapine, 30, 63, 125, 188–89

radiation therapy, 162 rapid eye movements behaviour disorder (REMBD), 6, 125, 172, 189, 199, 213 reminders, 26, 85, 127–29, 137, 153–55, 159, 163–64, 197–98 residential care. *See* care homes. resistance to care, 52, 238

Resources for Enhancing Alzheimer's Carer Health, 10 rigidity, 6, 104, 213, 237 risk factors, 114, 242 cardiovascular, 7, 23, 114, 115, 177, 180, 239, 242 cerebral, 42, 84 chronic disease, 92 diabetes, 42, 115 See also comorbidities. rivaroxaban, 57 rivastigmine, 10, 39-40, 42, 65, 125, 155, 164, 173 safety, 10, 30, 38, 121, 129, 209, 210, 215 salbutamol, 80, 124 screening tests. 7 See cognitive screening tests. secondary care, 4, 14-15, 102, 104, 208 selective serotonin reuptake inhibitors (SSRI), 67, 90, 102, 108, 135, 171, 197 self-awareness, 55, 73, 106, 108, 163, 166, 206 self-care, 31, 34, 35, 66-67, 69, 82-83, 126-27, 213 shopping, 24–26, 30, 36, 38, 60, 64, 66, 87 simvastatin, 45, 57, 98, 102, 124, 126, 173, 194 sleep, 6, 93, 94, 134, 137, 189, 198-99, 201 small vessel disease, 47, 84, 85-87, 88, 92, 110, 115, 184 smoking, 7, 215, 242 social centres, 11 social interaction, 35, 59, 61, 84, 113, 144, 182-83, 237 - 38social workers, 4-5, 13, 46, 79, 81, 117-19, 131, 186 - 88specialist care, 1-5, 11, 13-14, 15, 204, 206, 232, 236-37 SSRIs. See selective serotonin reuptake inhibitors. stress (carers), 86, 102, 117, 189, 204, 208, 212, 229 stroke, 6, 95-97, 127-31, 177, 179, 180-82, 203, 212-13

255

strontium ranelate, 169 subdural haematoma, 22-23, 98, 169, 216 subjective cognitive impairment, 7, 141, 155 suicidal ideation, 101-2, 189, 227 support services. See care and support services and resources. swallowing problems, 13, 66, 214 Swedish Dementia Registry, 3 task-sharing, 1, 14–15 task-shifting, 1, 2, 14-15, 16 temper. See irritability. temporal lobe degeneration, 57, 82-83, 110, 124 thalassemia, 197 thiamine, 189 three-step command task, 24, 30, 32, 42, 44, 51, 53, 56 thyroid function, 8

thyroid-stimulating hormone, 45, 96, 149, 150, 152, 154, 156, 173 toileting, 7, 9, 13, 120, 123, 126, 198, 202 See also incontinence. transferring, 66, 179, 202, 203 tremor, 50, 124, 213 triage, 13, 81, 153, 156, 212, 218, 219, 234 United Kingdom, 3, 5, 209, 215, 236 urinary incontinence, 47, 73, 96-97, 126, 142, 159, 196, 202 utilisation behaviours, 6

- valproate, 171, 197, 205 vascular dementia, 42, 96, 102, 106, 177–79, 181, 182, 184 verbal expression, 24, 26, 153,
 - 155, 183–84, 191, 203, 204 *See also* word finding.

- verbal function, naming, 22, 145, 146, 183, 186, 191 vildagliptin, 98, 173 visual hallucinations, 6, 50, 66, 160, 198–201, 213 visual impairment, 32, 154 *See also* cataract; glaucoma. waiting times, 2, 5, 13–15 warfarin, 45, 96, 126 wayfinding, 91, 114, 115, 143–44, 159, 183 word finding, 23, 90–91, 159, 50
- 109–10, 158–60, 180–82, 183, 190–91, 200–2 See also verbal expression; naming; GPCOG informant interview. World Alzheimer's Report, 1, 2, 13
- X-ray, 127, 134, 138, 140, 158, 160, 166

https://doi.org/10.1017/9781108989336.009 Published online by Cambridge University Press