Article: 1286

Topic: EPV07 - e-Poster 07: Cultural Psychiatry

What Can We Learn About Psycho-social Functioning of Individuals Diagnosed with Schizophrenia From Standardized Patients?

L. Lucic<sup>1</sup>, A. Khan<sup>2</sup>, L. Liharska<sup>3</sup>, B. Rothman<sup>3</sup>

<sup>1</sup>Social Science and Cultural Studies, Pratt Institute, Brooklyn, USA; <sup>2</sup>Psychopharmacology Research Program, Nathan S. Kline Institute for Psychiatric Research, New York, USA; <sup>3</sup>Research Department, ProPhase LLC, New York, USA

Treatment of patients with schizophrenia can be improved by precise administration of rating scales. The Positive and Negative Syndrome Scale (PANSS) is a commonly used tool to assess symptoms of schizophrenia. In order to develop a participatory learning program that aims to enhance raters training and increase inter-rater reliability we trained five actors to serve as standardized patients (SP). A training program for SPs was constructed using a narrative analysis scheme in the following manner: 1) Seven transcripts of SCI-PANSS interviews were analyzed and subdivided into thematic chapters including: symptom frequency, intensity, complexity, interference and impairment; 2) Script/story analysis of transcribed interviews were conducted where we identified scripted elements common among all seven interviews and storied elements that deviate from the script and have a function to individuate each of the interviews; 3) Linguistic markers (words, phrases, or discursive patterns indicative of symptoms and their severity) that are often seen as hallmarks of aetiology of schizophrenia were identified; 4) Symptoms were divided into two severity levels, moderate and severe, and three full SCI-PANSS interview scripts were built; 5) A workshop for five professional actors was conducted under the supervision of an expert PANSS interviewer. By studying the development of this program knowledge can be gained about a number of aspects related overall psycho-social functioning of individuals diagnosed with schizophrenia, such as the impact of semantic and discursive practices on interpersonal interactions and their further effect on the progress of the illness.