

Mario Milkov¹, Stefan Mirchev², Aleksandar Valkov², Georgi Nikolov³, Boris Duhlenki³, Hamaputra Vijayendra⁴

¹Prof. Paraskev Stoyanov Medical University of Varna Faculty of Dental Medicine, ²Medical University, Pleven, Bulgaria, ³Medical University of Pleven, Bulgaria, ⁴Vijaya E.N.T. care Centre, Belgaum, India

Learning Objectives: Although relatively rare in childhood, cholesteatoma deserves a special attention by not only by general practitioners but also by specialists in different mutually related disciplines in the family of neurosciences and, particular, in neurootology.

Introduction. Cholesteatoma in childhood is difficult to diagnose in spite of substantial advances in diagnostic approaches. The delayed detection of this complicated pathology necessitates adequate surgical management.

Methods: We report a child aged eight years presenting with congenital cholesteatoma accompanied by peripheral facial nerve palsy. Management has been initiated by a neurologist and later on, an otorhinolaryngologist has been involved in the therapeutic team because of the failed drug therapy. According to parents' report, three years ago the child complained of gait disturbances attributed mainly to his overweight. A comprehensive physician's examination included preoperative audiometry and computer tomography, intraoperative monitoring system enabling safe cholesteatoma removal in the area of bone destruction at second genu level as well as pre- and postoperative electromyography.

Results: We successfully identified the degree of morphological damage and of the disturbed function. Besides, the prognosis concerning facial nerve restoration was clarified. On the seventeenth day after the surgical intervention, there was convincing electromyographic evidence of significant improvement of the facial nerve function and positive change of facial appearance as documented by pictures in 4-day intervals. A concise review of recent publications dealing with cholesteatoma diagnosis and surgery illustrated the socio-medical importance of this entity in childhood (e.g., A. T. Harris et al., *J Laryngol Otol.* 2016;130:235; M. S. Cohen et al., *Laryngoscope.* 2016;126:732; J. B. Hunter et al., *Otolaryngol Head Neck Surg.* 2016; Mar 1, etc.).

Conclusion: Scanty initial clinical symptoms along with poor otoscopic findings hamper considerably the early exact diagnosis of cholesteatoma. Interdisciplinary collaboration between neurologists, otorhinolaryngologists, radiologists and neurosurgeons could warrant the proper therapeutic behaviour in children with cholesteatoma.

doi:10.1017/S0022215116006319

ID: IP135

Endoscope-i: transforming endoscopic technology and the delivery of patient care in ENT

Presenting Author: **Nina Mistry**

Nina Mistry¹, Jeyanthi Kulasegarah², Ajith George³, Christopher Coulson²

¹University Hospitals Birmingham NHS Foundation Trust, ²University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK, ³University Hospitals of North Midlands NHS Trust, Stoke, UK

Learning Objectives: To raise awareness amongst ENT surgeons of the potential role that digital and mobile device technology such as the endoscope-i can play in enhancing the delivery of healthcare services to patients.

Introduction: Digital and mobile device innovation in healthcare is a growing market. The introduction of the endoscope-i, the world's first endoscopic mobile imaging system, allows the acquisition of high definition images and videos, without the need for expensive and bulky stack systems.

Methods: The endoscope-i system was first developed by the senior authors in conjunction with experts in the engineering and software development fields. The system combines the e-i Pro camera app with a bespoke engineered endoscope-i adaptor which fits securely onto the iPhone. The system is portable and allows high definition endoscopic imaging of the ear, nose and throat, with the facility to capture still images and videos.

Results: The endoscope-i has far-reaching applications but comes into its own in when examining the ear endoscopically. The assessment of a variety of pathology such as tympanic membrane perforations, retraction pockets and cholesteatoma can be undertaken efficiently and relayed immediately to the patient. Not only does this provide instant feedback to the patient during the consultation, thereby improving patient education, it also allows the documentation of findings via a secure app which can be stored as part of the patient's record of care, replacing the need for previous hand-drawn notes.

Conclusions: The use of endoscopic mobile imaging systems has the potential to transform the way that healthcare is delivered in ENT. The endoscope-i system provides a cost effective device that is accessible, easy to use and which makes diagnostics simpler and quicker. Although there are clear applications in the field of otolaryngology, the technology is also being used by other medical specialties such as anaesthetics and urology as well as in veterinary medicine and engineering.

doi:10.1017/S0022215116006320

ID: IP136

The Novel Affordable Telford Temporal Bone Holder

Presenting Author: **Alistair Mitchell-Innes**