

Intake and food sources of nitrites and *N*-nitrosodimethylamine in Spain

Paula Jakszyn^{1,*}, Antonio Agudo¹, Antonio Berenguer¹, Raquel Ibáñez¹, Pilar Amiano², Guillem Pera¹, Eva Ardanaz³, Aurelio Barricarte³, María Dolores Chirlaque⁴, Miren Dorronsoro², Nerea Larrañaga², Carmen Martínez⁵, Carmen Navarro⁴, José R Quirós⁶, María José Sánchez⁵, María José Tormo⁴ and Carlos A González¹

¹Institut Català d'Oncologia (ICO-IDIBELL), L'Hospitalet de Llobregat, 08907 Barcelona, Spain: ²Dirección de Salud de Guipúzcoa, 20013 San Sebastián, Spain: ³Departamento de Salud de Navarra, 31003 Pamplona, Spain: ⁴Consejería de Sanidad y Consumo, 3008 Murcia, Spain: ⁵Escuela Andaluza de Salud Pública, 18080 Granada, Spain: ⁶Consejería de Sanidad y Servicios Sociales de Asturias, 33001 Oviedo, Spain

Submitted 27 January 2005: Accepted 27 July 2005

Abstract

Objective: To conduct a comprehensive assessment of dietary intakes of nitrites and *N*-nitrosodimethylamine (NDMA).

Subjects and setting: A study was conducted within the Spanish cohort of the European Prospective Investigation in Cancer and Nutrition (EPIC) to assess the intake and food sources of these compounds in Spanish adults. The study included 41 446 health volunteers, aged 29–69 years, from Northern and Southern regions. Usual food intake was estimated by in-person interviews using a computerised dietary questionnaire.

Results: The estimated geometric mean was 0.994 mg day⁻¹ for nitrites and 0.114 µg day⁻¹ for NDMA. For both compounds a positive trend in consumption with increasing energy intake was observed. Dietary NDMA was related to age and sex after energy adjustment, while nitrite consumption increased with higher intakes of vitamin C ($P < 0.001$). The food groups that contributed most to intakes were meat products, cereals, vegetables and fruits for nitrites, and processed meat, beer, cheese and broiled fish for NDMA. Current and past smokers, who had high levels of NDMA from tobacco exposure, were also identified as the highest consumers of dietary NDMA. Furthermore, smokers had low intakes of vitamin C (an inhibitor of endogenous nitrosation).

Conclusions: Intake levels of NDMA and nitrites in a Mediterranean cohort are currently relatively lower than those previously reported, although processed meat, beer and cured cheese still are the most important contributors to NDMA intake.

Keywords

Nitrites
Nitrosamines
Dietary intake
N-Nitrosodimethylamine
Cancer risk

Humans are exposed to *N*-nitroso compounds (NOC) from diet, tobacco smoke and other environmental sources, as well as from endogenous synthesis¹. Diet has been identified as the major source of exogenous exposure². Of the various NOC, nitrosamines (NA) have been studied the most extensively in foods. These compounds are formed through the reaction of a nitrosating agent (nitrogen oxides) with amino compounds. Nitrogen oxides are derived from two main sources: (1) the addition of nitrate and/or nitrite to foods and (2) the process of heating and/or drying of foods using combustion gas, during which molecular nitrogen can be oxidised. The highest concentrations of NA are found in cured meat and cheese, smoked preserved foods, foods preserved by drying, pickling and salting, and foods stored under humid

conditions³. The major NA in the diet are *N*-nitrosodimethylamine (NDMA), *N*-nitrosopyrrolidine and *N*-nitrosopiperidine.

In vitro and *in vivo* studies have shown that nitrates can be reduced to nitrites by bacterial or mammalian metabolic pathways. Thus, both nitrites and nitrates reduced to nitrites may react endogenously with amines and amides derived from proteins to form carcinogenic NOC³. Several studies have reported that endogenous synthesis could contribute 45–75% of total human exposure⁴. This process could be modified by several factors; for instance, vitamin C has been identified as an effective inhibitor of endogenous synthesis⁵. Nitrites are widely used as food preservatives and colouring agents by the food industry. They are mainly found in cured meat

*Corresponding author: Email paujak@iconcologia.catsalut.net

products, fish and cheeses. Nitrites may also be found in cereal products and vegetables as a result of formation during storage and/or cooking. Although nitrates can be reduced to nitrites, there is not much evidence about their relation to cancer risk.

Various NOC have been found to be carcinogenic in multiple organs in at least 40 animal species, including higher primates⁶. Many studies have reported that a high intake of NA may increase the incidence of nasopharyngeal, oesophageal, pancreatic, gastric, brain and colorectal cancers^{7,8}; however, their health effects in humans still have not been clearly established. NDMA and *N*-nitrosodiethylamine are classified by the International Agency for Research on Cancer as probable carcinogens for humans⁹.

In the majority of studies carried out to date, the estimation of dietary exposure to NA and NA precursors has focused only on those foods in which the content is high. The advantage of an assessment with a complete list of food items is that all dietary sources of the compound of interest are included. Foods with modest content consumed frequently by a substantial part of the population can be as important as infrequently consumed foods with higher content. In summary, more comprehensive assessment of intake from a wide array of potential dietary sources is needed in order to evaluate the health effects of these compounds in epidemiological research¹⁰.

Recently, we have compiled the information available on the content of nitrates, nitrites and NA in foods¹¹. In the present study we used the data to estimate the dietary intake of nitrites and NDMA in the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC). We also explored associations between intakes and several variables that might affect NDMA exposure.

Material and methods

Study population

Dietary data and other lifestyle factors from 25 812 women and 15 634 men aged 29–69 years, participants of the EPIC–Spain cohort, were obtained between October 1992 and July 1996. Participants were healthy volunteers recruited in five Spanish regions: three from the North (Asturias, Navarra and Guipúzcoa) and two from the South (Murcia and Granada). Most of the selected subjects were active blood donors. The mean age at recruitment was 50.8 and 48.4 years for men and women, respectively. The final participation rate was 55–60% across centres.

Dietary information

Usual food intake during the preceding year, taking into account seasonal variations, was estimated by an in-person interview using a computerised dietary history questionnaire which was developed and validated

specifically for the EPIC study in Spain¹². The questionnaire was structured according to occasions of food intake. Trained interviewers gathered data on preparation methods, average frequency of consumption, and usual portion sizes for each food consumed at least twice per month (or once per month for seasonal foods). Portion sizes were reported in natural units, household measures or with the aid of 35 sets of photographs prepared specifically for the study. The questionnaire included a list of 600 foods and beverages, and about 150 regional recipes. Data on vitamin C in foods were obtained from a computerised database of more than 700 food items compiled for the EPIC study in Spain¹³. We took only food sources into account because less than 4% of participants in the study took mineral or vitamin supplements during the previous week¹⁴.

Estimation of nitrite and NDMA intakes

Dietary intakes of nitrites and NDMA were estimated based on information compiled recently in a food database (FDB)¹¹. The FDB includes information about 278 and 313 food items for nitrites and NA, respectively. Since considerably more information is available for NDMA than for other NA found in foods, we decided to estimate only dietary intakes of NDMA. Since the food item could have several different source of information, we selected Spanish or European values when they were available. For food items for which content was not available, the value of the nearest comparable food was assigned. When we found several matches for a food item we assigned the mean of all the suitable values (<5% of the food items).

Statistical analysis

Means for nitrites and NDMA are reported as mg day^{-1} and $\mu\text{g day}^{-1}$, respectively. As nitrite and NDMA distributions were strongly right-skewed, we used the natural logarithm (\log_e) transformation to improve normality. Arithmetic means for the log-transformed intakes were later exponentiated to obtain the geometric means. Analysis of variance was used to assess differences in nitrite and NDMA intakes across categories of different variables, adjusting for sex, age, energy intake, centre and body mass index (BMI)¹⁵. The average estimates of dietary intakes of nitrites and NDMA for the whole cohort were age- and sex-standardised based on the structure of the Spanish population (aged 35–64 years)¹⁶.

In order to identify food sources of these compounds in our population, the individual contribution of each food item to the total compound intake was computed. Additionally, to identify which foods better explain the variability in intakes of NDMA, foods were ranked based on the proportion of variance in NDMA they explained (R^2), using linear regression analysis with stepwise selection¹⁵. Data were analysed using software from the R Foundation for Statistical Computing¹⁷.

Results

The number of food items from the diet history linked to the FDB was 149 for nitrites and 257 for NDMA. The estimated geometric mean was $0.994 \text{ mg day}^{-1}$ for nitrites and $0.114 \mu\text{g day}^{-1}$ for NDMA (age- and sex-standardised based on the Spanish population structure)¹⁶.

Geometric means of both compounds were computed separately for categories of sex, age, centre, smoking, BMI, energy intake and vitamin C intake (Table 1). Means were adjusted by sex, age, BMI, centre and energy consumption. For nitrites, we found a positive trend across categories of energy intake. The group with the highest energy intake had 46% more nitrites than the group with the lowest energy intake. When comparing intakes of nitrites in relation to intakes of vitamin C, we observed a positive gradient (Fig. 1). We found a 27% difference between the highest and the lowest quartiles of vitamin C consumption. Differences in the intake of nitrites across

categories of sex, age, centre, BMI and smoking did not follow a significant pattern.

NDMA intakes were 20% higher in men ($0.118 \mu\text{g day}^{-1}$) than in women ($0.098 \mu\text{g day}^{-1}$). Geometric mean NDMA intakes tended to decline with increasing age. The lowest values were observed in older people while the highest were found in young individuals (0.085 vs. $0.117 \mu\text{g day}^{-1}$). When comparing NDMA intake across levels of energy intake, the difference between the highest and the lowest energy intake group was 79%. There were no meaningful differences in NDMA intake across categories of centre, BMI, tobacco use and vitamin C.

The association between vitamin C consumption and NDMA intake was also examined after stratifying based on categories of tobacco use, as shown in Fig. 2. Current and past smokers in the bottom two quintiles of vitamin C consumption had substantially higher levels of NDMA than smokers in the top two quintiles of vitamin C intake.

Table 1 Intakes of nitrites and *N*-nitrosodimethylamine (NDMA) in categories of different variables

	<i>n</i>	Nitrites (mg day^{-1})		NDMA ($\mu\text{g day}^{-1}$)	
		Geometric mean*†	Proportion of change‡	Geometric mean*†	Proportion of change‡
Sex§					
Male	15 579	0.96	1.00	0.12	1.00
Female	25 752	0.96	0.99	0.10	0.80
Age (years)§					
28–34	300	0.97	1.00	0.12	1.00
35–44	13 808	0.97	0.99	0.12	0.98
45–54	16 075	0.96	0.98	0.11	0.89
55–64	10 693	0.95	0.97	0.09	0.80
65–74	455	0.94	0.96	0.09	0.72
Energy intake (kcal day^{-1})§					
Q1 (323–1750)	10 321	0.79	1.00	0.08	1.00
Q2 (1750–2180)	10 331	0.93	1.17	0.10	1.26
Q3 (2180–3270)	10 338	1.00	1.26	0.12	1.44
Q4 (2730–3102)	10 341	1.15	1.46	0.14	1.79
Centre§					
North Spain	25 109	1.01	1.00	0.10	1.00
South Spain	16 307	0.91	0.90	0.11	1.05
Body mass index (kg m^{-2})§					
< 25	9195	0.96	1.00	0.11	1.00
25–30	19 674	0.96	1.00	0.11	0.99
> 30	12 462	0.96	1.00	0.104	0.98
Vitamin C intake (mg day^{-1})§					
Q1 (1.45–94)	10 341	0.85	1.00	0.11	1.00
Q2 (94–137)	10 325	0.94	1.10	0.11	1.00
Q3 (137–191)	10 335	0.99	1.16	0.10	0.97
Q4 (191–1263)	10 330	1.08	1.27	0.10	0.95
Smoking§					
Never	22 927	0.96	1.00	0.10	1.00
Past	7285	0.97	1.01	0.11	1.05
Current	10 109	0.92	0.95	0.11	1.04

Q – quartile.

* Adjusted for sex, age, energy intake, body mass index (BMI) and centre.

† The values reported are calculated over the number of subjects with valid information; the numbers of subjects with missing information were as follows: sex, age, energy intake, BMI and vitamin C (115), centre (30) and tobacco smoking (1125).

‡ This value shows the proportional change taking the first category as reference.

§ In all cases *P*-values were <0.001, except for nitrites in categories of BMI.




Fig. 1 Intake* of nitrites by vitamin C decile (d). *Geometric means adjusted by age, sex, centre, energy intake and body mass index

Among never smokers, there was no association between NDMA and vitamin C.

The food groups that contributed most to intakes of each compound were also identified (Table 2). For NDMA, we identified the following food items: cured meats (14%), cheese (13%), frankfurters (12%), beer (11%), bacon (11%), broiled fish (5%) and fresh sausage (4%). Overall, processed meat contributed more than 40% to total NDMA intakes. For NDMA we also analysed the contribution of foods to the between-person variance. The major contributors to the variance were frankfurters, beer, bacon and cured meats ($R^2 = 0.71$) (data not shown). The food items selected by the model were the same as the




Fig. 2 Intake* of *N*-nitrosodimethylamine (NDMA) by vitamin C quintile (q) in past, current and never smokers. *Geometric means adjusted by age, sex, centre, energy intake and body mass index

main contributors to total intake, but their relative contribution in terms of explained variance or slopes in intake was not the same. The food items that contributed most to nitrite intakes were meat products (boiled ham, bacon, sausage, hamburger), cereals (bread, spaghetti), vegetables (potato, lettuce, leek, tomato, cucumber) and fruits (orange, apple, pear, peach) (Table 2).

Discussion

In the present study we estimated dietary intakes and food sources of nitrites and NDMA in Spanish adults. The geometric mean was $0.994 \text{ mg day}^{-1}$ and $0.114 \text{ µg day}^{-1}$ for nitrites and NDMA, respectively.

Analysis of the NDMA distribution suggests that dietary intakes are 20% higher for men than for women. Several other studies have also shown that intake in men could be up three times higher^{18–20}. A possible explanation for sex differences in this study is that men consumed significantly more beer and processed meat than women, given that these foods are strongly related to NDMA intake.

The relationship between NDMA and age was negative, as intakes in younger people were higher than in older individuals. The difference between the oldest and youngest individual was 28%. This pattern was attributable to the higher intakes of processed meat and beer among younger subjects.

For both nitrites and NDMA a positive association with energy intake was observed. This could be a consequence of the substantial contribution of key food sources of nitrites and NDMA to total energy intakes. This link with energy intake was stronger for NDMA than for nitrites, probably because NDMA is concentrated in food groups with high energy density (kcal g^{-1}) such as processed meat and beer, while the key sources of nitrites include foods with low energy density, such as vegetables and fruits.

Patterns of intake across other variables analysed were not meaningful in spite of their statistical significance, which is probably due to the large sample size.

Figure 1 shows a direct relationship between intake of nitrites and consumption of vitamin C. Ascorbic acid has been identified as an important factor in the inhibition of NA formation because it reduces nitrites to nitric oxide⁵. In our study, individuals who had higher intakes of nitrites also consumed higher levels of vitamin C, which can help to inhibit their transformation to NA. In order to identify groups of individuals who have higher overall exposure to NA, we explored the relationship between vitamin C consumption and NDMA intakes across categories of tobacco use. These results showed that, among current and past smokers (also exposed to NA from tobacco) in our population, those with low intakes of vitamin C also had high dietary intakes of NDMA. These low intakes of vitamin C in this group could exacerbate the problem of endogenous formation of NA. This pattern of exposure

Table 2 Contributions of food items to total intakes of *N*-nitrosodimethylamine (NDMA) and nitrites

Food item	Contribution to total intake of NDMA		Food item	Contribution to total intake of nitrites	
	Percentage	Cumulative percentage		Percentage	Cumulative percentage
Cured meat	14.26	14.26	Boiled ham	15.08	15.08
Cured cheese	13.16	27.42	Bread	10.17	25.25
Frankfurt sausage	11.17	38.52	Hamburger	7.28	32.53
Beer	11.00	49.59	Bacon	6.70	39.23
Bacon	10.56	60.15	Spaghetti	6.50	45.73
Broiled fish	4.66	64.81	Potato	4.80	50.53
Fresh sausage	4.31	69.12	Sausage	3.87	54.40
Hamburger	2.89	72.12	Lettuce	2.81	57.21
Boiled ham	1.26	73.27	Orange	2.54	59.75
Wine	0.68	73.95	Apple	2.51	62.26
Whisky	0.11	74.06	Leek	2.16	64.42
			Cucumber	2.12	66.63
			Tomato	1.98	68.61
			Spinach	1.91	70.52

reflects the fact that, in our population, smokers had the most unhealthy dietary profile, consuming less vegetables and fruits²¹ and more processed meat and alcohol²² than non-smokers. Similar findings have been reported in other populations. For instance, in a meta-analysis carried out by Dallongeville *et al.*, smokers consumed 16.5% less vitamin C and 9% more saturated fats than non-smokers²³. Moreover, smoking itself decreases levels of vitamin C in plasma, increasing dietary requirements²⁴. This could further exacerbate total NA exposure in this population.

For NDMA our estimated intake value is within the range of 0.08 to 0.55 $\mu\text{g day}^{-1}$ found in previous studies^{25–32}. Although data suggest that mean intakes in our population are relatively low, it is important to note that 10% of the population had consumption levels above 0.40 $\mu\text{g day}^{-1}$, while 5% had intakes five-fold higher than the estimated mean (0.78 $\mu\text{g day}^{-1}$). To evaluate differences in estimated intakes across studies, several methodological issues must be taken into account. First, various dietary assessment instruments (food-frequency questionnaire, diet history, 48-hour recall or duplicate sampling) have been utilised. Second, there are large differences between the numbers of food items used to estimate total intakes of NDMA (28 to 474 items). Finally, the results reported in previous studies have been expressed as arithmetic means or medians, while in our study the values are shown as the geometric mean. In the present study, median and arithmetic mean were 0.109 and 0.19 $\mu\text{g day}^{-1}$, respectively.

Another important influence on intakes is the fact that, during the 1980s, reductions in the use of nitrates and nitrites for curing meat and modifications in malting techniques in the brewing industry resulted in significant reductions in the levels of NA in foods³. For instance, the value used in the present study for beer was 10 times lower than values used in previous studies. Since beer is an important contributor to total intakes, it is not surprising

that NDMA exposure in our population is lower than in the majority of the studies published in the past. These results show that, at the present, the contribution from dietary sources to total exposure is lower than in previous decades.

In accordance with other studies, intakes of NDMA were also strongly dependent on the consumption of processed meats, especially frankfurters, sausages, bacon and cured meats, followed by beer and cheese. Moreover, we found that broiled fish was one of 10 main contributors, with 5% of intake. This finding is relevant because despite the fact that broiled fish does not have a high content of NDMA, intakes in Mediterranean countries are very high. On the other hand, foods with a high content of NDMA¹¹, such as smoked or salted fish, are not important contributors to the total intake because they are consumed by a very low percentage of the EPIC–Spain population (0.30% and 3%, respectively). Different patterns of consumption have been found in studies published in Finland and France, in which smoked fish contributed 22–25% to total intakes of NDMA^{18,33}. The highest values of NDMA intake have been reported in Japan, due to high intakes of dried and smoked fish³⁴.

When the contribution by foods to total intake and to between-person variance of NDMA was analysed, the food items selected were similar but differed in their relative importance. In the present study, only four foods (frankfurters, beer, bacon and salami; $R^2 = 0.71$) explained 70% of the variance in NDMA consumption (data not shown). These same four foods jointly contributed 49% of total NDMA intake. As is generally found in other studies, fewer items are necessary explain a given proportion of between person-variance than the same proportion of total intakes^{35,36}.

Intakes of nitrites were comparable with previously published values^{2,18}, which ranged between 0.8 and 5.3 mg day^{-1} . For nitrites the major contributors to total intake were meat products (32%), cereals and derivatives

(22%), vegetables (16%) and fruits (12%). We found that bread is one of the most important contributors to total intake of nitrites. It is relevant because despite the fact that bread does not have a high content of nitrites, intakes in Spanish adults are very high. The present study indicates that major sources of nitrites in Spanish adults are the same as those reported in previous investigations from other countries³⁷.

One limitation of our study could be that the Spanish EPIC study is based on a non-representative sample of the general population. However, the number of subjects was very large, the participation rate was relatively high, subjects came from different social levels and different geographical areas, and the pattern of dietary intake and other lifestyle factors was very similar to that observed in population-based surveys carried out in the Spanish regions included in the present study^{38–40}. In order to minimise this limitation, the reported geometric means for the whole cohort were standardised using the age and sex distribution of the Spanish population.

The limitations of assigning values to foods based on published reports include the limited availability of data on levels of these compounds in some items, and the variation in levels found in similar items across countries. We addressed this problem by taking the most suitable value for our population taking into account the analytical method, the origin and the year of publication.

Among the strengths of the present study are the large dataset (41 446 subjects), the number of food items included for analysis and the validated methodology used in the dietary assessment¹². This is important, taking into account that even high-quality food composition tables will produce poor estimates when combined with low-quality data due to deficient validity and/or reliability of a dietary questionnaire. Several studies have reported intakes of these compounds, but in some cases the accuracy of the methods used to assess dietary intakes was limited.

In summary, we have estimated exogenous dietary exposure to nitrites and NDMA in Spanish adults (aged 35–64 years). These findings may contribute to improved ability to classify subjects according to their dietary intake patterns, and, in the future, to better understanding of the role of such compounds in cancer risk. In our study we also found that current and past smokers, groups with low intakes of vitamin C (an inhibitor of endogenous nitrosation), also had high dietary NDMA exposure. Thus, these groups could be considered a high-risk population. Additional research to estimate endogenous synthesis is needed for an accurate assessment of total exposure.

Acknowledgements

This study is part of the EPIC–Spain Study. EPIC is coordinated by the Unit of Nutrition and Cancer of the International Agency for Research on Cancer. This project has received financial support from the Health Research

Fund (FIS, 02/0652) of the Spanish Ministry of Health and the ‘Red de Centros de Investigación Cooperativa en Epidemiología y Salud Pública’ (RCESP, C03/09). Participation of P.J. in the project was supported by a grant of the ‘Fundació Privada Institut d’Investigació Biomèdica de Bellvitge’ (IDIBELL).

References

- 1 Mirvish SS. Role of *N*-nitroso compounds (NOC) and *N*-nitrosation in etiology of gastric, esophageal, nasopharyngeal and bladder cancer and contribution to cancer of known exposures to NOC. *Cancer Letters* 1995; **93**(1): 17–48.
- 2 US Assembly of Sciences, National Research Council. *The Health Effects of Nitrate, Nitrite and Nitrosocompounds. Part I of a Two-part Study by the Committee on Nitrate and Alternative Curing Agents in Foods*. Washington, DC: National Academy Press, 1981.
- 3 Tricker AR, Preussmann R. Carcinogenic *N*-nitrosamines in the diet: occurrence, formation, mechanisms and carcinogenic potential. *Mutation Research* 1991; **259**(3–4): 277–89.
- 4 Tricker AR. *N*-Nitroso compounds and man: sources of exposure, endogenous formation and occurrence in body fluids. *European Journal of Cancer Prevention* 1997; **6**(3): 226–68.
- 5 Mirvish SS. Blocking the formation of *N*-nitroso compounds with ascorbic acid *in vitro* and *in vivo*. *Annals of the New York Academy of Sciences* 1975; **258**: 175–80.
- 6 Bogovski P, Bogovski S. Animal species in which *N*-nitroso compounds induce cancer. *International Journal of Cancer* 1981; **27**(4): 471–4.
- 7 World Cancer Research Fund (WCRF)/American Institute for Cancer Research (AICR). *Food, Nutrition and the Prevention of Cancer: A Global Perspective*. Washington, DC: WCRF/AICR, 1997.
- 8 Eichholzer M, Gutzwiller F. Dietary nitrates, nitrites, and *N*-nitroso compounds and cancer risk: a review of the epidemiologic evidence. *Nutrition Reviews* 1998; **56**(4 Pt 1): 95–105.
- 9 International Agency for Research on Cancer (IARC). *Evaluation of Carcinogenic Risk to Humans*. IARC Monographs Supplement 7. Lyon: IARC, 1987: 40–74.
- 10 Gangolli SD, van den Brandt PA, Feron VJ, Janzowsky C, Koeman JH, Speijers GJ, *et al.* Nitrate, nitrite and *N*-nitroso compounds. *European Journal of Pharmacology* 1994; **292**(1): 1–38.
- 11 Jakszyn P, Agudo A, Ibanez R, Garcia-Closas R, Pera G, Amiano P, *et al.* Development of a food database of nitrosamines, heterocyclic amines, and polycyclic aromatic hydrocarbons. *Journal of Nutrition* 2004; **134**(8): 2011–4.
- 12 EPIC Group of Spain. Relative validity and reproducibility of a diet history questionnaire in Spain. II. Nutrients. *International Journal of Epidemiology* 1997; **26**(Suppl. 1): S100–9.
- 13 Slimani N, Farran A, Torrent M, Farriol N, Moreuo I, Hemon B, *et al.* *European Prospective Investigation into Cancer and Nutrition (EPIC) Food Composition Tables – Spain*, 2nd ed. Lyon: International Agency for Research on Cancer, 1996.
- 14 Pera G, González CA y grupo EPIC-España. Consumo de suplementos vitamínicos y minerales en población adulta sana de cinco provincias de España. *Gaceta Sanitaria* 1999; **13**(4): 326–7.
- 15 Armitage P, Berry G, Matthews J. *Statistical Methods in Medical Research*. London: Blackwell Science, 2002.
- 16 Instituto Nacional de Estadística (INE). *Cifras oficiales de Población. Padron Municipal* [online], 2003. Available at <http://www.ine.es/inebase/index.html>.

- 17 R Development Core Team. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing, 2004. Also available at <http://www.R-project.org>
- 18 Dich J, Jarvinen R, Knekt P, Penttila PL. Dietary intakes of nitrate, nitrite and NDMA in the Finnish Mobile Clinic Health Examination Survey. *Food Additives and Contaminants* 1996; **13**(5): 541–52.
- 19 Spiegelhalter B, Eisenbrand G, Preussmann R. Volatile nitrosamines in food. *Oncology* 1980; **37**(4): 211–6.
- 20 Wilkens LR, Kadir MM, Kolonel LN, Nomura AM, Hankin JH. Risk factors for lower urinary tract cancer: the role of total fluid consumption, nitrites and nitrosamines, and selected foods. *Cancer Epidemiology, Biomarkers & Prevention* 1996; **5**(3): 161–6.
- 21 Agudo A, Pera G. Vegetable and fruit consumption associated with anthropometric, dietary and lifestyle factors in Spain. EPIC Group of Spain. European Prospective Investigation into Cancer. *Public Health Nutrition* 1999; **2**(3): 263–71.
- 22 Martinez-Gonzalez MA, Perez-Gutierrez R, Martinez-Gonzalez J, Garcia-Martin M, Bueno-Cavanillas A. Dietary intake of some food items in smokers and non-smokers in a Mediterranean population. *European Journal of Public Health* 1997; **7**: 40–4.
- 23 Dallongeville J, Marecaux N, Fruchart JC, Amouyel P. Cigarette smoking is associated with unhealthy patterns of nutrient intake: a meta-analysis. *Journal of Nutrition* 1998; **128**(9): 1450–7.
- 24 Preston AM. Cigarette smoking – nutritional implications. *Progress in Food & Nutrition Science* 1991; **15**(4): 183–217.
- 25 Pobel D, Riboli E, Cornee J, Hemon B, Guyader M. Nitrosamine, nitrate and nitrite in relation to gastric cancer: a case–control study in Marseille, France. *European Journal of Epidemiology* 1995; **11**(1): 67–73.
- 26 Yamamoto M, Iwata R, Ishiwata H, Yamada T, Tanimura A. Determination of volatile nitrosamine levels in foods and estimation of their daily intake in Japan. *Food and Chemical Toxicology* 1984; **22**(1): 61–4.
- 27 Tricker AR, Pfundstein B, Theobald E, Preussmann R, Spiegelhalter B. Mean daily intake of volatile *N*-nitrosamines from foods and beverages in West Germany in 1989–1990. *Food and Chemical Toxicology* 1991; **29**(11): 729–32.
- 28 La Vecchia C, D'Avanzo B, Airoldi L, Braga C, Decarli A. Nitrosamine intake and gastric cancer risk. *European Journal of Cancer Prevention* 1995; **4**(6): 469–74.
- 29 De Stefani E, Boffetta P, Mendilaharsu M, Carzoglio J, Deneo-Pellegrini H. Dietary nitrosamines, heterocyclic amines, and risk of gastric cancer: a case–control study in Uruguay. *Nutrition and Cancer* 1998; **30**(2): 158–62.
- 30 Gonzalez CA, Riboli E, Badosa J, Batiste E, Cardona T, Pita S, *et al.* Nutritional factors and gastric cancer in Spain. *American Journal of Epidemiology* 1994; **139**(5): 466–73.
- 31 Rogers MA, Vaughan TL, Davis S, Thomas DB. Consumption of nitrate, nitrite, and nitrosodimethylamine and the risk of upper aerodigestive tract cancer. *Cancer Epidemiology, Biomarkers & Prevention* 1995; **4**(1): 29–36.
- 32 Ellen G, Egmond E, Van Loon JW, Sahertian ET, Tolsma K. Dietary intakes of some essential and non-essential trace elements, nitrate, nitrite and *N*-nitrosamines, by Dutch adults: estimated via a 24-hour duplicate portion study. *Food Additives and Contaminants* 1990; **7**(2): 207–21.
- 33 Cornee J, Lairon D, Velema J, Guyader M, Berthezene P. An estimate of nitrate, nitrite and *N*-nitrosodimethylamine concentrations in French food products or food groups. *Sciences des Aliments* 1992; **12**: 155–97.
- 34 Maki T, Tamura Y, Shimamura Y, Naoi Y. Estimate of the volatile nitrosamine content of Japanese food. *Bulletin of Environmental Contamination and Toxicology* 1980; **25**(2): 257–61.
- 35 Stryker WS, Salvini S, Stampfer MJ, Sampson L, Colditz GA, Willett WC. Contributions of specific foods to absolute intake and between-person variation of nutrient consumption. *Journal of the American Dietetic Association* 1991; **91**(2): 172–8.
- 36 Byers T, Marshall J, Fiedler R, Zielezny M, Graham S. Assessing nutrient intake with an abbreviated dietary interview. *American Journal of Epidemiology* 1985; **122**(1): 41–50.
- 37 Knight TM, Forman D, Al-Dabbagh SA, Doll R. Estimation of dietary intake of nitrates and nitrites in Great Britain. *Food and Chemical Toxicology* 1987; **25**(4): 277–85.
- 38 Aranceta J, Perez C, Eguileor I, Gonzalez de Goldeano L, Mataix J, Saez de Buruaga J. *Nutritional Survey, 1990*. Technical Documents of Public Series A, No. 9. Vitoria: Government of the Basque Country, 1990.
- 39 Mataix Verú J, Llopis González J, Martínez de Victoria E, Montellano Delgado MA, Lopez Frías M, Aranda Ramirez P. *Evaluation of the Nutritional Status in the Autonomous Community of Andalucía*. Granada: Public Health Service & Institute of Nutrition and Food Technology of the University of Granada, 1999.
- 40 Agudo A, Pera G, Rodriguez M, Quiros JR, Navarro C, Martinez C, *et al.* Changes in smoking habits in adults: results from a prospective study in Spain. *Annals of Epidemiology* 2004; **14**(4): 235–43.