

Crops and Soils Review

Cite this article: George NA, Coorey R, Dixon K, Bennett SJ (2023). A review of the *de novo* domestication and cultivation of edible Australian native plants as food crops. *The Journal of Agricultural Science* **161**, 778–793. <https://doi.org/10.1017/S0021859624000078>

Received: 2 May 2023

Revised: 30 November 2023

Accepted: 21 December 2023

First published online: 22 February 2024

Keywords:

agronomic research; bush tucker; crop domestication; germplasm collection; native foods

Corresponding author:

Nicholas Alexander George;

Email: nicholas.george@curtin.edu.au

A review of the *de novo* domestication and cultivation of edible Australian native plants as food crops

Nicholas Alexander George , Ranil Coorey, Kingsley Dixon and Sarita Jane Bennett

School of Molecular and Life Sciences, Curtin University, Bentley, Perth, WA 6102, Australia

Abstract

Australia has a diverse and unique native flora with thousands of edible plant taxa, many of which are wild relatives of important food crops. These have the potential to diversify and improve the sustainability of Australian farming systems. However, the current level of domestication and cultivation of Australian plants as food crops is extremely limited by global standards. This review examines the current status and potential for future *de novo* domestication and large-scale cultivation of Australian plants as food crops. This is done in the context of international new crop development and factors that impact the success or failure of such efforts. Our review finds considerable potential for native Australian plants to be developed as food crops, but the industry faces several significant challenges. The current industry focuses on niche food markets that are susceptible to oversupply. It also suffers from inconsistent quantity and quality of product, which is attributed to a reliance on wild harvesting and the cultivation of unimproved germplasm. More active cultivation is necessary for industry growth, but attempts have historically failed due to poorly adapted germplasm and a lack of agronomic information. The *de novo* domestication and large-scale cultivation of Australian plants as food crops will require an investment in publicly supported multidisciplinary research and development programmes. Research programmes must prioritize the exploration of plants throughout Australia and the collection and evaluation of germplasm. Programmes must also seek to engage relevant stakeholders, pursue participatory research models and provide appropriate engagement and benefit-sharing opportunities with Indigenous Australian communities.

Introduction

Australia has a diverse and unique native flora, spanning major biome types from tropical to arid to alpine, with thousands of edible plant taxa, many of which are wild relatives of important global food crops. Over the last 40 years, individuals have argued for the domestication and cultivation of edible Australian plants as crops (e.g. Yen, 1993; Considine, 1996; Bell *et al.*, 2011; Abdelghany *et al.*, 2021; Drake *et al.*, 2021), yet, the commercial cultivation of Australian plants for food remains limited. The Macadamia nut (*Macadamia integrifolia* Maiden & Betche, *M. tetraphylla* L.A.S. Johnson, and their hybrids), native to coastal areas of the states of Queensland and New South Wales and domesticated in Hawaii from the 1920s, remains the only widely grown food crop endemic to the Australian continent (Shigeura and Ooka, 1984; Johnson and Burchett, 1996).

The lack of domesticated native Australian food crops is surprising, given that multiple food crops have been derived from the native flora of every other inhabited continent (Stalker *et al.*, 2021). However, the absence of native Australian domesticates should not imply a lack of suitability of Australian plants to become food crops. In this review, we argue that there is considerable potential for the *de novo* domestication and cultivation of Australian plants as food crops and that investing in such domestication and cultivation could assist in diversifying Australian farming systems, providing environmental and economic sustainability benefits (Lin, 2011; Kahane *et al.*, 2013; Isbell *et al.*, 2017; Burchfield *et al.*, 2019). We review the prior research and development of Australian native food plants in light of international new crop development efforts and the factors that impact the success or failure of these efforts. Constraints to developing native Australian crops and associated farming industries are identified, along with a framework for overcoming these constraints.

The need for greater crop diversity in Australian farming systems

Australia has around 60 million hectares of actively cultivated farmland (ABARES, 2022a). Around one-third of this area, predominantly in the continent's southwest, south and east,

comprises monocultures of rainfed, annual grain crops (ABARES 2022a, 2022b). Three crops, wheat (*Triticum aestivum* L.), barley (*Hordeum vulgare* L.) and canola (*Brassica napus* L.), represent approximately 90% of the planted area (ABARES, 2022a, 2022b), and contribute significantly to globally traded staple foods and international food security (ABARES, 2022a, 2022b; FAOSTAT, 2022). These intensive, high-input, low-diversity monocultures of annual crops are not considered sustainable, given their negative environmental impacts and lack of resilience to disturbances such as climate change (FAO, 2017; Pretty *et al.*, 2018). Lack of diversity in agricultural systems is not limited to Australia: globally, farming systems are underpinned by an increasingly limited number of major annual crop taxa, and global diets are becoming less diverse, which negatively impacts the resilience of global food systems (Khoury *et al.*, 2014; Martin *et al.*, 2019; Bentham *et al.*, 2020).

Factors such as interannual weather variability, water insecurity, soil degradation and loss, ecosystem disturbance, pests and diseases pressure and changing global markets for agricultural commodities threaten the long-term viability of Australian agricultural systems, and such disturbances are likely to worsen in the future (Keating and Carberry, 2010; Cresswell *et al.*, 2021). Climate change poses a particularly serious challenge. Productivity in Australian grain farming has already been negatively impacted by the aridification of previously mesic production environments (Sudmeyer *et al.*, 2016; Hochman *et al.*, 2017). It is predicted that Australian agricultural industries and the agricultural sector worldwide will need to make significant changes to agronomic management and species selection to adapt to future climatic conditions (Howden *et al.*, 2010).

Increasing agrobiodiversity is a well-recognized strategy to improve the resilience and sustainability of agricultural systems (Jacobsen *et al.*, 2015; Isbell *et al.*, 2017; Li *et al.*, 2023). Agrobiodiversity can be increased through the production of minor crops, the introduction of exotic crops or the *de novo* domestication of new taxa (Massawe *et al.*, 2016; Toensmeier, 2016; Mustafa *et al.*, 2019; N'Danikou and Tchokponhoue, 2019). *De novo* domestication means the domestication and cultivation of species with little or no prior history of domestication or cultivation. New species provide opportunities for diversification of farming systems and enable transformational changes required for long-term sustainability (Rickards and Howden, 2012; Petersen and Snapp, 2015; Pretty *et al.*, 2018). For example, the use of high-diversity agricultural systems which favour perennial species, termed *perennial polycultures*, is proposed as one strategy for increased agricultural sustainability (Brummer *et al.*, 2011; Iverson *et al.*, 2014; Toensmeier, 2016; Crews *et al.*, 2018), but is difficult to achieve in Australia given existing crop species options (Hatton and Nulsen, 1999; Hobbs and O'Connor, 1999; Pate and Bell, 1999; Bell *et al.*, 2010; Loomis, 2022). Native Australian food crops could potentially provide economically viable perennial species that are well-adapted to local production environments, making perennial polycultures more feasible (Shelef *et al.*, 2017).

Many of Australia's most economically important agricultural industries were developed only recently (Nelson and Hawthorne, 2000; Salisbury *et al.*, 2016). Nearly three-quarters of the total value of crop production in Australia from the 1950s to 1990s is derived from new crops and emerging agricultural industries (Fletcher, 2002; Salvin *et al.*, 2004; Foster, 2014). Along with diversification benefits, native Australian food crops could, therefore, also lead to new, economically valuable agricultural industries. Globally, many governments and organizations recognize

the value of new crops and invest in developing new crop species to enable similar economic opportunities (Janick *et al.*, 1996; Williams, 2005; Foster, 2014).

Can Australian flora be a source of new food crops?

Australia's native flora comprises approximately 20 000 recognized taxa of vascular land plants, around 85% of which are endemic (DEWR, 2007; Chapman, 2009; Broadhurst and Coates, 2017). Many individual taxa are known to be edible, with all plant food groups represented – cereals, pulses, nuts, roots and tubers, fruits and vegetables (Isaacs, 1987; Low, 1991; Latz, 1995; Bindon, 1996). Lists of edible native Australian plants have been compiled, derived predominately from records of plants traditionally eaten by Indigenous Australians (e.g. Isaacs, 1987; Low, 1991; Latz, 1995; Bindon, 1996; Hansen and Horsfall, 2019). Hansen and Horsfall (2019) and Latz (1995) provide comprehensive and regionally specific lists. The former documents approximately 400 edible taxa in southwest Western Australia, and the latter documents 110 taxa in the central desert region. Southwestern Australia has approximately 8000 native vascular plant taxa (FloraBase, 2021), and the central desert 1500 (FloraNT, 2021; AVH, 2023), suggesting 5–7% of local plant species are edible. This is comparable to or slightly lower than global estimates, suggesting that 10–20% of local flora in temperate regions globally could be edible (Rapoport and Drausal, 2013).

Lists of edible species are likely incomplete due to the loss of traditional Indigenous knowledge following European colonization, a lack of comprehensive documentation, cultural preferences in plant use and differing definitions of 'edible' (Rapoport and Drausal, 2013). To illustrate this, southwestern Australia has 39 genera in the legume sub-family Faboideae, representing 500 currently named taxa, with many endemic (FloraBase, 2021). Central Australia has 41 genera representing 138 currently named taxa in Faboideae (FloraNT, 2021). Hansen and Horsfall (2019) and Latz (1995) do not report the seed of these taxa as having been traditionally eaten, and there are no widely published reports of the seed of any Faboideae being eaten elsewhere in Australia. This is despite some taxa being crop wild relatives, such as *Glycine* Willd. and *Vigna* Savi. The seed composition of Australian Faboideae is not well studied, but anti-nutritional and potentially toxic compounds commonly occur in legumes (Tiwari *et al.*, 2011; Kumar *et al.*, 2022), and may have limited the traditional use of Australian taxa as food. Such compounds have been reduced or eliminated via appropriate food preparation techniques and breeding in domesticated legumes and could potentially be eliminated in Australian native legumes (Bell *et al.*, 2011; Bohra *et al.*, 2022; Zhang *et al.*, 2022). Many Australian Faboideae could, therefore, be considered 'potentially edible' and worth investigating for *de novo* domestication (Bell *et al.*, 2011). Including even a small number of the Faboideae expands the edible proportion of Australian flora to the high end of global estimates (Rapoport and Drausal, 2013). This simple estimate illustrates how Australia could have 4000 or more plant species suitable for exploration as potential food crops.

What is the potential of edible Australian flora for producing new crops?

Various traits influence crop domestication potential, the specific traits favouring domestication vary between species and crop type

(DeHaan *et al.*, 2016; Fuller *et al.*, 2023), and some plant taxa are more straightforward to domesticate than others (DeHaan *et al.*, 2016; Stalker *et al.*, 2021). Given the diversity of edible plant taxa native to the continent, it seems probable that some Australian species will have a combination of traits favouring *de novo* domestication. The development and global success of the Macadamia nut industry illustrates that some Australian species have traits that make them suitable for domestication. Furthermore, related plant taxa have often been independently domesticated in geographically separate regions, most probably because these taxa share common traits favouring their domestication (Wang *et al.*, 2014; Renny-Byfield *et al.*, 2016; Wu *et al.*, 2018). At least 130 Australian taxa are crop wild relatives (Rapoport and Drausal, 2013; Norton *et al.*, 2017), including species from *Oryza* L. (rice) (Henry, 2019; Abdelghany *et al.*, 2021), *Sorghum* (L.) Moench (Ananda *et al.*, 2020), *Vigna* Savi (beans) (Lawn 2015) and *Glycine* Willd. (soybean) (Hwang *et al.*, 2019). As well as providing a genetic resource for associated breeding programmes of domestic crops (Henry, 2023), such taxa are likely to share some traits that favoured their relatives' domestication, increasing their potential for *de novo* domestication. This suggests that edible Australian flora has good potential for producing new crops.

An overview of the current Australian native food industry

Historical use of Australian plants as food by humans

Australia's edible native flora has been extensively utilized by people since their arrival on the continent some 65 000 years ago, although a debate about whether plant cultivation was practised on the Australian continent before European colonization is ongoing (Pascoe, 2014; Keen 2021; Sutton and Walshe, 2021; Denham and Donohue, 2023). Plant domestication and cultivation, where plants have diverged morphologically and genetically from wild ancestors due to human selection and the reliance of human communities on these plants for most of their food intake, does not appear to have occurred in the Australian continent before the arrival of Europeans (Sutton and Walshe, 2021). Plant domestication is a continuum, however, without a well-defined start and endpoint (Winterhalder and Kennett, 2006; Meyer *et al.*, 2012; Zeder, 2015; Stetter *et al.*, 2017; Stalker *et al.*, 2021; Fuller *et al.*, 2023). Some 'early' plant cultivation and domestication are not readily distinguished from other forms of plant exploitation, particularly in the archaeological record (Zeder, 2015; Denham and Donohue, 2023).

There is evidence for the intensive management and manipulation of Australian flora by people via practices such as the use of fire and the translocation of food plants (Hallam, 1989; Bowman, 1998; Clarke, 2011; Ens *et al.*, 2017; Silcock, 2018; Lullfitz *et al.*, 2020a, 2020b; Keen, 2021; Fahey *et al.*, 2022), and 'non-agricultural' cultures are also known to have engaged in 'niche constructive' behaviours that maintained or increased the productivity of their environments (Smith, 2011; Anderson, 2013; Lightfoot *et al.*, 2013; Thompson *et al.*, 2021a, 2021b). Such activities can result in lasting changes in the geographic distribution and genetic composition of plant taxa (Levis *et al.*, 2017; Coughlan and Nelson, 2018; Levis *et al.*, 2018; Pavlik *et al.*, 2021). This appears to have resulted in detectable changes in the genetics of some Australian taxa (Rangan *et al.*, 2015; Lullfitz *et al.*, 2020a, 2020b) and may have also resulted in phenotypic changes. For example, it has been proposed that the large

grain size in some native Australian *Oryza* may reflect human selection (Henry, 2019). This may impact efforts to domesticate Australian species as crops in the future.

Features of the current industry

The possibility of cultivating edible Australian plants as crops has been acknowledged for over a century (Maiden, 1889). However, the Australian 'native foods industry' did not commence until approximately the 1980s (Cherikoff and Brand, 1983; Brand-Miller and Cherikoff, 1985; Cherikoff and Brand, 1988). Commercial native food production now takes place in all Australian states and territories (Clarke, 2012; Sultanbawa and Sultanbawa, 2016), but the current industry is small in terms of total production and economic value. Excluding Macadamia, total output was estimated to average only 8 tonnes in 2010 (Clarke, 2013), with a farm-gate value of \$21 million in 2019 (Laurie, 2020), in comparison to the total gross value of Australian agriculture of \$55 billion in 2015–16 (ABARES 2022a, 2022b).

Most of the production and economic value of the industry is represented by only 11 taxa (Table 1) (Clarke, 2013; Laurie, 2020). These came to dominate the industry through multiple 'organic' routes over four decades and are mainly used as 'niche' food additives or flavourants (i.e. herbs and spices) or as fruits that are processed into value-added products. Feedstock material used by the food industry is obtained from wild and cultivated sources (Clarke, 2013; Laurie, 2020). Most of the taxa are native to eastern and northern Australia's tropical, subtropical and oceanic climate zones (Fig. 1A), except for *Solanum centrale* (Fig. 1B), which grows in arid zones of Australia. Since the native ranges of the taxa do not overlap with the Australian grain production regions, these species offer little potential to diversify existing grain industries with locally adapted crops. Those taxa that do grow in the grain belt include *Santalum acuminatum* (Quandong) (Fig. 1C) and multiple species of *Acacia* (Table 1). Quandong is an obligate root hemiparasite that requires a host tree and produces a fleshy fruit with an edible nut (Ahmed and Johnson, 2000; Lee, 2013). These traits give it limited potential for broadscale planting in grain-producing regions. Conversely, the various species of *Acacia* produce a grain legume (or *pulse*) and offer the prospect of large-scale planting for bulk food production (Ahmed and Johnson, 2000; Bartle *et al.*, 2002; Lee, 2013). Aside from the taxa in Table 1, around 40 other Australian plant taxa are sold for food (Table 2) (CNFS, 2022; Tucker Bush, 2022). These represent a greater diversity of food types than the taxa in Table 1 and are native to a broader range of environments. However, the majority are native to eastern Australia and are used only as niche food additives.

Literature dating back to the late 1990s has examined the utilization of Australian native plants as food crops and the development of the native food industry. Authors have consistently concluded there is a good market potential for Australian food plants, especially those considered novel and with exceptional nutritional profiles (Cherikoff, 2000; Konczak *et al.*, 2009; Clarke, 2012; Sultanbawa and Sultanbawa, 2016; Birch *et al.*, 2023). They also conclude that Australian plants offer valuable opportunities for diversifying the continent's agricultural systems with well-adapted new crops that can enhance environmental and economic sustainability (Considine, 1996; Ahmed and Johnson, 2000; Bell *et al.*, 2011; Abdelghany *et al.*, 2021; Drake *et al.*, 2021; Canning, 2022). However, the various authors have

Table 1. Plant taxa and their relatives that are the current focus of the Australian native food industry (Clarke, 2013; Laurie, 2020)

Species	Common name	Family	Food type	Approximate native range	Climate
<i>Acacia</i> spp. Commonly <i>A. victoriae</i> ; <i>A. adsurgens</i> ; <i>A. aneura</i> ; <i>A. coleii</i> ; <i>A. coriacea</i> ; <i>A. cowleana</i> ; <i>A. kempeana</i> ; <i>A. murrayana</i> ; <i>A. tenuissim</i> ; <i>A. pycnantha</i> ; <i>A. retinodes</i> ; <i>A. sophorae</i>	Wattle	Fabaceae	Seed	Continent wide genus	
<i>Acronychia acidula</i> and <i>A. oblongifolia</i>	Lemon Aspen, Pigeon Berry, Southern Lemon Aspen, White Aspen	Rutaceae	Herb/spice	Coastal Qld, NSW and Vic	Cfb, Cfa, Cwa
<i>Backhousia anisata</i> (syn <i>Syzygium anisatum</i>) and <i>Backhousia citriodora</i>	Anise myrtle, Aniseed myrtle, Ringwood, Lemon myrtle, Lemon Ironwood, Lemon scented myrtle, Sweet Verbena tree	Myrtaceae	Herb/spice	Mainly coastal Qld	Cfa, Cwa, Aw
<i>Citrus glauca</i> (syn <i>Eremocitrus glauca</i>), <i>C. australasica</i> (syn <i>Microcitrus australasica</i>)	Finger Lime, Desert lime	Rutaceae	Fruit	Coastal Qld and NSW, Inland areas of Qld, NSW and SA.	Cfa, Bsh, Bsk, Bwh
<i>Davidsonia jerseyana</i> , <i>D. johnsonii</i> , <i>D. pruriens</i>	Davidson Plum, Queensland Itchtree, Ooray Smooth Davidsonia	Cunoniaceae	Fruit	Coastal Qld and NSW	Am, Af, Cfa, Cwa
<i>Kunzea pomifera</i>	Muntries, Emu apple	Myrtaceae	Fruit	South-eastern SA and western Vic	Csb, Csa, Bsk
<i>Santalum acuminatum</i> and <i>S. spicatum</i>	Quandong, Sweet Quandong, Wild Peach, Desert Peach, Native Peach, Guwandhuna, Gutchu, Wanjanu, Mangata, Goorti, Wadjal	Santalaceae	Fruit/nut	Southern Australia	All except tropical
<i>Solanum centrale</i>	Bush tomato, Desert raisin	Solanaceae	Fruit	Inland areas mainly WA, NT and SA	Bwh, Bsh
<i>Syzygium luehmannii</i> , <i>S. australe</i> and <i>S. oleosum</i>	Riberry, Small-leaved Lilly Pilly, Clove Lilli Pilli	Myrtaceae	Fruit	Coastal Qld and NSW	Am, Af, Cfa, Cwa
<i>Tasmannia lanceolata</i>	Tasmanian pepper, Mountain pepper, Native pepper, Pepperberry, Pepperleaf	Winteraceae	Herb/spice	Tas, Vic and southeast NSW	Cfb
<i>Terminalia ferdinandiana</i>	Kakadu Plum, Billy Goat Plum, Green Plum, Salty Plum, Wild Plum, Murunga, Marnybi, Manmohpan, Kullari Plum, Gubinge	Combretaceae	Fruit	Northern NT and northwest WA	Aw, Bsh

The approximate native range obtained from the Australasian Virtual Herbarium (AVH 2023).

WA, Western Australia; NT, Northern Territory; SA, South Australia; Qld, Queensland; NSW, New South Wales; Vic, Victoria; Tas, Tasmania; Af, Tropical rainforest; Am, Tropical monsoon; Aw, Tropic Savanna with dry winter; Bsh/Bsk, semi-arid hot; Bwh, arid hot; Cfa, humid sub-tropical; Cfb, oceanic; Csa, Mediterranean hot summer; Csb, Mediterranean warm summer; Cwa, dry-winter humid subtropical.

also identified significant challenges. The current emphasis on niche food markets makes them susceptible to oversupply (Clarke, 2012; Clarke, 2013), necessitating the development of new crops capable of supplying larger markets. The industry also grapples with inconsistent quantity and quality of supply, attributed mainly to a reliance on wild harvesting and the cultivation of unimproved germplasm, so there is a need for more active cultivation and the use of improved cultivars to address this problem (Stynes, 1997; Salvin *et al.*, 2004; Lee, 2013; Abdelghany *et al.*, 2021). Although some active cultivation efforts are underway, they face obstacles such as a lack of information on cultivation methods, challenges posed by pests and diseases and an overreliance on manual labour (Ahmed and Johnson, 2000; Clarke, 2012, 2013; Lee, 2013; Sultanbawa and Sultanbawa, 2016). To advance the industry, ongoing research is required, with a critical need for cultivar development, general agronomy and market development (Gorst, 2002; Salvin *et al.*, 2004; Lee

and Six, 2010; Clarke, 2012, 2013; Sultanbawa and Sultanbawa, 2016).

Research on individual taxa

To quantify the extent of research on individual taxa published in the 20 years between 2001 and 2021 (the time since the last scholarly review of the industry by Ahmed and Johnson, 2000), the Thomson Reuters Web of Science database was searched for publications in scholarly journals relating to the taxa in Table 1 and Table 2. We identified 234 research articles mentioning at least one of the taxa from Table 1 (Fig. 2). *Acacia* was excluded, given the large number of individual taxa in the genus, taxa found outside Australia and the non-food use of the species globally. No relevant published work was found for the taxa in Table 2. For comparison, the total research output for the taxa with the most published research is comparable to the




Figure 1. The maps in panels A–D show the distribution of taxa listed in Table 1. The maps are based on collection information from the Australian Virtual Herbarium (AVH 2023). The Australian grain production zone is shown in grey. The distribution of (a) *Acronychia acidula* and *A. oblongifolia*, *Backhousia anisata* (syn *Syzygium anisatum*) and *B. citriodora*, *Citrus glauca* (syn *Eremocitrus glauca*), *C. australasica* (syn *Microcitrus australasica*), *Davidsonia jerseyana*, *D. johnsonii*, *D. pruriens*, *Kunzea pomifera*, *Tasmania lanceolata* and *Terminalia ferdinandiana*. (b) *Solanum central*; (c) *Santalum acuminatum* and *S. spicatum*, (d) *Acacia victoriae*; *A. adsurgens*; *A. aneura*; *A. coleii*; *A. coriacea*; *A. cowleana*; *A. kempeana*; *A. murrayana*; *A. tenuissim*; *A. pycnantha*; *A. retinodes*; *A. sophorae*.

Australian research output for minor crops like kiwifruit, with 41 papers (*Actinidia*), or Blueberry (*Vaccinium*), with 106 papers over the same period (ABARES, 2022a, 2022b). Publications covered by the Web of Science Core Collection are assigned to at least one subject area category. We found that the journal articles relating to the taxa mainly related to food sciences, chemistry or nutrition (Fig. 2). Papers addressing other areas relevant to the development of the industry, such as agronomy and genetics, represented only a quarter of all publications. Ahmed and Johnson (2000) observed that most published research at the time of their review was focused on the compositional analysis of native food plants, and other critical areas were lacking. Our literature search shows this trend has continued, and whilst compositional analysis is essential for industry development, lack of research in other areas has likely contributed to the slow pace of industry development.

Agrifutures Australia, previously called the Rural Industries Research & Development Corporation (RIRDC), is a statutory authority established by the Australian Federal Government to support new and emerging industries. Since 2000, Agrifutures Australia has produced 43 reports dealing wholly or partly with different aspects of the Australian native foods industry (Agrifutures, 2022), but only one has been published since 2017

(Table 1). While agronomy and germplasm improvement are addressed more frequently than in the published literature, food science or compositional analyses remain the most common area of research (Fig. 3). Like the scientific literature, it is a relatively small number of publications for any individual taxon (Fig. 4).

Some individual edible taxa have had more concerted research aimed at their domestication as crops. Germplasm screening and selection, genetics and reproductive biology studies and some agronomy have been undertaken for *Kunzea pomifera* (Page, 2004; Page et al., 2006a, 2000b; Do et al., 2014, 2018a, 2018b). Native *Acacia* have been the focus of development as a grain crop in Australia and elsewhere (Lister et al., 1996; Maslin et al., 1998; Bartle et al., 2002; Hele, 2002; Rinaudo et al., 2002; Midgley and Turnbull, 2003; Rinaudo and Cunningham, 2008). Native Australian legumes, aside from *Acacia*, have also been explored as pulses, with the examination of grain yield and seed composition (Rivett et al., 1983; Bell et al., 2011; Ryan et al., 2011; Bell et al., 2012). Several commercial *Citrus* varieties have been developed by hybridizing Australian native *Citrus* spp. with domestic *Citrus* spp. (Sykes, 1997; Hele, 2001; Agrifutures, 2017). Native *Oryza* spp. has been considered as source of germplasm for improving domestic rice, but germplasm collection and

Table 2. The edible plant taxa sold by the Tuckerbush and Creative Native Food Service companies at the time of writing in 2023

Species	Common name	Family	Type	Approximate range
<i>Adansonia gregorii</i>	Boab Roots	Malvaceae	Root/tuber	Northwest WA
<i>Alpinia caerulea</i>	Australia Ginger	Zingiberaceae	Herb/spice	Coastal Qld and NSW
<i>Antidesma bunius</i> and <i>A. erostre</i>	Bignay/Wild Grape	Phyllanthaceae	Fruit	Coastal Qld
<i>Apium annuum</i> and <i>prostratum</i>	Native celery parsley	Apiaceae	Vegetable	Coastal southern and eastern Australia
<i>Araucaria bidwillii</i>	Bunya	Araucariaceae	Nut	Coastal Qld and NSW
<i>Arthropodium strictum</i>	Chocolate lily	Asparagaceae	Roots/ tubers	Mainly SA, Vic, NSW, Qld
<i>Athertonia diversifolia</i>	Atherton Almonds	Proteaceae	Nut	Coastal Qld and NSW
<i>Atractocarpus fitzalanii</i>	Native Gardenia	Rubiaceae	Fruit	Coastal Qld
<i>Atriplex nummularia</i>	Saltbush	Amaranthaceae	Vegetable	Inland areas of all mainland states
<i>Austromyrtus dulcis</i>	Midyim Berries	Myrtaceae	Fruit	Coastal Qld and NSW
<i>Barbarea australis</i>	Native wintercress	Brassicaceae	Vegetable	Tas
<i>Brachychiton populneus</i>	Jurrajin seed	Malvaceae	Seed	Qld, NSW, Vic
<i>Chamelaucium uncinatum</i>	Geraldton Wax	Myrtaceae	Herb/spice	Southwestern WA
<i>Cymbopogon ambiguus</i>	Native lemongrass	Poaceae	Herb/spice	All mainland states
<i>Dianella revoluta</i>	Dianella	Asphodelaceae	Fruit	Southern and eastern Aust.
<i>Dimocarpus australianus</i>	Rainforest lychee	Sapindaceae	Fruit	North Coastal Qld
<i>Dioscorea hastifolia</i>	Warrine	Dioscoreaceae	Roots/ tubers	Southwestern WA
<i>Diospyros australis</i>	Black Plum	Ebenaceae	Fruit	Coastal Qld and NSW
<i>Diploglottis australis</i> and <i>D. campbellii</i>	Large and small leaf Tamarind	Sapindaceae	Fruit	Coastal Qld and NSW
<i>Disphyma crassifolium</i>	Karkalla Beach bananas	Aizoaceae	Herb/spice	Southern Aust.
<i>Elaeocarpus angustifolius</i>	Blue quandone	Elaeocarpaceae	Fruit	Coastal Qld
<i>Enchylaena tomentosa</i>	Ruby saltbush	Amaranthaceae	Fruit	All mainland states
<i>Eremophila alternifolia</i>	Scented Emu Bush	Scrophulariaceae	Herb/spice	Predominantly southern inland areas of WA, SA
<i>Eupomatia laurina</i>	Bolwarra	Eupomatiaceae	Herb/spice	Coastal Qld and NSW
<i>Ficus coronata</i>	Sandpaper fig	Moraceae	Fruit	Coastal Qld and NSW
<i>Haemodorum spicatum</i>	Mean	Haemodoraceae	Roots/ tubers	Southwestern WA
<i>Hibiscus heterophylla</i>	Native rosella	Malvaceae	Fruit/flower	Coastal Qld and NSW, northern WA, NT
<i>Melastoma affine</i>	Blue Toungue	Melastomataceae	Fruit	Coastal Qld and NSW, northern WA, NT
<i>Mentha satuireioides</i> and <i>M. australis</i>	Bush mint	Lamiaceae	Herb/spice	NSW, Vis, southern SA and Qld
<i>Microseris lanceolata</i> and <i>M.</i> <i>scapigera</i>	Yam daisy	Asteraceae	Roots/ tubers	Southwestern Aust.
<i>Myoporum insulare</i> and <i>M.</i> <i>montanum</i>	Boobialla/Waterbush	Scrophulariaceae	Fruit	Coastal southern Australia
<i>Olearia axillaris</i>	Coastal Rosemary/Coastal daisy	Asteraceae	Herb/spice	Coastal southern and western Australia
<i>Pipturus argenteus</i>	Native mulberry	Urticaceae	Fruit	Coastal Qld
<i>Platysace deflexa</i>	Youlk	Apiaceae	Roots/ tubers	Southwestern WA
<i>Plectranthus graveolens</i> (<i>Coleus</i> <i>graveolens</i>)	Basil bush	Lamiaceae	Herb/spice	Coastal Qld and NSW
<i>Pleiogynium timorense</i>	Burdekin plum	Anacardiaceae	Fruit	Coastal Qld
<i>Podocarpus elatus</i>	Illawara plum	Podocarpaceae		Coastal Qld and NSW

(Continued)

Table 2. (Continued.)

Species	Common name	Family	Type	Approximate range
<i>Pouteria australis</i>	Black apple	Sapotaceae	Fruit	Coastal southern Qld and NSW
<i>Prostanthera rotundifolia</i> and <i>P. incisa</i>	Native mints, Native Thyme	Lamiaceae		NSW, Vic, Tas
<i>Rubus probus</i>	Atherton raspberry	Rosaceae	Fruit	Coastal Qld
<i>Scaevola spinescens</i>	Maroon bush	Goodeniaceae	Fruit	Inland areas of all mainland state
<i>Sterculia quadrifida</i>	Peanut tree	Malvaceae	Seed	Coastal Qld, NT and northern WA
<i>Tetragonia tetragonoides</i>	Warrigal greens	Aizoaceae	Vegetable	Southern Aust.

WA, Western Australia, NT, Northern Territory, SA, South Australia, Qld, Queensland, NSW, New South Wales, Vic, Victoria, Tas, Tasmania.

characterization and the systematic identification of research priorities have also taken place with the aim of *de novo* domestication (Henry *et al.*, 2010; Henry, 2012; Henry, 2019; Abdelghany *et al.*, 2021). Germplasm screening (Davies *et al.*, 2005), genetic analysis (Shapter *et al.*, 2013; Mitchell *et al.*, 2015) and commercialization of elite lines have been undertaken for *Microlaena stipodes*, a widespread native grass that produces an edible grain similar to rice (Chivers *et al.*, 2015; Shapter and Chivers, 2015). Several other grass species are also being actively investigated as potential grain crops (Khoddami *et al.*, 2020; Drake *et al.*, 2021). Despite this research and development effort, a large-scale commercial agricultural industry has yet to develop for these taxa.

Indigenous engagement and benefit sharing

Australian plants can hold cultural and spiritual significance to people in the Indigenous Australian community (Clarke, 2011), but their engagement and benefit-sharing with the native foods industry has historically been limited (Considine, 1996; Stynes, 1997; Ahmed and Johnson, 2000; Clarke, 2013; Lingard and Martin, 2016; Sultanbawa and Sultanbawa, 2016; Drake *et al.*, 2021). 'The Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising from their Utilization' aims to implement the access and benefit-sharing obligations of the International Convention on Biological Diversity (Lee, 2013; Leha *et al.*, 2019; Sherman and Henry, 2020; Fyfe *et al.*, 2021). Australia ratified the Convention on Biological Diversity, and while it is not presently a participant in the Nagoya Protocol, current domestic measures purport to align with the principles outlined in the protocol (DCCEEW, 2021). However, others have found that traditional knowledge and intellectual property regarding edible native plants currently lack comprehensive legal protection in Australia or may not be adequately protected by existing laws (Leha *et al.*, 2019). Food is globally recognized as an intangible cultural heritage, eligible for special recognition and protection (Di Giovine and Brulotte, 2016; Galanakis, 2019). So even without clear intellectual property ownership or legal frameworks for protecting traditional biological knowledge, moral and ethical obligations remain for those seeking to develop native Australian food crops (Leha *et al.*, 2019; Jarvis *et al.*, 2021; Maclean *et al.*, 2022). Indigenous Australian support for developing native food industries is generally considered conditional on ensuring such industries recognize and respect culturally or spiritually significant plants, along with their traditional uses (NLE, 2022). Any research and development targeting

Australian native food plants must, therefore, acknowledge the ongoing cultural connections of Indigenous Australian peoples with native flora and take steps to ensure Indigenous Australian communities have opportunities to engage, lead and benefit from the industry.

We believe current ambiguities around Indigenous Australian engagement and benefit sharing will likely hinder research and development activities, and this must be addressed if the industry is to develop. Firstly, even if legislation regarding intellectual property ownership and benefit-sharing is enacted, ownership of traditional knowledge and biological resources and appropriate avenues for benefit-sharing are often unclear. This is the case when information regarding taxa is well-documented, in the public domain, and has been so for a prolonged period. There is ambiguity in the Nagoya Protocol regarding historical germplasm collections and information (Sherman and Henry, 2020). Moreover, some edible plant taxa have wide native ranges spanning many Indigenous Australian communities, and under these circumstances, intellectual property ownership and appropriate benefit-sharing avenues are also unclear. Surveys of Indigenous Australian stakeholders have found that most respondents support the development of a native food industry (NLE, 2022). However, stakeholders still have divergent opinions about whether the commercial development of native food plants as crops should occur and what form the industry should take (Ahmed and Johnson, 2000; Clarke, 2013; Drake *et al.*, 2021). This makes it challenging to identify, engage and coordinate among owners of traditional biological knowledge and find consensus regarding the appropriate way to domesticate and cultivate some species.

The complexity of creating legal and ethical frameworks that both protect and allow the use of traditional ethnobotanical knowledge is a globally recognized problem, as is a lack of engagement and benefit sharing with traditional owners when commercializing traditional foods (Zimmerer and De Haan, 2017; Antonelli, 2023). A discussion of a possible framework for the ethical development of native Australian crops that can address the challenges described above is outside the scope of this review, but the challenges will act as a major obstacle to research and development activities, and nationally consistent legislation and best practice guidelines to address them are urgently needed.

What are the ways forward for native Australian food crops?

Six key areas stand out that would support the use of Australian native plants as food: (i) active cultivation; (ii) germplasm




Figure 2. The number of scholarly journal articles in the Web of Science database that mention: (a) at least one of the plant taxa prioritized by the Australian native food industry (Table 1), published between 2001 and 2021. (b) The number of scholarly publications of Australian native species relating to individual Web of Science subject categories.

collection, characterization and improvement; (iii) basic research; (iv) sustained public funding of critical R&D; (v) greater diversity of food types and cultivation regions; (vi) engagement with Indigenous stakeholders and participatory approaches to research and (vii) consideration of the implications of domestication for conservation, and Indigenous traditional knowledge and use. These are discussed in detail below.

Active cultivation

Wild harvesting remains the primary source of material in the native food industry for many taxa. Wild harvests can provide

economic returns to communities engaged in the collection and are attractive to those advocating for 'ecological' approaches to agribusiness industries (Ahmed and Johnson, 2000; Clarke, 2013; Lee, 2013). However, wild harvesting is also associated with challenges such as inconsistent and unpredictable yields and product quality, limited supply, limited scope for expansion, high demand for labour and possible negative impacts on natural ecosystems (Miers, 2004; RIRDC, 2008; Clarke, 2013; Sultanbawa and Sultanbawa, 2016; Laurie, 2020). Additionally, wild harvesting is not risk-free for workers. For example, wild harvesting of native *Oryza* risks attack by saltwater crocodiles (Abdelghany *et al.*, 2021). Consequently, although wild



Figure 3. The number of reports published by Agrifutures Australia that address subject matter relating to Australian native food plants.

harvesting may be viable for specific regions and species, potentially augmented by ‘active management’ or ‘enrichment planting’ of otherwise wild plant communities (Lee and Courtenay, 2016), industry growth will require active cultivation.

Germplasm collection, characterization and improvement

The wild phenotype of most, if not all, plant taxa is sub-optimal for commercial utilization (Wilson, 2007; Brummer *et al.*, 2011; DeHaan *et al.*, 2016), and suboptimal germplasm is a significant obstacle to viable cultivation of edible Australian plants (Stynes, 1997; Salvin *et al.*, 2004; Lee, 2013; Abdelghany *et al.*, 2021). Germplasm screening and improvement will therefore be essential

for the active cultivation of edible Australian plants to meet the needs of growers and consumers.

Even if wild taxa are identified that pose few challenges to *de novo* domestication, research and development will still be needed to address problematic traits (DeHaan *et al.*, 2016; Toensmeier, 2016). Information regarding commercially important traits is unavailable for most Australian edible taxa, making it impossible to assess their potential as crops or to set research and development priorities. Furthermore, any available information often relates to germplasm of unknown provenance or collections with minimal genetic variation (e.g. see discussions in Sultanbawa and Sultanbawa (2016)). Many edible Australian taxa have broad distributions spanning a considerable range of




Figure 4. The number of reports published by Agrifutures Australia that address the native food taxa.

climatic and edaphic conditions and display high genetic and morphological diversity, including variation in economically important traits (Davies *et al.*, 2005; Ariati *et al.*, 2007; Mitchell *et al.*, 2015; Shapter and Chivers, 2015; Broadhurst *et al.*, 2017; Snowball *et al.*, 2021). Drawing conclusions regarding a species' suitability for domestication from samples of minimal genetic diversity is, therefore, of limited value or even potentially misleading. For example, Ryan *et al.* (2011) identified the representation of genetic diversity in germplasm collections and its evaluation under controlled conditions as a critical gap in assessments of Australian native pulses.

Assembling genetically diverse germplasm collections and then evaluating and selecting improved cultivars is an effective strategy for cultivar development, and remains an essential method for plant breeding globally (Murphy, 2007; Acquah, 2015; Rebetzke *et al.*, 2019), particularly for minor crops that lack resources for research and development (Jacobsen *et al.*, 2015). However, crop domestication initially involved, on average, the modification of only three major traits, which were controlled primarily by single genes (Meyer *et al.*, 2012; Østerberg *et al.*, 2017; Stetter *et al.*, 2017). So, the use of molecular breeding techniques that target limited numbers of single gene traits to 'mimic' domestication events could increase the speed and efficiency of new crop development from Australian flora (Smykal *et al.*, 2018; Gasparini *et al.*, 2021; Luo *et al.*, 2022; Bartlett *et al.*, 2023; Henry, 2023). Such approaches come with risks, though, including that a focus on single genes may over-simplify domestication or neglect the importance of agronomy and genotype-by-environment interactions in the crop phenotype (Passioura, 2020; Van Tassel *et al.*, 2020; Bartlett *et al.*, 2023), and should therefore be used in conjunction with more traditional approaches. Centralized breeding may also not address the specific localized needs of growers (Fadda *et al.*, 2020), or provide opportunities to engage with Indigenous communities (Bartlett *et al.*, 2023). Thus, we believe a range of strategies, including 'traditional' approaches in the early stages of crop development, genetic screening to ensure sufficient diversity and strategic investment in molecular breeding, are appropriate.

Germplasm collection and characterization must consider the historical use of taxa by people, which may have influenced the genetic diversity and geographic distribution of edible taxa. Taxa with high levels of anthropogenic translocation may show a lack of population genetic structure or a structure corresponding to human activity (Lullfittz *et al.*, 2020a, 2020b). Therefore, phylogeographic patterns resulting from human translocation of food plants should be considered in germplasm collection documentation and activities. Plant populations from areas with intensive utilization could be targeted for germplasm collection because these populations may exhibit a higher frequency of individuals with useful genotypes. Further work is needed to investigate the anthropogenic influence on Australian plant genetics and phylogeographic patterns to inform germplasm collection, characterization and crop development activities.

Habitat loss due to clearing native vegetation for agriculture and subsequent land degradation has heavily impacted many ecosystems in Australia (Cresswell *et al.*, 2021). As a result, the remaining native vegetation is often highly fragmented (Hobbs and Yates, 2003; Hopper and Gioia, 2004; Coates *et al.*, 2014; Broadhurst and Coates, 2017), and faces ongoing pressure from pests, disease and climate change (Cresswell *et al.*, 2021). Preservation of genetic diversity is essential for agricultural sustainability globally, as well as conservation efforts. Care must

also be taken as the greater use of native plants as crops brings risks such as gene flow between domesticated and wild populations, posing potential threats to wild populations (Haygood *et al.*, 2003), particularly if wild populations are small and highly fragmented.

The need for basic research

Poorly adapted germplasm, a lack of agronomic information and insufficient investment to address these issues are also commonly identified as significant obstacles to the growth and expansion of the Australian native food industry (Salvin *et al.*, 2004; Clarke, 2012; Clarke, 2013). Developing productive and economically viable farming systems in Australia and globally has relied on basic agricultural research (Hunt *et al.*, 2019; Zaidi *et al.*, 2019; Hunt *et al.*, 2021). There is a need globally for more significant investment in basic research to increase agrobiodiversity and food security (Jacobsen *et al.*, 2015; Toensmeier, 2016), not just in Australia. Many wild plant taxa have been explored as potential crops (Janick, 1996; Janick, 1999; Janick and Whipkey, 2002; Janick and Whipkey, 2007), but few are now commercially viable and widely grown (DeHaan *et al.*, 2016). A common issue is that basic research needed to understand and address problematic plant traits that inhibit economically viable production is missing, as is research to underpin commercially viable agronomy (Jolliff, 1990; Blade and Slinkard, 2002; Wilson, 2007; Abbo *et al.*, 2014; DeHaan *et al.*, 2016). Basic applied research will, therefore, be essential for developing the Australian native food industry.

The need for sustained public funding of R&D

Developing new crops and associated agricultural industries, particularly from undomesticated taxa, requires a sustained, long-term, and multidisciplinary research effort (Wollenweber *et al.*, 2005; Runck *et al.*, 2014; DeHaan *et al.*, 2016). Successful new crop industries in Australia and elsewhere have relied on sustained public research investment in multi-decade and multidisciplinary research programmes (Williams, 2005; Collins and Norton, 2019; Pratley and Kirkegaard, 2019). This is illustrated by the introduction and development of canola (*Brassica napus*) (Colton and Potter, 1999; Salisbury *et al.*, 2016) and edible lines of lupins (*Lupinus angustifolius*) (Nelson and Hawthorne, 2000). The successful development of native Australian crops will require similar research and development efforts. Public funding for such research work in Australia is relatively limited, unsustainable or often non-existent and has also not attracted investment from private enterprises. This stands as a significant hurdle to the further development of the industry.

A possible funding and industry development model already exists in Australia in the form of the Research and Development Corporations (RDC). Supported partly by levies on producers, RDCs have brought demonstrable benefits to several agricultural industries (CRRDC, 2016). Agrifutures Australia is the RDC responsible for supporting research and industry development for edible Australian plants as part of a broader mandate to support new agricultural industries. However, it does not currently provide sustained funding of the sort needed for *de novo* domestication of food crops. Dedicating a specific RDC to native crop domestication could meet global calls for governments to support more diverse and locally adapted food systems built partly on non-conventional crops (Antonelli, 2023). By administering research through national and regional RDC panels comprised of stakeholders,

including members of the Indigenous Australian community, such a model could also provide a mechanism to address engagement and benefit-sharing challenges. When intellectual property ownership is complex, disputed or distributed, such a body could collect levies and administer a consolidated fund.

Greater diversity of food types and regions

The Australian native food industry is biased towards niche markets. Expanding the number of taxa under consideration to encompass more food types, such as grains or pulses, that can supply large-scale staple food markets and to include taxa adapted to a more diverse range of agroecosystems, particularly the grain production zones, will increase opportunities for large-scale native food production to diversify existing extensive agricultural industries. This necessitates research and development towards a more diverse range of edible species.

Participatory research approaches

Participatory germplasm improvement and agronomic research are increasingly common in Australia and globally (Walters *et al.*, 2018; Snapp *et al.*, 2019; Colley *et al.*, 2021; Lacoste *et al.*, 2022). Participatory research includes stakeholders in evaluating and selecting germplasm, developing research targets and conducting agronomic research to address industry constraints (Shelton *et al.*, 2016; Walters *et al.*, 2018; Snapp *et al.*, 2019; Lacoste *et al.*, 2022). Participatory breeding has been used successfully to improve the productivity and quality of crops in several regions, notably in some minor crops (Ceccarelli, 2015; Shelton *et al.*, 2016; Ceccarelli and Grando, 2020; Fadda *et al.*, 2020). On-farm agronomic research can better understand and address complex genotype, management and environmental interactions (Rotili *et al.*, 2020), identify industry needs, and encourage more rapid adoption of new crops and farming practices (Hunt *et al.*, 2019). Participatory research can yield efficiencies in a stretched research funding environment, complement traditional research programmes and create additional avenues for engagement and empowerment of Australian Aboriginal communities.

Conclusions

There is considerable potential for the *de novo* domestication and cultivation of native Australian plants as food crops. Such crops could provide valuable new agricultural industries that increase the long-term sustainability of Australian agricultural systems and contribute to global food security. The primary impediment is inadequate funding and policy needed to underpin appropriate research and development, particularly basic cultivar development and agronomic research needed for active cultivation. Historically, successful new crop programmes show that developing native Australian food crops will require a sustained investment in publicly supported multidisciplinary research and development. This could happen through established Australian agricultural funding frameworks like the RDCs. Research and development activities must commence with collecting and evaluating a range of edible taxa, from throughout the continent, targeting species with the potential for large-scale staple food markets and adapted to a diverse range of agroecosystems. Finally, development programmes must also engage all relevant stakeholders and provide appropriate engagement and benefit-sharing opportunities with Indigenous communities.

Acknowledgements. We sincerely thank Dr Angela Pattison, The University of Sydney, and Professor Sally Thompson, The University of Western Australia, for providing comments on drafts of this review.

Author contributions. N. A. G. and K. D. conceived the review topic and N. A. G. researched and wrote the manuscript. S. J. B., R. C. and K. D. revised the manuscript and provided additional intellectual content.

Funding statement. This research did not receive any specific funding.

Competing interests. None.

Ethical standards. Not applicable.

References

- ABARES (2022a) Australian commodity statistics 2020. Canberra, Australian Government. Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES).
- ABARES (2022b) Snapshot of Australian Agriculture 2022. ABARES Insights. Canberra, Australian Bureau of Agricultural and Resource Economics and Sciences 14.
- Abbo S, van-Oss RP, Gopher A, Saranga Y, Ofner I and Peleg Z (2014) Plant domestication versus crop evolution: a conceptual framework for cereals and grain legumes. *Trends in Plant Science* **19**, 351–360.
- Abdelghany G, Wurm P, Hoang LTM and Bellairs SM (2021) Commercial cultivation of Australian wild *Oryza* spp.: a review and conceptual framework for future research needs. *Agronomy* **12**, 42.
- Acquaah G (2015) Conventional plant breeding principles and techniques. In Al-Khayri JM, Jain SM and Johnson DV (Eds). *Advances in Plant Breeding Strategies: Breeding, Biotechnology and Molecular Tools*. Switzerland: Springer International Publishing, pp. 115–158.
- AgriFutures (2017) Desert Lime, AgriFutures. Available at <https://www.agrifutures.com.au/farm-diversity/desert-lime/>
- AgriFutures. (2022) “AgriFutures Australia. Available at <https://www.agrifutures.com.au/>”.
- Ahmed AK and Johnson KA (2000) Horticultural development of Australian native edible plants. *Australian Journal of Botany* **48**, 417–426.
- Ananda GK, Myrans H, Norton SL, Gleadow R, Furtado A and Henry RJ (2020) Wild sorghum as a promising resource for crop improvement. *Frontiers in Plant Science* **11**, 1108.
- Anderson MK (2013) *Tending the Wild: Native American Knowledge and the Management of California's Natural Resources*. Berkeley, CA: University of California Press.
- Antonelli A (2023) Indigenous knowledge is key to sustainable food systems. *Nature* **613**, 239–242.
- Ariati SR, Murphy DJ, Gardner S and Ladiges PY (2007) Morphological and genetic variation within the widespread species *Acacia victoriae* (Mimosaceae). *Australian Systematic Botany* **20**, 54–62.
- AVH (2023, August 2023) The Australasian Virtual Herbarium, Council of Heads of Australasian Herbaria. Available at <https://avh.chah.org.au>.
- Bartle J, Cooper D, Olsen G and Carslake J (2002) *Acacia* species as large-scale crop plants in the Western Australian wheatbelt. *Conservation Science Western Australia* **4**, 96–108.
- Bartlett ME, Moyers BT, Man J, Subramaniam B and Makunga NP (2023) The power and perils of *de novo* domestication using genome editing. *Annual Reviews of Plant Biology* **74**, 727–750.
- Bell LW, Wade LJ and Ewing MA (2010) Perennial wheat: a review of environmental and agronomic prospects for development in Australia. *Crop and Pasture Science* **61**, 679–690.
- Bell LW, Bennett RG, Ryan MH and Clarke H (2011) The potential of herbaceous native Australian legumes as grain crops: a review. *Renewable Agriculture and Food Systems* **26**, 72–91.
- Bell LW, Ryan MH, Bennett RG, Collins MT and Clarke HJ (2012) Growth, yield and seed composition of native Australian legumes with potential as grain crops. *Journal of the Science of Food and Agriculture* **92**, 1354–1361.
- Bentham J, Singh GM, Danaei G, Green R, Lin JK, Stevens GA, Farzadfar F, Bennett JE, Di Cesare M and Dangour AD (2020) Multidimensional characterization of global food supply from 1961 to 2013. *Nature food* **1**, 70–75.

- Bindon P** (1996) *Useful Bush Plants*. Perth, WA.: Western Australian Museum.
- Birch J, Benkendorff K, Liu L and Luke H** (2023) The nutritional composition of Australian native grains used by First Nations people and their re-emergence for human health and sustainable food systems. *Frontiers in Sustainable Food Systems* 7, 1237862.
- Blade SF and Slinkard AE** (2002) New crop development: the Canadian experience. In Janick J and Whipkey A (eds). *Trends in New Crops and New Uses*. Alexandria, VA: ASHS, pp. 62–75.
- Bohra A, Tiwari A, Kaur P, Ganie SA, Raza A, Roorkiwal M, Mir RR, Fernie AR, Smýkal P and Varshney RK** (2022) The key to the future lies in the past: insights from grain legume domestication and improvement should inform future breeding strategies. *Plant and Cell Physiology* 63, 1554–1572.
- Bowman DM** (1998) The impact of Aboriginal landscape burning on the Australian biota. *New Phytologist* 140, 385–410.
- Brand-Miller JC and Cherikoff V** (1985) Australian Aboriginal bushfoods: the nutritional composition of plants from arid and semi-arid areas. *Australian Aboriginal Studies* 2, 38–46.
- Broadhurst L and Coates D** (2017) Plant conservation in Australia: current directions and future challenges. *Plant Diversity* 39, 348–356.
- Broadhurst L, Breed M, Lowe A, Bragg J, Catullo R, Coates D, Encinas-Viso F, Gellie N, James E and Krauss S** (2017) Genetic diversity and structure of the Australian flora. *Diversity and Distributions* 23, 41–52.
- Brummer EC, Barber WT, Collier SM, Cox TS, Johnson R, Murray SC, Olsen RT, Pratt RC and Thro AM** (2011) Plant breeding for harmony between agriculture and the environment. *Frontiers in Ecology and the Environment* 9, 561–568.
- Burchfield EK, Nelson KS and Spangler K** (2019) The impact of agricultural landscape diversification on US crop production. *Agriculture, Ecosystems & Environment* 285, 106615.
- Canning AD** (2022) Rediscovering wild food to diversify production across Australia's agricultural landscapes. *Frontiers in Sustainable Food Systems* 6, 865580.
- Ceccarelli S** (2015) Efficiency of plant breeding. *Crop Science* 55, 87–97.
- Ceccarelli S and Grando S** (2020) Participatory plant breeding: who did it, who does it and where? *Experimental Agriculture* 56, 1–11.
- Chapman AD** (2009) Numbers of living species in Australia and the world. Report for the Australian Biological Resources Study. Canberra., The Australian Government Department of the Environment, Water, Heritage and the Arts.: 84.
- Cherikoff V** (2000) Marketing the Australian native food industry. Rural Industries Research and Development Corporation (RIRDC) 39.
- Cherikoff V and Brand J** (1983) Nutrient composition of Aboriginal bush foods. *Australian Aboriginal Studies* 2, 78–79.
- Cherikoff V and Brand JC** (1988) Is there a trend towards indigenous foods in Australia. Food habits in Australia: proceedings of the first Deakin/Sydney Universities symposium on Australian nutrition. A. S. Truswell and M. L. Wahlqvist. North Balwyn, Vic, University of Sydney. 178–183.
- Chivers I, Warrick R, Bomman J and Evans C** (2015) Native grasses make new products. A review of current and past uses and assessment of potential. Rural Industries Research and Development Corporation (RIRDC) 30.
- Clarke PA** (2011) *Aboriginal People and Their Plants*. Kenthurst, N.S.W.: Rosenberg Publishing.
- Clarke M** (2012) Australian native food industry stocktake. Rural Industries Research and Development Corporation (RIRDC) 89.
- Clarke M** (2013) Native foods R&D priorities and strategies 2013–2018. Rural Industries Research and Development Corporation (RIRDC) 53.
- CNFS** (2022) Creative Native Food Service. Available at <https://creativenativefoods.com.au/>
- Coates D, Byrne M, Cochrane A, Dunne C, Gibson N, Keighery G, Lambers H, Monks L, Thiele K and Yates C** (2014) Conservation of the kwongan flora: threats and challenges. In Lambers H (ed). *Plant Life on the Sandplains in Southwest Australia, a Global Biodiversity Hotspot*. Crawley, Western Australia: UWA Publishing, pp. 263–284.
- Colley M, Dawson J, McCluskey C, Myers J, Tracy W and van Bueren EL** (2021) Exploring the emergence of participatory plant breeding in countries of the Global North – a review. *The Journal of Agricultural Science* 159, 320–338.
- Collins M and Norton R** (2019) Diversifying the cropping phase. In Pratley JE and Kirkegaard JA (eds). *Australian Agriculture in 2020: From Conservation to Automation*. Wagga Wagga, N.S.W.: Australian Society for Agronomy, pp. 307–322.
- Colton B and Potter T** (1999) History. Canola in Australia – The First 30 Years. P. A. Salisbury, T. D. Potter, G. McDonald and A. G. Green, Organising Committee of 10th International Rapeseed Congress). Available at www.regional.org.au/au/gcirc/canola/: 1–4.
- Considine JA** (1996) Emerging Indigenous Crops of Australia. Progress in New Crops. J. Janick. Alexandria, VA. ASHS Press: p. 26–36.
- Coughlan MR and Nelson DR** (2018) Influences of native American land use on the colonial Euro-American settlement of the South Carolina Piedmont. *PLoS ONE* 13, e0195036.
- Cresswell I, Janke T and Johnston E** (2021) Australia state of the environment 2021: overview, independent report to the Australian Government Minister for the Environment, Commonwealth of Australia, Canberra: 274.
- Crews TE, Carton W and Olsson L** (2018) Is the future of agriculture perennial? Imperatives and opportunities to reinvent agriculture by shifting from annual monocultures to perennial polycultures. *Global Sustainability* 1, e11.
- CRRDC** (2016) Cross-RDC Impact Assessment and Performance Reporting Update. Stage 1: Cross-RDC Impact Assessment for the Period 1 July 2009 to 30 June 2015 The Council of Rural Research and Development Corporations. Submitted by Agrtrans Research, AgEconPlus, and EconSearch: 53.
- Davies CL, Waugh DL and Lefory EC** (2005) Variation in seed yield and its components in the Australian native grass *Microlaena stipoides* as a guide to its potential as a perennial grain crop. *Australian Journal of Agricultural Research* 56, 309–316.
- DCCEEW** (2021) Australian Government Department of Climate Change, Energy, the Environment and Water. The Nagoya Protocol – Convention on Biological Diversity. Available at <https://www.dcceew.gov.au/science-research/australias-biological-resources/nagoya-protocol-convention-biological> (Jan 2024).
- DeHaan LR, Van Tassel DL, Anderson JA, Asselin SR, Barnes R, Baute GJ, Cattani DJ, Culman SW, Dorn KM, Hulke BS, Kantar M, Larson S, Marks MD, Miller AJ, Poland J, Ravetta DA, Rude E, Ryan MR, Wyse D and Zhang X** (2016) Pipeline strategy for grain crop domestication. *Crop Science* 56, 917–930.
- Denham T and Donohue M** (2023) Putting the Dark Emu debate into context. *Archaeology in Oceania* 58, 275–295.
- DEWR** (2007) Australia's native vegetation: a summary of Australia's major vegetation groups. Canberra, ACT, Australian Government. Department of the Environment and Water Resources.
- Di Giovine MA and Brulotte RL** (2016) Food and foodways as cultural heritage. In Brulotte RL and Di Giovine MA (eds). *Edible Identities: Food as Cultural Heritage*. London: Routledge, pp. 1–27.
- Do C, Panakera-Thorpe L, Delaporte K and Schultz C** (2014) *Kunzea pomifera* (muntries): selection validation and evaluation of important horticultural traits. XXIX International Horticultural Congress on Horticulture: Sustaining Lives, Livelihoods and Landscapes (IHC2014): II 1117.
- Do CM, Delaporte KL, Pagay V and Schultz CJ** (2018a) Salinity tolerance of muntries (*Kunzea pomifera* F. Muell.), a native food crop in Australia. *HortScience* 53, 1562–1569.
- Do CM, Panakera-Thorpe LC, Delaporte KL, Croxford AE and Schultz CJ** (2018b) Genic simple sequence repeat markers for measuring genetic diversity in a native food crop: a case study of Australian *Kunzea pomifera* F. Muell.(muntries). *Genetic Resources and Crop Evolution* 65, 917–937.
- Drake A, Keitel C and Pattison A** (2021) The use of Australian native grains as a food: a review of research in a global grains context. *The Rangeland Journal* 43, 223–233.
- Ens E, Walsh F and Clarke P** (2017) Aboriginal people and Australia's vegetation: past and current interactions. In Keith DA (ed). *Australian Vegetation*. Cambridge, United Kingdom: Cambridge University Press, pp. 89–112.
- Fadda C, Mengistu DK, Kidane YG, Dell'Acqua M, Pè ME and Van Etten J** (2020) Integrating conventional and participatory crop improvement for smallholder agriculture using the Seeds for Needs Approach: a review. *Frontiers in Plant Science* 11, 559515.
- Fahey M, Rossetto M, Ens E and Ford A** (2022) Genomic screening to identify food trees potentially dispersed by precolonial indigenous peoples. *Genes* 13, 476.

- FAO (2017) *The Future of Food and Agriculture – Trends and Challenges*. Rome: Food and Agriculture Organization of the United Nations, 180.
- FAOSTAT (2022) <http://faostat3.fao.org>. Retrieved August 2022.
- Fletcher RJ (2002) International new crops development incentives, barriers, processes and progress: an Australian perspective. *Trends in New Crops and New Uses*. J. Janick and A. Whipkey. Alexandria, VA, ASHS: 15.
- FloraBase (2021) FloraBase: the Western Australian Flora. Available at <http://florabase.dec.wa.gov.au>, The Western Australian Department of Environment and Conservation.
- FloraNT (2021) FLoraNT. Northern Territory Flora Online. Available at <http://eflora.nt.gov.au/>, Northern Territory Government.
- Foster M (2014) Emerging animal and plant industries – their value to Australia. Rural Industries Research and Development Corporation (RIRDC) 194.
- Fuller DQ, Denham T and Allaby R (2023) Plant domestication and agricultural ecologies. *Current Biology* **33**, R636–R649.
- Fyfe S, Smyth HE, Schirra HJ, Rychlik M and Sultanbawa Y (2021) The framework for responsible research with Australian native plant foods: a food chemist's perspective. *Frontiers in Nutrition* **8**, 738627.
- Galanakis CM (2019) *Innovations in Traditional Foods*. Duxford, United Kingdom: Woodhead Publishing.
- Gasparini K, dos Reis Moreira J, Peres LEP and Zsögön A (2021) De novo domestication of wild species to create crops with increased resilience and nutritional value. *Current Opinion in Plant Biology* **60**, 102006.
- Gorst J (2002) Indigenous fruits of Australia. ISHS Acta Horticulturae 575: International Symposium on Tropical and Subtropical Fruits(575), 555–561.
- Hallam S (1989) Plant usage and management in Southwest Australian Aboriginal societies. In Harris DR and Hillman G (eds). *Foraging and Farming: The Evolution of Plant Exploitation*. London: Unwin Hyman Ltd, pp. 136–151.
- Hansen V and Horsfall J (2019) *Noongar Bush Tucker*. Perth, WA: UWA Publishing.
- Hatton T and Nulsen R (1999) Towards achieving functional ecosystem mimicry with respect to water cycling in southern Australian agriculture. *Agroforestry Systems* **45**, 203–214.
- Haygood R, Ives AR and Andow DA (2003) Consequences of recurrent gene flow from crops to wild relatives. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**, 1879–1886.
- Hele A (2001) Australian native citrus – wild species, cultivars and hybrids., Primary Industries and Resources South Australian Available at www.pir.sa.gov.au/factsheets: 7.
- Hele A (2002) Issues in the commercialisation of wattle seed for food. *Conservation Science Western Australia* **4**, 181–184.
- Henry RJ (2012) Next-generation sequencing for understanding and accelerating crop domestication. *Briefings in Functional Genomics* **11**, 51–56.
- Henry R (2019) Australian wild rice populations: a key resource for global food security. *Frontiers in Plant Science* **10**, 1354.
- Henry RJ (2023) Genomic characterization supporting the development of new food and crop options from the Australian flora. *Sustainable Food Technology* **1**, 337–347.
- Henry RJ, Rice N, Waters DL, Kasem S, Ishikawa R, Hao Y, Dillon S, Crayn D, Wing R and Vaughan D (2010) Australian *Oryza*: utility and conservation. *Rice* **3**, 235–241.
- Hobbs RJ and O'Connor M (1999) Designing mimics from incomplete data sets: salmon gum woodland and heathland ecosystems in South West Australia. *Agroforestry Systems* **45**, 365–394.
- Hobbs RJ and Yates CJ (2003) Impacts of ecosystem fragmentation on plant populations: generalising the idiosyncratic. *Australian Journal of Botany* **51**, 471–488.
- Hochman Z, Gobbett DL and Horan H (2017) Climate trends account for stalled wheat yields in Australia since 1990. *Global Change Biology* **23**, 2071–2081.
- Hopper SD and Gioia P (2004) The southwest Australian floristic region: evolution and conservation of a global hot spot of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **35**, 623–650.
- Howden SM, Gifford RG and Meinke H (2010) Grains. In Stokes C and Howden M (eds). *Adapting Agriculture to Climate Change: Preparing Australian Agriculture, Forestry and Fisheries for the Future*. Collingwood, Vic.: CSIRO publishing, pp. 36–70.
- Hunt J, Kirkegaard J, Celestina C and Porker K (2019) Transformational agronomy: restoring the role of agronomy in modern agricultural research. In Pratley JE and Kirkegaard J (eds), *Australian Agriculture in 2020: From Conservation to Automation*. Wagga Wagga, NSW, Australia: Agronomy Australia and Charles Sturt University, pp. 373–388.
- Hunt JR, Kirkegaard JA, Harris FA, Porker KD, Rattey AR, Collins MJ, Celestina C, Cann DJ, Hochman Z and Lilley JM (2021) Exploiting genotype×management interactions to increase rainfed crop production: a case study from south-eastern Australia. *Journal of Experimental Botany* **72**, 5189–5207.
- Hwang E-Y, Wei H, Schroeder SG, Fickus EW, Quigley CV, Elia P, Araya S, Dong F, Costa L and Ferreira ME (2019) Genetic diversity and phylogenetic relationships of annual and perennial *Glycine* species. *G3: Genes, Genomes, Genetics* **9**, 2325–2336.
- Isaacs J (1987) *Bush Food: Aboriginal Food and Herbal Medicine*. McMahons Point, NSW: Weldon Publishing.
- Isbell F, Adler PR, Eisenhauer N, Fornara D, Kimmel K, Kremen C, Letourneau DK, Liebman M, Polley HW and Quijas S (2017) Benefits of increasing plant diversity in sustainable agroecosystems. *Journal of Ecology* **105**, 871–879.
- Iverson AL, Marín LE, Ennis KK, Gonthier DJ, Connor-Barrie BT, Remfert JL, Cardinale BJ and Perfecto I (2014) Do polycultures promote win-wins or trade-offs in agricultural ecosystem services? A meta-analysis. *Journal of Applied Ecology* **51**, 1593–1602.
- Jacobsen S-E, Sorensen M, Pedersen SM and Weiner J (2015) Using our agrobiodiversity: plant-based solutions to feed the world. *Agronomy for Sustainable Development* **35**, 1217–1235.
- Janick J (ed.) (1996) *Progress in New Crops*. Alexandria, VA: ASHS Press.
- Janick J (1999) New crops and the search for new food resources. In Janick J (ed.), *Perspectives on New Crops and New Uses*. Alexandria, VA: ASHS Press, pp. 104–110.
- Janick J and Whipkey A (eds) (2002) *Trends in New Crops and New Uses*. Alexandria, VA: ASHS Press.
- Janick J and Whipkey A (eds) (2007) *Issues in New Crops and New Uses*. Alexandria, VA: ASHS Press.
- Janick J, Blase MG, Johnson DL, Jolliff GD and Myers RL (1996) Diversifying U.S. crop production. CAST Issue Paper 6. Ames, Iowa., Council of Agric. Sci. and Tech. 98–109.
- Jarvis D, Maclean K and Woodward E (2021) The Australian Indigenous-led bush products sector: insights from the literature and recommendations for the future. *Ambio* **51**, 226–240.
- Johnson K and Burchett M (eds) (1996) *Native Australian Plants: Horticulture and Uses*. Sydney: UNSW Press.
- Jolliff GD (1990) Strategic planning for new-crop development. In Janick J and Simon JE (eds), *Advances in New Crops*. Portland, OR: Timber Press, pp. 29–40.
- Kahane R, Hodgkin T, Jaenicke H, Hoogendoorn C, Hermann M, Keatinge JDHD, Hughes JdA, Padulosi S and Looney N (2013) Agrobiodiversity for food security, health and income. *Agronomy for Sustainable Development* **33**, 671–693.
- Keating B and Carberry P (2010) Emerging opportunities and challenges for Australian broadacre agriculture. *Crop and Pasture Science* **61**, 269–278.
- Keen I (2021) Foragers or farmers: dark emu and the controversy over Aboriginal agriculture. *Anthropological Forum* **31**, 106–128.
- Khoddami A, Drake A, Pattison A, Craige C, Badaoui C, Keitel C, Roth G, Leung H, Lee JH, Cross R, Phillips S and Bell T (2020) *Native Grains from Paddock to Plate*. University of Sydney, Institute of Agriculture, 40.
- Khoury CK, Bjorkman AD, Dempewolf H, Ramirez-Villegas J, Guarino L, Jarvis A, Rieseberg LH and Struik PC (2014) Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences* **111**, 4001–4006.
- Konczak I, Zabarar D, Dunstan M, Aguas P, Roulfe P and Pavan A (2009) Health benefits of Australian native foods – an evaluation of health-enhancing compounds, Rural Industries Research and Development Corporation (RIRDC) 52.
- Kumar Y, Basu S, Goswami D, Devi M, Shivhare US and Vishwakarma RK (2022) Anti-nutritional compounds in pulses: implications and alleviation methods. *Legume Science* **4**, e111.

- Lacoste M, Cook S, McNee M, Gale D, Ingram J, Bellon-Maurel V, MacMillan T, Sylvester-Bradley R, Kindred D, Bramley R, Tremblay N, Longchamps L, Thompson L, Ruiz J, García FO, Maxwell B, Griffin T, Oberthür T, Huyghe C, Zhang W, McNamara J and Hall A (2022) On-farm experimentation to transform global agriculture. *Nature Food* 3, 11–18.
- Latz P (1995) *Bushfires and Bushucker: Aboriginal Plant Use in Central Australia*. Alice Springs, NT: IAD Press.
- Laurie S (2020) Australian native foods and botanicals – 2019/20 market study. Australian Native Foods and Botanicals. The University of Sydney. 56.
- Lawn RJ (2015) The Australian Vigna species: a case study in the collection and conservation of crop wild relatives. In Redden RJ, Yadav SS, Maxted N, Dulloo ME, Guarino L and Smith P (eds). *Crop Wild Relatives and Climate Change*. Hoboken, NJ: John Wiley & Sons, Incorporated, pp. 318–355.
- Lee LS (2013) Horticultural development of bush food plants and rights of Indigenous people as traditional custodians – the Australian bush tomato (*Solanum centrale*) example: a review. *The Rangeland Journal* 34, 359–373.
- Lee LS and Courtenay K (2016) Enrichment plantings as a means of enhanced bush food and bush medicine plant production in remote arid regions: a review and status report. Learning Communities. International Journal of Learning in Social Contexts. Special Issue: Synthesis & Integration Writing Forum. R. Wallace, Charles Darwin University. The Cooperative Research Centre for Remote Economic Participation. Number 19: 64–75.
- Lee JH and Six J (2010) Effect of climate change on field crop production and greenhouse gas emissions in the California's Central Valley. Proceedings of the 19th World Congress of Soil Science: Soil solutions for a changing world. 1–6 August 2010. R. Gilkes and N. Prakongkep. Brisbane, Australia., Australian Society of Soil Science Inc.
- Leha D, Cubillo C and Janke T (2019) IP food for thought: intellectual property and protecting indigenous bush foods. Merinj Kaardijn: Aboriginal Food Knowledge Forum. Albany, WA, 22 & 23 November 2019: 14.
- Levis C, Costa FR, Bongers F, Peña-Claros M, Clement CR, Junqueira AB, Neves EG, Tamanaha EK, Figueiredo FO and Salomão RP (2017) Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science* 355, 925–931.
- Levis C, Flores BM, Moreira PA, Luize BG, Alves RP, Franco-Moraes J, Lins J, Konings E, Peña-Claros M and Bongers F (2018) How people domesticated Amazonian forests. *Frontiers in Ecology and Evolution* 5, 171.
- Li C, Stomph T-J, Makowski D, Li H, Zhang C, Zhang F and van der Werf W (2023) The productive performance of intercropping. *Proceedings of the National Academy of Sciences* 120, e2201886120.
- Lightfoot KG, Cuthrell RQ, Striplen CJ and Hylkema MG (2013) Rethinking the study of landscape management practices among hunter-gatherers in North America. *American Antiquity* 78, 285–301.
- Lin BB (2011) Resilience in agriculture through crop diversification: adaptive management for environmental change. *BioScience* 61, 183–193.
- Lingard K and Martin P (2016) Strategies to support the interests of Aboriginal and Torres Strait Islander peoples in the commercial development of gourmet bush food products. *International Journal of Cultural Property* 23, 33–70.
- Lister PR, Holford P, Haigh T and Morrison DA (1996) Acacia in Australia: ethnobotany and potential food crop. Progress in new crops. J. Janick. Alexandria, VA., ASHS Press: p. 228–236.
- Loomis RS (2022) Perils of production with perennial polycultures. *Outlook on Agriculture* 51, 22–31.
- Low T (1991) *Wild Food Plants of Australia*. North Ryde, N.S.W.: Angus and Robertson.
- Lullfitz A, Byrne M, Knapp L and Hopper SD (2020a) *Platysace* (Apiaceae) of south-western Australia: silent story tellers of an ancient human landscape. *Biological Journal of the Linnean Society* 130, 61–78.
- Lullfitz A, Dabb A, Reynolds R, Knapp L, Pettersen C and Hopper SD (2020b) Contemporary distribution of *Macrozamia dyeri* (Zamiaceae) is correlated with patterns of Nyungar occupation in south-east coastal Western Australia. *Austral Ecology* 45, 933–947.
- Luo G, Najafi J, Correia PM, Trinh MDL, Chapman EA, Østerberg JT, Thomsen HC, Pedas PR, Larson S and Gao C (2022) Accelerated domestication of new crops: yield is key. *Plant and Cell Physiology* 63, 1624–1640.
- Maclean K, Woodward E, Jarvis D, Turpin G, Rowland D and Rist P (2022) Decolonising knowledge co-production: examining the role of positionality and partnerships to support Indigenous-led bush product enterprises in northern Australia. *Sustainability Science* 17, 333–350.
- Maiden JH (1889) *The Useful Native Plants of Australia: (including Tasmania)*. Sydney, N.S.W.: Turner and Henderson.
- Martin AR, Cadotte MW, Isaac ME, Milla R, Vile D and Violle C (2019) Regional and global shifts in crop diversity through the Anthropocene. *PLoS ONE* 14, e0209788.
- Maslin BR, Thomson LAJ, McDonald BW and Hamilton-Brown S (1998) Edible wattle seeds of Southern Australia. A review of species for use in semi-arid regions. Perth, Australia, CSIRO Australia.
- Massawe F, Mayes S and Cheng A (2016) Crop diversity: an unexploited treasure trove for food security. *Trends in Plant Science* 21, 365–368.
- Meyer RS, DuVal AE and Jensen HR (2012) Patterns and processes in crop domestication: an historical review and quantitative analysis of 203 global food crops. *New Phytologist* 196, 29–48.
- Midgley S and Turnbull J (2003) Domestication and use of Australian acacias: case studies of five important species. *Australian Systematic Botany* 16, 89–102.
- Miers G (2004) Cultivation and sustainable wild harvest of bushfoods by Aboriginal Communities in Central Australia, Rural Industries Research and Development Corporation (RIRDC) 79.
- Mitchell M, Stodart B and Virgona J (2015) Genetic diversity within a population of *Microlaena stipoides*, as revealed by AFLP markers. *Australian Journal of Botany* 62, 580–586.
- Murphy DJ (2007) *People, Plants and Genes*. New York: Oxford University Press.
- Mustafa MA, Mayes S and Massawe F (2019) Crop diversification through a wider use of underutilised crops: a strategy to ensure food and nutrition security in the face of climate change. In Sarkar A, Sensarma S and vanLoon G (eds), *Sustainable Solutions for Food Security*. Cham: Springer, pp. 125–149.
- N'Danikou S and Tchokponhoue DA (2019) Plant domestication for enhanced food security. In Leal Filho W, Azul A, Brandli L, Özuyar P and Wall T (eds). *Zero Hunger*. Encyclopedia of the UN Sustainable Development Goals. New York, NY: Springer, Cham, pp. 644–654.
- Nelson P and Hawthorne WA (2000) Development of lupins as a crop in Australia. Linking Research and Marketing Opportunities for Pulses in the 21st Century. Current Plant Science and Biotechnology in Agriculture, vol 34. K. R., Springer: 549–559.
- NLE (2022) Yoordaning-bah, Coming Together., Noongar Land Enterprise Group. Food Innovation Australia Ltd. (FIAL): 31.
- Norton SL, Khoury CK, Sosa CC, Castañeda-Álvarez NP, Achicanoy HA and Sotelo S (2017) Priorities for enhancing the ex situ conservation and use of Australian crop wild relatives. *Australian Journal of Botany* 65, 638–645.
- Østerberg JT, Xiang W, Olsen LI, Edenbrandt AK, Vedel SE, Christiansen A, Landes X, Andersen MM, Pagh P and Sandøe P (2017) Accelerating the domestication of new crops: feasibility and approaches. *Trends in Plant Science* 22, 373–384.
- Page T (2004) *Muntries: The Domestication and Improvement of Kunzea Pomifera*. Canberra, ACT: Rural Industries Research and Development Corporation, 95.
- Page T, Moore G, Will J and Halloran G (2006a) Onset and duration of stigma receptivity in *Kunzea pomifera* (Myrtaceae). *Australian Journal of Botany* 54, 559–563.
- Page T, Moore G, Will J and Halloran G (2006b) Pollen viability in *Kunzea pomifera* (Myrtaceae) as influenced by sucrose concentration and storage. *Australian Journal of Botany* 54, 553–558.
- Pascoe B (2014) *Dark Emu Black Seeds: Agriculture or Accident?*. Broome, Western Australia: Magabala Books.
- Passioura JB (2020) Translational research in agriculture. Can we do it better? *Crop and Pasture Science* 71, 517–528.
- Pate J and Bell T (1999) Application of the ecosystem mimic concept to the species-rich Banksia woodlands of Western Australia. *Agroforestry Systems* 45, 303–341.
- Pavlik BM, Louderback LA, Vernon KB, Yaworsky PM, Wilson C, Clifford A and Coddling BF (2021) Plant species richness at

- archaeological sites suggests ecological legacy of Indigenous subsistence on the Colorado Plateau. *Proceedings of the National Academy of Sciences* **118**, e2025047118.
- Petersen B and Snapp S** (2015) What is sustainable intensification? Views from experts. *Land Use Policy* **46**, 1–10.
- Pratley JE and Kirkegaard J** (eds) (2019) Australian Agriculture in 2020: from conservation to automation. Available at <https://www.csu.edu.au/research/grahamcentre/publications/e-books/australian-agriculture-in-2020>, Graham Center for Agricultural Innovation.
- Pretty J, Benton TG, Bharucha ZP, Dicks LV, Flora CB, Godfray HCJ, Goulson D, Hartley S, Lampkin N and Morris C** (2018) Global assessment of agricultural system redesign for sustainable intensification. *Nature Sustainability* **1**, 441–446.
- Rangan H, Bell KL, Baum DA, Fowler R, McConvell P, Saunders T, Spronck S, Kull CA and Murphy DJ** (2015) New genetic and linguistic analyses show ancient human influence on baobab evolution and distribution in Australia. *PLoS ONE* **10**, e0119758.
- Rapport EH and Drausal BS** (2013) Edible plants. In Scheiner SM (Ed), *Encyclopedia of Biodiversity*, Vol. 3. Minneapolis, MN: Elsevier, pp. 127–132.
- Rebetzke G, Ingvordsen C, Bovill W, Trethowan R and Fletcher A** (2019) Breeding evolution for conservation agriculture. In Pratley JE and Kirkegaard JA (eds). *Australian Agriculture in 2020: From Conservation to Automation*. Wagga Wagga, N.S.W.: Australian Society for Agronomy, pp. 273–287.
- Renny-Byfield S, Page JT, Udall JA, Sanders WS, Peterson DG, Arick MA, Grover CE and Wendel JF** (2016) Independent domestication of two old world cotton species. *Genome Biology and Evolution* **8**, 1940–1947.
- Rickards L and Howden SM** (2012) Transformational adaptation: agriculture and climate change. *Crop and Pasture Science* **63**, 240–250.
- Rinaudo A and Cunningham P** (2008) Australian acacias as multi-purpose agro-forestry species for semi-arid regions of Africa. *Muelleria* **26**, 79–85.
- Rinaudo A, Patel P and Thomson L** (2002) Potential of Australian Acacias in combating hunger in semi-arid lands. *Conservation Science Western Australia* **4**, 161–169.
- RIRDC** (2008) Native foods R&D priorities and strategies 2007–2012. Rural Industries Research and Development Corporation (RIRDC) 24.
- Rivett D, Tucker D and Jones G** (1983) The chemical composition of seeds from some Australian plants. *Australian Journal of Agricultural Research* **34**, 427–432.
- Rotili DH, de Voil P, Eyre J, Serafin L, Aisthorpe D, Maddonni GÁ and Rodriguez D** (2020) Untangling genotype x management interactions in multi-environment on-farm experimentation. *Field Crops Research* **255**, 107900.
- Runk BC, Kantar MB, Jordan NR, Anderson JA, Wyse DL, Eckberg JO, Barnes RJ, Lehman CL, DeHaan LR and Stupar RM** (2014) The reflective plant breeding paradigm: a robust system of germplasm development to support strategic diversification of agroecosystems. *Crop Science* **54**, 1939–1948.
- Ryan M, Bell L, Bennett R, Collins M and Clarke H** (2011). Native legumes as a grain crop for diversification in Australia. Rural Industries Research and Development Corporation (RIRDC) 68.
- Salisbury PA, Cowling WA and Potter TD** (2016) Continuing innovation in Australian canola breeding. *Crop & Pasture Science* **67**, 266–272.
- Salvin S, Bourke M, Byrne AM and Byrne T** (eds) (2004) *The New Crop Industries Handbook*. Barton, ACT: Australian Government. Rural Industries Research and Development Corporation.
- Shapter FM and Chivers I** (2015) Commercialisation of elite lines of *Microlaena stipoides* as a perennial grain. Rural Industries Research and Development Corporation (RIRDC) 38.
- Shapter FM, Cross M, Ablett G, Malory S, Chivers IH, King GJ and Henry RJ** (2013) High-throughput sequencing and mutagenesis to accelerate the domestication of *Microlaena stipoides* as a new food crop. *PLoS ONE* **8**, e82641.
- Shelef O, Weisberg PJ and Provenza FD** (2017) The value of native plants and local production in an era of global agriculture. *Frontiers in Plant Science* **8**, 2069.
- Shelton AC, Tracy WF, Kapuscinski AR and Locke KA** (2016) Participatory plant breeding and organic agriculture: a synergistic model for organic variety development in the United States participatory plant breeding and organic agriculture: a synergistic model. *Elementa: Science of the Anthropocene* **4**, 000143.
- Sherman B and Henry RJ** (2020) The Nagoya Protocol and historical collections of plants. *Nature Plants* **6**, 430–432.
- Shigeura G and Ooka H** (1984) *Macadamia Nuts in Hawaii: History and Production*. Hawaii: University of Hawaii, 13.
- Silcock J** (2018) Aboriginal translocations: the intentional propagation and dispersal of plants in Aboriginal Australia. *Journal of Ethnobiology* **38**, 390–405.
- Smith BD** (2011) General patterns of niche construction and the management of ‘wild’ plant and animal resources by small-scale pre-industrial societies. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 836–848.
- Smykal P, Nelson MN, Berger JD and Von Wettberg EJ** (2018) The impact of genetic changes during crop domestication. *Agronomy* **8**, 119.
- Snapp SS, DeDecker J and Davis AS** (2019) Farmer participatory research advances sustainable agriculture: lessons from Michigan and Malawi. *Agronomy Journal* **111**, 2681–2691.
- Snowball R, Norman H and D’Antuono M** (2021) Investigation of two native Australian perennial forage legumes for their potential use in agriculture: *Indigofera australis* subsp. *hesperia* and *Glycyrrhiza acanthocarpa*. *Crop and Pasture Science* **72**, 311–323.
- Stalker HT, Warburton ML and Harlan JR** (2021) *Harlan’s Crops and Man: People, Plants and Their Domestication*. Hoboken, NJ: John Wiley & Sons.
- Stetter MG, Gates DJ, Mei W and Ross-Ibarra J** (2017) How to make a domesticate. *Current Biology* **27**, R896–R900.
- Stynes B** (1997) Opportunities for contributing to the development of Aboriginal food plants. *Tropical Grasslands* **31**, 311–314.
- Sudmeyer R, Edward A, Fazakerley V, Simpkin L and Foster I** (2016) *Climate Change: Impacts and Adaptation for Agriculture in Western Australia*. Perth, Western Australia: Department of Agriculture and Food, 177.
- Sultanbawa Y and Sultanbawa F** (eds) (2016) Australian native plants: cultivation and uses in the health and food industries. In *Traditional Herbal Medicines for Modern Times*. Boca Raton, Florida: CRC Press, p. 409.
- Sutton P and Walshe K** (2021) *Farmers or Hunter-Gatherers?: The Dark Emu Debate*. Melbourne, V.c.: Melbourne Univ. Publishing.
- Sykes S** (1997) Australian native limes (*Eremocitrus* and *Microcitrus*); a citrus breeder’s viewpoint. *Australian Bush Foods Magazine* **3**, 12–15.
- Thompson JC, Wright DK and Ivory SJ** (2021a) The emergence and intensification of early hunter-gatherer niche construction. *Evolutionary Anthropology: Issues, News, and Reviews* **30**, 17–27.
- Thompson JC, Wright DK, Ivory SJ, Choi J-H, Nightingale S, Mackay A, Schilt F, Otárola-Castillo E, Mercader J and Forman SL** (2021b) Early human impacts and ecosystem reorganization in southern-central Africa. *Science Advances* **7**, eabf9776.
- Tiwari BK, Gowen A and McKenna B** (eds) (2011) *Pulse Foods: Processing, Quality and Nutraceutical Applications*. Sydney, N.S.W.: Academic Press.
- Toensmeier E** (2016) *The Carbon Farming Solution: A Global Toolkit of Perennial Crops and Regenerative Agriculture Practices for Climate Change Mitigation and Food Security*. Chelsea, Vermont: Chelsea Green Publishing.
- Tucker Bush** (2022) Tucker Bush. Edible Australian Tucker Bush Available at <https://tuckerbush.com.au/about-us/>
- Van Tassel DL, Tesdell O, Schlautman B, Rubin MJ, DeHaan LR, Crews TE and Streit Krug A** (2020) New food crop domestication in the age of gene editing: genetic, agronomic and cultural change remain co-evolutionarily entangled. *Frontiers in Plant Science* **11**, 789.
- Walters J, Milne R and Thompson H** (2018) Online farm trials: a national web-based information source for Australian grains research, development and extension. *Rural Extension and Innovation Systems Journal* **14**, 117–123.
- Wang M, Yu Y, Haberer G, Marri PR, Fan C, Goicoechea JL, Zuccolo A, Song X, Kudrna D and Ammiraju JS** (2014) The genome sequence of African rice (*Oryza glaberrima*) and evidence for independent domestication. *Nature genetics* **46**, 982–988.
- Williams KA** (2005) An overview of the US National Plant Germplasm System’s exploration program. *HortScience* **40**, 297–301.

- Wilson RF** (2007) Strategies for narrowing the gap between R&D and commercialization of new crops. In Janick J and Whipkey A (eds), *Issues in New Crops and New Uses*. Alexandria, VA: ASHS Press, pp. 4–6.
- Winterhalder B and Kennett DJ** (2006) Behavioral ecology and the transition from hunting and gathering to agriculture. In Kennett DJ and Winterhalder B (eds), *Behavioral Ecology and the Transition to Agriculture*. Berkeley, CA: University of California Press, pp. 1–21.
- Wollenweber B, Porter JR and Lübberstedt T** (2005) Need for multidisciplinary research towards a second green revolution. *Current Opinion in Plant Biology* **8**, 337–341.
- Wu J, Wang Y, Xu J, Korban SS, Fei Z, Tao S, Ming R, Tai S, Khan AM and Postman JD** (2018) Diversification and independent domestication of Asian and European pears. *Genome Biology* **19**, 1–16.
- Yen DE** (1993) The origins of subsistence agriculture in Oceania and the potentials for future tropical food crops. *Economic Botany* **47**, 3–14.
- Zaidi SS-e-A, Vanderschuren H, Qaim M, Mahfouz MM, Kohli A, Mansoor S and Tester M** (2019) New plant breeding technologies for food security. *Science* **363**, 1390–1391.
- Zeder MA** (2015) Core questions in domestication research. *Proceedings of the National Academy of Sciences* **112**, 3191–3198.
- Zhang H, Mascher M, Abbo S and Jayakodi M** (2022) Advancing grain legumes domestication and evolution studies with genomics. *Plant and Cell Physiology* **63**, 1540–1553.
- Zimmerer KS and De Haan S** (2017) Agrobiodiversity and a sustainable food future. *Nature Plants* **3**, 1–3.