

- 1996;128:862-863. Letter.
4. Rubin LG, Tucci V, Cercenado E, Eliopoulos G, Isenberg HD. Vancomycin-resistant *Enterococcus faecium* in hospitalized children. *Infect Control Hosp Epidemiol* 1992;13:700-705.
 5. Fletcher CV, Giese RM, Rodman JH. Pharmacist interventions to improve prescribing of vancomycin and tobramycin. *Am J Hosp Pharm* 1986;43:2198-2202.
 6. McCormack JP, Lynd LD, Pfeifer NM. Vancomycin cost containment through a therapeutic and pharmacokinetic drug monitoring service. *Canadian Journal of Hospital Pharmacy* 1986;42:3-9.
 7. Misan GMH, Martin ED, Smith ER, Somogyi AA, Bartholomeusz RCA, Bochner F. Drug utilization review in a teaching hospital: experience with vancomycin. *Eur J Clin Pharmacol* 1990;39:457-461.
 8. Johnson SV, Hoey LL, Vance-Bryan K. Inappropriate vancomycin prescribing based on criteria from the Centers for Disease Control and Prevention. *Pharmacotherapy* 1995;15:579-585.
 9. Evans ME, Kortes KJ. Vancomycin use in a university medical center: comparison with Hospital Infection Control Practices Advisory Committee guidelines. *Infect Control Hosp Epidemiol* 1996;17:356-359.
 10. Logsdon BA, Lee KR, Barrett FE. Correct dosing of vancomycin in infants and children. *Pediatrics* 1995;96:1177. Letter.

Outbreak of *Mycoplasma Pneumonia*

Researchers from the CDC and the US Merchant Marine Academy reported an outbreak of *Mycoplasma pneumoniae* infection among students at a Federal Service Academy. During a 2-month outbreak, 164 of 233 (70%) of freshmen and 154 of 354 (44%) upperclassmen questioned reported having a respiratory illness consistent with *M pneumoniae* infection. Of the 199 ill students who had complete laboratory evaluation, 65 (55%) had laboratory evidence of *M pneumoniae* infection. The first phase of the outbreak involved freshmen;

the second, occurring 2 to 3 weeks later, involved predominantly upperclassmen, who had arrived on campus 2 weeks after the freshmen. In a multivariate analysis of the first phase, ill students were more likely than well ones to have multiple visits to the academy's health clinic before the onset of respiratory illness. A case-control study of freshman visiting the clinic showed that transmission was more likely when more than six people were present in the waiting room. The authors concluded that crowding in the health clinic was

a significant risk factor for *M pneumoniae* infection in this outbreak. A separate waiting room for patients with respiratory symptoms during an outbreak of *M pneumoniae* may limit transmission.

FROM: Feikin D, Moroney J, Talkington D, et al. New implications for institutional outbreaks of *Mycoplasma pneumoniae*. Presented at the 35th Annual Meeting of the Infectious Disease Society of America; September 14-16, 1997; San Francisco, California. Abstract 37.