VOL. XII.—1897.

THE

JOURNAL OF LARYNGOLOGY,

RHINOLOGY, AND OTOLOGY;

AN ANALYTICAL RECORD OF CURRENT LITERATURE

RELATING TO

THE THROAT, NOSE, AND EAR.

PUBLISHED MONTHLY.

Kondon:

THE REBMAN PUBLISHING COMPANY, LIMITED,

SUCCESSORS TO F. J. REBMAN, 11, ADAM STREET, STRAND.

AMERICAN AGENTS:

THE F. A. DAVIS COMPANY, 1914-16, CHERRY STREET, PHILADELPHIA, PA.

ENTERED AT STATIONERS' HALL.

THE JOURNAL OF LARYNGOLOGY, RHINOLOGY, AND OTOLOGY.

Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN

EDITORS:

JOHN MACINTYRE, M.B., F.R.S.E.,
DUNDAS GRANT, M.A., M.D., F.R.C.S.Eng.,
ARTHUR SANDFORD, M.D., M.Ch.R.U.I.

W. MILLIGAN, M.D. (Manchester), Sub-Ed.
R. LAKE, F.R.C.S. (London), Managing Sub-Ed.

WITH THE CO-OPERATION OF

DRS. BARON (Bristol), BETTMAN (Rome), BOTEY (Barcelona),

CARTAZ (Paris), BRYSON DELAYAN (New York), DODD (Chicago),

DRAISPUL (St. Petersburg), GRAZZI (Florence), GUYE (Amsterdam),

HICGUET (Brussels), MIDDLEMASS HUNT (Liverfool), HUTCHISON (Glaszow),

JOAL (Paris), KARWOWSKI (Warsaw), KELLY (Glasgow),

LACOARRET (Toulouse), LICHTWITZ (Bordeaux),

LIEVEN (Aix-la-Chapelle), PROF. JOHN N. MACKENZIE (Baltimore),

PROF. MASSEI (Naples), MEYJES (Amsterdam), E. MEYER (Berlin),

HOLGER MYGIND (Copenhagen), PORTER (St. Louis),

STGEORGE REID (London), RUEDA (Madrid), SAJOUS (Paris),

SENDZIAK (Warsaw), SOTA (Seville), STCLAIR THOMSON (London),

WAGGETT (London), WOODS (Dublin), AND ZIEM (Pantzic).

CONTENTS.

Original Articles— Papilloma of Larynx (illustrated). By J. McNbill Whistler, M.D
Societies' Meetings— LARYNGOLOGICAL SOCIETY OF LONDON.—Laryngectomy, Spencer—Ulcerative Lacunar Tonsillitis, Lake—Cleft Palate and Adenoids, Lave—Cyst of Naso-Pharynx, Bond—Membranous Laryngitis, Bond—Recurrent Laryngeal Growth, Bond—Syringomyelia, with Paralysis of Palate and Vocal Cord, Horne—Cancer of Esophagus and Trachea, Beale—Unilateral Paralysis and Displacement of Arytenoid, Waggett—Chronic Laryngitis after Typhoid, etc., Thomson—Pages 426 to 433
New York Academy of Medicine.—Obscure Laryngeal Disease, <i>Phillips</i> .—Frontal Sinus Disease, <i>Myles</i> .—Specific Adhesions of Soft Palate, <i>Berens</i> .—Tumour of Ventricle, <i>Gleitsmann</i> .—Nasal Saddle, <i>Colien</i> .—Sand Figures, <i>Curtis</i> .—Adenoids and Growth of the Palate, <i>Cleitsmann</i>
AMERICAN LARVNGOLOGICAL ASSOCIATION.—Tumour of Nose, Bryan — Frontal Sinus Disease, Bryan —Atrophic Rhinitis, Casselberry — Simulated Sarcoma of Tonsil, Delavan — Enlargements of Inferior Turbinateds, Delavan — Subglottic Tumour, Farlow — Edematous Nasal Polypi and Adenomata, Wright — Lupus of Larynx, Mayer — Laryngeal Phthisis, Murray — Nasal Sarcoma, Nichols. Pages 433 to 445
Austrian Otological Society.—Epidemic Meningitis and Suppurative Ottis, Alt——Symmetrical Exostoses, Politzer——Atypical Mastoid Operation, Kaufmann—Paracentesis and Vaseline Injections in Plastic Ottis, Kaufmann——Hyperostoses, Gruber—Discussion: Hydrogen PeroxidePages 445 to 449
Deutschen Gesellschaft für Chirurgie.—Retained Intubation Tube, König——Plastic Operations, Larynx and Trachea, König, jun.—Laryngectomy, Graf——Carcinoma of the Pharynx, Krönlein——Lymphatics and Carcinoma of Tongue, Kuttner——Hare Lip, Samter
Für Aerzte in Zürich.—Sinusitis, Laubi, 452 to 455
Editorial— Tuning Fork Tests Page 455 Jubilee Honours , 457
Abstracts— Diphtheria.—Antitoxin, Brown, Slater, Cameron, "Lancet"—Avian, Ferré.—Umbilical, Pitts— Subcutaneous Emphysema, Sharman and D'Esterre
Mouth.—Exaggerated Mobility, Bourdette—Tonsillar Lympho-Adenitis, Geronti—Tonsillitis and Milk, Grey-Edwards—Acute Tonsillitis, Sepsis, and Death, Haddaeus—Artificial Alimentary Canal, Jaboulay—Retro-Pharyngeal Abscess, Munger—Actinowycosis, Ququet—Acute Macroglossitis, Sabrazes—Tuberculosis of Tonsils, Somers. Pages 460 to 462
Nose.—Tri-Valve Speculum, Armstrong.—Nose in Scarlet Fever, Clark.—Frontal Sinus Anatomy, Collier.—Polypi of Choanæ, Gaudier.—Meyer's Tonsil, Gourc.—Ozeana, Gradenigo.—Frontal Sinusitis, Gradenigo.—Nasal Reflex, Hobbs.—Fibro-Sarcoma, Nose, Hubbard.—Congenital Occlusion Right Naris, Kugh.—Direct Inspection of Meyer's Tonsil, Lindt.—Frontal Empyema, Photiades.—Professional Rhinitis, Prota.—Atrophic Rhinitis, Rhodes.—Deformed Nose, Roe.—Infantile Nasal Atresia, Stout.—Pages 452 to 467.
Larynx.—Cancer of Larynx, Ardonin.—Airol, Fasano.—Laryngeal Stenosis, Fischer.—Laryngeal Tuberculosis, Goris.—Hæmorthage after Tracheotomy, Hecker.—Combined Laryngoscopy, Kirstein.—Recurrent Paralysis, Lermoyez.—Amyloid Neoplasm, Martuscelli.—Laryngeal Tuberculosis, Massei.—Epithelioma, Phillips.—Laryngocele Ventriculaire, Rangé.—Cancer, Schmiegelow.—Artificial Larynx, Stuart.—Paralysis, Wallenberg
Ear. —Pyogenic Brain Disease, Baker—New Treatment, Cohen-Kysper—Conducting Portion of Labyrinth, Cheatle—Pyæmia, Eulenstein—Surgical Treatment of Deafness, Garnault—Teaching the Deaf, Goldstein—Tubercle of Middle Ear, Haenel—Sinus Thrombosis, Hoffman—Cerumen, Lannois Pages 471 to 474
Reviews— Fages 471 to 474 Love.—Deaf-Mutism Page 474 Manhattan Reports " 475
New Preparations, 476
GUIDE TO ADVERTISEMENTS.
Bullock & Co
Burroughs, Wellcome & Co

	PAGE
Bullock & Co	vii.
Burroughs, Wellcome & Co	vi.
Central London Throat Hospital	xiv.
Globe Manufacturing Company	x.
Ingram & Royle	vii xi.
Globe Manufacturing Company Ingram & Royle Lambert Pharmacal Co.	ii.
Mayer & Meltzer	xii .
Pearl Life Assurance Company	x
Pearl Life Assurance Company	i., ix, x
Slater, Bros. & Crossley	iv.
Woodruff	νii.

"TABLOID" and "SOLOID" MEDICATION IN LARYNGOLOGY.

A "TABLOID," when intended for the treatment of the throat and voice, is compressed hard so as to dissolve slowly and produce a prolonged action on the mucous membrane. There are no sugary or gummy substances present to interfere with the direct application of the medicament to the affected surfaces. Placed upon the tongue, a "Tabloid" is immediately acted upon by the saliva forming a highly-concentrated solution, which remains in continuous contact with the affected mucous membrane until the whole of the "Tabloid" is dissolved.

"TABLOID" RED GUM.



Red Gum, an exudation from the Australian *Eucalyptus rostrata*, is a mild styptic and antiseptic, and an excellent local and general astringent. The *Practitioner* reports:—"When a Red Gum 'Tabloid' is allowed to dissolve in the mouth, the continuous action of the drug on the throat is fully secured. Great care has been exercised so to adjust the relative portions of the ingredients that the full therapeutic effect of the drug may be obtained without offending the palate or affecting the appetite."

Supplied in boxes of 30 and bottles of 100, at 5d. and 1s. each.

"TABLOID" TAR.

In a paper, entitled "A Mode of Administering Tar," which appeared in the *Medical Press and Circular*, a distinguished physician reports as follows:—

"I have recently had my attention called to 'Tabloids' of Tar, containing one grain; and these, I think, constitute by far the best mode of administering the drug. They contain the whole of the constituents of tar and are palatable and speedily disintegrate. The new mode of administration is three or four every four hours; but a better plan is to suck one frequently—in fact, every time the cough is troublesome."

Supplied in bottles of 50 and 100 at 5d. and 9d. each.

"SOLOID" EUCAINE HYDROCHLORIDE. (Gr. 1 and Gr. 5.

Eucaine is a synthetic compound allied to cocaine and having a similar chemical composition. Its local anæsthetic action is not unlike that of cocaine, and it has not been found to produce untoward after-effects. The anæsthesia of eucaine develops more slowly than that of cocaine, but lasts longer. In nose and throat surgery, where a large quantity of anæsthetic has occasionally to be used, it is found to act efficiently, and to be without the poisonous effect previously produced in the same patients by similar doses of cocaine.

Owing to the sparing solubility of eucaine hydrochloride in cold water, and to the fact that it freely crystallizes from hot water on cooling, it is necessary that solutions for anæsthetic purposes should be made only at the moment they are required. For this purpose "Soloid" Eucaine Hydrochloride is most convenient. By dissolving one or more in definite quantities of warm water, solutions of any desired strength can be instantly prepared, and the operator is assured of obtaining the full anæsthetic effect of the drug. In laryngological practice eucaine may be used, either as a spray or as an injection, in the same strength as cocaine.

Supplied in bottles of 25, 1 gr. and 5 gr., at 1s. 8d. and 7s. per bottle.

BURROUGHS, WELLCOME &. Co.,

SNOW HILL BUILDINGS, LONDON.

Australasian Branch: 108, PITT STREET, SYDNEY, N.S.W.

Cable and Telegraphic Address: "BURCOME, LONDON."