




On ordinary differentially large fields

Omar León Sánchez and Marcus Tressl 

Abstract. We provide a characterization of differentially large fields in arbitrary characteristic and a single derivation in the spirit of Blum axioms for differentially closed fields. In the case of characteristic zero, we use these axioms to characterize differential largeness in terms of being existentially closed in the differential algebraic Laurent series ring, and we prove that any large field of infinite transcendence degree can be expanded to a differentially large field even under certain prescribed constant fields. As an application, we show that the theory of proper dense pairs of models of a complete and model-complete theory of large fields, is a complete theory. As a further consequence of the expansion result we show that there is no real closed and differential field that has a prime model extension in closed ordered differential fields, unless it is itself a closed ordered differential field.

1 Introduction

The class of differentially large fields was introduced and studied by the authors in [LST24]. Evidenced by the results in that paper, this class can be considered as the differential analog of the class of large fields. We recall that a field K is said to be large (aka ample) if every irreducible algebraic variety over K with a smooth K -point has a Zariski dense set of K -points. Equivalently, K is e.c. (existentially closed) in the field of formal Laurent series $K((t))$.

In [LST24], a differential field (K, Δ) of characteristic zero with commuting derivations $\Delta = \{\delta_1, \dots, \delta_m\}$ is defined to be differentially large if K is large (as a field) and for every differential extension (L, Δ) , if K is e.c. in L (as a field), then (K, Δ) is e.c. in (L, Δ) as a differential field (see [LST24, 2.1] for the algebraic meaning of e.c.). Several foundational properties and applications are explored in [LST24]. In particular, it is shown that (K, Δ) is differentially large if and only if (K, Δ) is e.c. in $(K((t_1, \dots, t_m)), \Delta)$, where the derivations on $K((t_1, \dots, t_m))$ are the natural ones extending those on K that commute with meaningful sums and satisfy $\delta_j(t_i) = \frac{dt_i}{dt_j}$.

The first-order characterization of differential largeness provided in [LST24, 4.7] makes reference to the somewhat elaborate axiom scheme UC from [Tre05, Corollary 4.5]. In Theorem 2.8 below, we give a significant simplification of this axiom scheme in the ordinary case, i.e., the case of a single derivation, so $\Delta = \{\delta\}$. The new scheme resembles the Blum axioms for differentially closed fields of characteristic 0 (DCF₀) and at the same time allows an extension of the notion of differential largeness to

Received by the editors August 1, 2024; revised December 16, 2024; accepted December 16, 2024.

Published online on Cambridge Core December 23, 2024.

The first author was partially supported by EPSRC grant EP/V03619X/1.

AMS subject classification: 12H05, 12E99, 03C60.

Keywords: Differential fields, large fields, formal Laurent series.



arbitrary characteristic (cf. Definition 2.1). In subsequent sections, we give applications of our new simple description of differential largeness in the ordinary case as follows.

Henceforth we restrict to a single derivation. An immediate consequence of the new axioms is the new characterization Corollary 2.9 of closed ordered differential fields (CODF), in the sense of Singer [Sin78], which does not make reference to the order. A further Corollary 2.12 provides geometric axioms for differentially large fields in arbitrary characteristic in terms of D-varieties, in the spirit of the Pierce–Pillay axioms for DCF_0 , see [PP98].

In the rest of the paper, we readopt the characteristic zero assumption. In Section 3, we prove that differential largeness can be characterized in terms of being existentially closed in the differential algebraic formal Laurent series, see Corollary 3.5. Our proof uses an approximation-type statement that resembles that of Denef–Lipshitz in [DL84]. We then use this to produce a new way (or rather an improvement of the construction in [LST24] for the ordinary case) to construct differentially large fields using iterated differential algebraic Laurent series, see Theorem 3.8.

In Section 4, we show that for any ordinary differential field (K, d) and any given large field $L \supseteq K$ of transcendence degree over K at least the size of K , there is an extension δ of d to L such that (L, δ) is differentially large, see Theorem 4.3. This has two consequences: Firstly, large fields of infinite transcendence degree (over \mathbb{Q}) are characterized in Corollary 4.5 as exactly those fields that possess a derivation d for which (L, d) is differentially large (significantly generalizing an earlier result by Christian Michaux saying that \mathbb{R} carries a CODF structure). Secondly, we show in Theorem 6.2 that no real closed field equipped with any derivation has a prime model extension in CODF, unless it is already a CODF; this strengthens a result from [Sin78] stating that the theory CODF does not have a prime model.

Theorem 4.3 is significantly strengthened in Section 5 in the case when the constant field C of K is dense for the étale open topology of L (see 5.5 for its definition). Namely we show in Theorem 5.8 that L can be expanded to a differentially large field whose constant field is algebraic over C . This Theorem has an interesting consequence for dense pairs of large fields: In Corollary 5.12, we show that for any complete and model complete theory T of large fields of characteristic 0 in the language \mathcal{L} of rings, possibly extended by constants, the theory of proper dense pairs of models of T is complete and inherits various neostability theoretic properties from T .

By a *differential ring* in this paper, we always mean a commutative unital ring furnished with a single derivation.

2 Blum-style axioms for ordinary differentially large fields

In [LST24], differentially large fields in characteristic zero were introduced. The definition there makes sense also for ordinary differential fields of characteristic $p > 0$.

Definition 2.1 A differential field (K, d) , of arbitrary characteristic, is said to be differentially large if it is large as a field and for every differential field extension $(L, \delta)/(K, d)$, if K is e.c. in L as a field, then (K, d) is e.c. in (L, δ) .

Examples of differentially large fields in characteristic $p > 0$ are differentially closed fields in the sense of Wood [Woo73], and also separably differentially closed fields

in the sense of Ino and the first author [IL23]. Recall that a differential field (K, δ) is said to be separably differentially closed if for every differential field extension $(L, \delta)/(K, \delta)$ with L/K separable (as fields), (K, δ) is e.c. in (L, δ) . To see that this class of differential fields is differentially large one only needs to note that if K is e.c. in L as a field, then L/K is separable.

Let (K, δ) be a differential field (of arbitrary characteristic). In what follows, we freely and interchangeably view any differential polynomial $f \in K\{x\}$ of order n as a differential polynomial in the differential variables $x = (x_1, \dots, x_m)$ and also as a polynomial in $m(n+1)$ algebraic variables $x, \delta x, \dots, \delta^n x$. It will be clear from the context which view we are taking; for instance, if $a \in K^{m(n+1)}$ and we write $f(a) = 0$, we mean viewing f as a polynomial in $m(n+1)$ variables.

In Theorem 2.8 below, we provide Blum-style axioms for ordinary differentially large fields of arbitrary characteristic. The proof relies on the following fact and its consequences, about extending derivations.

Fact 2.2 [Jac64, Theorem 18, Section IV.7] *Suppose L/K is a separable field extension. If $\delta : K \rightarrow L$ is a derivation, then δ can be extended to a derivation $L \rightarrow L$.*

Corollary 2.3 *Let $(K, \delta) \subseteq (L, \delta)$ be an extension of differential fields and let E be a subset of L with $L/K(E)$ separable. Then there is a derivation $\partial : K(E \cup \delta(E)) \rightarrow K(E \cup \delta(E))$ that restricts to δ on $K(E)$.*

If E is finite, then for each such ∂ there is some $f \in K[E \cup \delta(E)]$ such that ∂ restricts to a derivation of the localisation $K[E \cup \delta(E)]_f \rightarrow K[E \cup \delta(E)]_f$.

Proof Since $\delta(K(E)) \subseteq K(E \cup \delta(E))$ we may apply Fact 2.2 to the derivation $\delta|_{K(E)} : K(E) \rightarrow K(E \cup \delta(E))$ and get a derivation $\partial : K(E \cup \delta(E)) \rightarrow K(E \cup \delta(E))$ that restricts to δ on $K(E)$. Assume then that E is finite. There is some nonzero $f \in K[E \cup \delta(E)]$ such that $f \cdot \partial(\delta(a)) \in K[E \cup \delta(E)]$ for each $a \in E$. Obviously f has the required property. ■

Proposition 2.4 *Let K be a differential field and let $S = (S, \delta)$ be a differentially finitely generated K -algebra and a domain such that S/K is separable, i.e., $\text{Quot}(S)/K$ is a separable field extension. Let A be a finitely generated K -subalgebra of S . Then there are an element $f \in S$, a finitely generated K -subalgebra B of S_f containing A , a derivation ∂ on B and a differential K -algebra homomorphism $S \rightarrow (B, \partial)$ that restricts to the identity map on A . In particular $\partial a = \delta a$ for all $a \in A$.*

Proof Let $b \in S^n$ be such that S is the differential K -algebra generated by b and $A \subseteq K[b]$. Let $\mathfrak{p} = \{f \in K\{x\} \mid f(b) = 0\}$ be the differential vanishing ideal of b over K . Then \mathfrak{p} is a separable prime differential ideal; separability is due to fact that $K\{x\}/\mathfrak{p}$ is K -isomorphic to S . By the differential basis theorem of Kolchin [Kol73, Corollary 4, Section III.5], there is a finite set $\Sigma \subseteq \mathfrak{p}$ that generates \mathfrak{p} as a radical differential ideal. Take $d \geq 1$ such that each derivative of any x_1, \dots, x_n occurring in some polynomial from Σ has order $\leq d$. Finally take

$$E = \{\delta^k b_i \mid i \in \{1, \dots, n\}, k \leq d\} \subseteq S.$$

By possibly taking a larger d , a result of Kolchin appearing in [Kol73, Lemma 1, Section III.2] tells us that $S/K(E)$ is separable. By Corollary 2.3, there are $f \in K[E \cup \delta(E)]$

and a derivation ∂ of $B := K[E \cup \delta(E)]_f$ that restricts to δ on $K[E]$. Then $\partial^k b_i = \delta^k b_i$ for all $i \in \{1, \dots, n\}$, $k \leq d$ and therefore b is a solution to $\Sigma = 0$ in (B, ∂) . Consequently, the identity map of $K \cup \{b_1, \dots, b_n\}$ extends to a differential K -algebra homomorphism $\varphi : S \rightarrow (B, \partial)$. By choice of b , the map φ restricts to the identity map of A . ■

Corollary 2.5 *Let Σ be a set of differential polynomials over (K, δ) in finitely many differential variables. Suppose $\Sigma = 0$ has a solution in some differential field extension (L, δ) with L/K separable. Then there is a finitely generated K -subalgebra B of L and a derivation ∂ of B such that (B, ∂) has a solution to $\Sigma = 0$. In particular, (B, ∂) is differentially algebraic over (K, δ) and B/K is separable.*

Notice that if K is e.c. in L as a field then K is also e.c. in B as a field.

Proof By assumption, there is a solution of $\Sigma = 0$ in a differentially finitely generated K -subalgebra S of L . Now apply Proposition 2.4 to S and $A = K$. ■

Remark 2.6 In the case of several commuting derivations statements similar to Proposition 2.4 and Corollary 2.5 fail in general. This follows from examples produced by Johnson, Reinhart, and Rubel [JRR95, Theorem 2]. In particular, working over $(\mathbb{C}(z_1, z_2), \delta_1 \equiv \frac{\partial}{\partial z_1}, \delta_2 \equiv \frac{\partial}{\partial z_2})$, they prove that the PDE

$$\delta_2(x) = \left(1 - \frac{z_1}{z_2}\right)x + 1$$

has no differential algebraic solutions (equivalently, has no solution in a differential field extension of finite transcendence degree over \mathbb{C}).

Given a differential field K and a differential polynomial $f \in K\{x\}$, where x is a single differential variable, we denote by s_f the separant of f ; namely, the formal partial derivative of f with respect to its highest order variable. We write $[f]$ for the differential ideal generated by f in $K\{x\}$ and

$$[f] : s_f^\infty = \{g \in K\{x\} : s_f^m g \in [f] \text{ for some } m \geq 0\}.$$

Observation 2.7 *Let K be a differential field and let $f \in K\{x\}$ for x a single differential variable. Let $n = \text{ord}(f) \geq 0$ and let $a \in K^{n+1}$ with $f(a) = 0$ and $s_f(a) \neq 0$. Then there is an irreducible factor h of f with $\text{ord}(h) = n$, $h(a) = 0$ and $s_h(a) \neq 0$.*

Proof Let $f_0, f_1 \in K\{x\}$, with f_0 irreducible, $f = f_0 \cdot f_1$ and $\text{ord}(f_0) = n$. Then

$$(*) \quad s_f = \frac{\partial f}{\partial x_n} = \frac{\partial f_0}{\partial x_n} \cdot f_1 + f_0 \cdot \frac{\partial f_1}{\partial x_n}.$$

If $f_0(a) = 0$, then $(*)$ implies $s_{f_0}(a) = \frac{\partial f_0}{\partial x_n}(a) \neq 0$. If $f_0(a) \neq 0$, then $f_1(a) = 0$ and $(*)$ shows $s_{f_1}(a) = \frac{\partial f_1}{\partial x_n}(a) \neq 0$; hence also $\text{ord}(f_1) = n$ and in this case we may replace f by f_1 and proceed by induction. ■

We now come to the promised axiomatization.

Theorem 2.8 *Let (K, δ) be an ordinary differential field of arbitrary characteristic. The following conditions are equivalent.*

- (i) (K, δ) is differentially large.
 (ii) K is large as a field and for every pair $f, g \in K\{x\}$, where x is a single differential variable, with g nonzero and $\text{ord}(f) > \text{ord}(g)$, if the system

$$f(x) = 0 \text{ \& } s_f(x) \neq 0$$

has an algebraic solution in K , then $f(x) = 0 \text{ \& } g(x) \neq 0$ has a differential solution in K .¹

- (iii) For every pair $f, g \in K\{x\}$, where x is a single differential variable, with $\text{ord}(f) \geq 1$ and $\text{ord}(f) \geq \text{ord}(g)$, if the system

$$f(x) = 0 \text{ \& } g(x) \cdot s_f(x) \neq 0$$

has an algebraic solution in K , then it has infinitely many differential solutions in K . Notice that each of the properties (ii) and (iii) gives an axiom scheme for a first order axiomatization of differential largeness in the language of differential rings.

Proof (i) \Rightarrow (iii). Let $f, g \in K\{x\}$ with $\text{ord}(f) \geq 1$ and $\text{ord}(f) \geq \text{ord}(g)$ and assume

$$(\dagger) \quad f(x) = 0 \text{ \& } g(x) \cdot s_f(x) \neq 0$$

has an algebraic solution in K . Let $n = \text{ord}(f)$. By Observation 2.7, we may assume that f is irreducible. Let $\mathfrak{p} = [f] : s_f^\infty$. Since $s_f \neq 0$, Theorem 3.1(2) of [IL23] says that \mathfrak{p} is a separable prime differential ideal of $K\{x\}$. We write $a = x \bmod \mathfrak{p}$. Now, an algebraic solution of $f(x) = 0 \text{ \& } s_f(x) \neq 0$ in K is a smooth K -rational point of

$$K[x_0, \dots, x_n]/(f) \cong_K K[a, \dots, a^{(n)}].$$

The largeness of K yields that K is e.c. in $K(a, \dots, a^{(n)})$. Since the latter is equal to the differential field $K\langle a \rangle$ generated by a over K , differential largeness implies that (K, δ) is e.c. in $(K\langle a \rangle, \delta)$.

Since $\text{ord}(f) \geq \text{ord}(g)$ and (\dagger) has an algebraic solution in K , Lemma 3.6(1) of [IL23] implies that $g \cdot s_f \notin \mathfrak{p}$. Hence a is a differential solution of (\dagger) in $K\langle a \rangle$. As (K, δ) is e.c. in $(K\langle a \rangle, \delta)$ also K has a differential solution α of (\dagger) . To argue that there are infinitely many solutions, note that $g \cdot (x - \alpha)$ has again order at most $\text{ord}(f)$. By largeness of K and the assumption $\text{ord}(f) \geq 1$, there is an algebraic solution of the new system where we replace g with $g \cdot (x - \alpha)$. It follows, by repeating the above argument, that there are infinitely many differential solutions of (\dagger) in K .

(iii) \Rightarrow (ii) It suffices to show that K is large as a field. By [Jar11, Lemma 5.3.1, p. 67], a field K is large if and only if for every absolutely irreducible polynomial $F(X, Y) \in K[X, Y]$, if there is a point $(a, b) \in K^2$ with $F(a, b) = 0$ and $\frac{\partial F}{\partial Y}(a, b) \neq 0$, then there are infinitely many such points.

So take an absolutely irreducible polynomial $F(X, Y) \in K[X, Y]$ and some $(a, b) \in K^2$ with $F(a, b) = 0$ and $\frac{\partial F}{\partial Y}(a, b) \neq 0$. Consider the differential polynomial $f(x) = F(x, x')$. Then $f(x) = 0 \text{ \& } s_f(x) \neq 0$ has an algebraic solution in K , namely (a, b) . By (iii) there are infinitely many differential solutions in K . But then there are infinitely many solutions to $F(X, Y) = 0$ and $\frac{\partial F}{\partial Y}(X, Y) \neq 0$ in K as well.

(ii) \Rightarrow (i). To prove differential largeness, let F be a differential field extension of K such that K is e.c. in F as a field. Note that then F/K is separable. We need to show

¹By 2.7 we may also assume that f is irreducible in this condition.

that K is e.c. in F as a differential field. Let Σ be a system of differential polynomials in n differential variables over K and assume that $\Sigma = 0$ has a solution $a \in F^n$. We may assume that $F = K\langle a \rangle$. By 2.5 applied to F , we may assume that F is differentially algebraic over K (and F/K remains separable).

Condition (ii) guarantees that $[K : C_K]$ is infinite; hence, by the differential primitive element theorem [Kol73, Proposition 9, Section II.8, p. 103], the differential field F is differentially generated over K by a single element $b \in F$. Let \mathfrak{p} be the prime differential ideal of $K\{x\}$ associated with b . Note that \mathfrak{p} is separable (over K).

Then, by Theorem 3.1(1) of [IL23], $\mathfrak{p} = [f] : s_f^\infty$ for $f \in \mathfrak{p}$ irreducible of minimal rank. Write $a = (a_1, \dots, a_n)$ and let $f_i, g \in K\{x\}$ with $a_i = \frac{f_i(b)}{g(b)}$. By the differential division algorithm [Kol73, Section I.9] there are $h \in K\{x\}$ reduced with respect to f and some $r \geq 0$ with

$$(i_f s_f)^r g \equiv h \pmod{[f]}.$$

Since $f(b) = 0$ and $i_f(b) \cdot s_f(b) \neq 0$ we get $i_f^r(b) s_f^r(b) g(b) = h(b) \neq 0$. Hence, we may replace g by h and f_i by $(i_f s_f)^r \cdot f_i$ if necessary and assume that g is reduced with respect to f . Notice that $a_i \in K\{b\}_{g(b)}$.

Now, since K e.c. in F as a field, the system $f(x) = 0$ & $s_f(x) \neq 0$ has an algebraic solutions in K . By condition (ii), the set

$$\{f = 0\} \cup \{q \neq 0 \mid q \in K\{x\} \text{ is nonzero and } \text{ord}(q) < \text{ord}(f)\}$$

is finitely satisfiable in the differential field K . Hence there is an elementary extension L of the differential field K having a differential solution c to $f(x) = 0$ such that $q(c) \neq 0$ for all $q \in K\{x\}$ with $\text{ord}(q) < \text{ord}(f)$. Since f is irreducible, it follows that $q(c) \neq 0$ for all $q \in K\{x\}$ that are reduced with respect to f .

In particular $f(c) = 0$ & $g(c) \neq 0$. Since $K < L$ there is some $d \in K$ with $f(d) = 0$ & $g(d) \neq 0$. This means there is a differential K -homomorphism $(K\{x\}/\mathfrak{p})_{g \bmod \mathfrak{p}} \rightarrow K$. By choice of \mathfrak{p} we have $(K\{x\}/\mathfrak{p})_{g \bmod \mathfrak{p}} \cong K\{b\}_{g(b)}$ as differential K -algebras. Since $K\{a_1, \dots, a_n\} \subseteq K\{b\}_{g(b)}$, we obtain a differential K -algebra homomorphism $K\{a_1, \dots, a_n\} \rightarrow K$ and this corresponds to a differential solution of $\Sigma = 0$ in K^n . ■

When K is real closed, the above theorem yields a new axiomatization of the theory CODF. A differential field (K, δ) is a model of CODF if and only if it is an existentially closed model of the theory of ordered differential fields. Axioms for CODF appear in [Sin78]. While the axioms there make explicit reference to the order, our new axioms are purely in the differential field language, namely:

Corollary 2.9 *Let (K, δ) be a differential field. The following are equivalent.*

- (i) $(K, \delta) \models \text{CODF}$
- (ii) K is real closed and for every pair $f, g \in K\{x\}$, where x is a single differential variable, with g nonzero and $\text{ord}(f) > \text{ord}(g)$, if the system

$$f(x) = 0 \text{ \& \> } s_f(x) \neq 0$$

has an algebraic solution in K , then $f(x) = 0$ & $g(x) \neq 0$ has a differential solution in K .

Notice that every field K is algebraically closed in the large field $K((t))$, but not every field is large. In the differential phrasing this changes:

Corollary 2.10 *Let L/K be an extension of differential fields. If K is differentially algebraically closed in L and L is differentially large, then K is differentially large as well.*

Proof We verify 2.8(iii). Take $f, g \in K\{x\}$, x a single differential variable, with $\text{ord}(f) \geq 1$ and $\text{ord}(f) \geq \text{ord}(g)$, and assume that $f(x) = 0$ & $g(x) \cdot s_f(x) \neq 0$ has an algebraic solution in K . Since L is differentially large, it has infinitely many differential solutions to $f(x) = 0$ & $g(x) \cdot s_f(x) \neq 0$. But then each of these solutions is differentially algebraic over K . Hence all these solutions are in K . ■

Remark 2.11 We note that the condition of a differential field (K, δ) being differentially algebraically closed in some extension (L, δ) is quite strong. Arguably, being differentially algebraically closed in an extension is not quite the right differential analog of being algebraically closed in the field sense. We do not know whether the assumption in Corollary 2.10 can be weakened to only assuming that K is constrainedly closed in L (namely, every finite tuple from L which is constrained over K , in the sense of Kolchin [Kol73, Section III.10], is from K).

We conclude this section with a geometric characterization of being differentially large. Namely, in terms of algebraic D-varieties. Recall that an algebraic D-variety over K is a pair (V, s) where V is an algebraic variety over K and $s : V \rightarrow \tau V$ is a section over K of the prolongation of V (see [KP05, Section 2], for instance). The latter is the algebraic bundle $\pi : \tau V \rightarrow V$ with the characteristic property that for any differential field extension (L, δ) of (K, δ) we have that if $a \in V(L)$ then $(a, \delta a) \in \tau V$.

Corollary 2.12 *Let K be a large field of arbitrary characteristic and let δ be a derivation of K . The following conditions are equivalent.*

- (i) (K, δ) is differentially large
- (ii) *Let V and W be K -irreducible algebraic varieties with $W \subseteq \tau V$. If $\pi|_W : W \rightarrow V$ is a separable morphism and W has a smooth K -point, then the set*

$$\{(a, \delta a) \in W : a \in V(K)\}$$

is Zariski dense in W .

- (iii) *Let (V, s) be a K -irreducible algebraic D-variety. If V has a smooth K -point, then the set*

$$\{a \in V(K) : s(a) = (a, \delta(a))\}$$

is Zariski dense in V .

Proof (i) \Rightarrow (ii) Let (a, b) be a K -generic point of W . Since $\pi_W : W \rightarrow V$ is a separable morphism, we obtain that a is K -generic in V and $K(a, b)/K(a)$ is a separable extension. Since $W \subseteq \tau V$, there is a derivation $\delta : K(a) \rightarrow K(a, b)$ extending the one on K such that $\delta(a) = b$. As $K(a, b)/K(a)$ is separable, by 2.2, we can extend the derivation to $K(a, b) \rightarrow K(a, b)$. Then, for any nonempty Zariski-open $O_W \subseteq W$ over K , in the differential field extension $(K(a, b), \delta)$ we can find a solution to $x \in V$ and $(x, \delta x) \in O_W$ (namely, the tuple a). Since W has a smooth K -point, we get that K is e.c.

in $K(W) = K(a, b)$ as a field. By differential largeness, (K, δ) is e.c. in $(K(a, b), \delta)$, and so we can find the desired solution in K .

(ii) \Rightarrow (iii) If we let $W = s(V) \subseteq \tau V$, then the pair V and W satisfy the conditions of (ii) (note that if b is a smooth point of V then $(b, s(b))$ is a smooth point of W). It follows that the set of points in W of the form $(a, \delta a)$ with $a \in V(K)$ is Zariski dense in W . But then, as $W = s(V)$, the set of points $a \in V$ such that $s(a) = (a, \delta a)$ must be Zariski dense in V .

(iii) \Rightarrow (i) We verify 2.8(ii). Let $f, g \in K\{x\}$ with $\text{ord}(g) < \text{ord}(f)$ and g nonzero. Assume the system

$$f(x) = 0 \text{ \& } s_f(x) \neq 0$$

has an algebraic solution in K . In particular, $s_f \neq 0$. By Observation 2.7, we may assume that f is irreducible. By Theorem 3.1(1) of [IL23], $\mathfrak{p} = [f] : s_f^\infty$ is a separable prime differential ideal of $K\{x\}$. Let $a = x + \mathfrak{p}$ in the fraction field of $K\{x\}/\mathfrak{p}$. Letting $n = \text{ord}(f)$, we see that $(a, \delta a, \dots, \delta^{n-1}a)$ is algebraically independent over K and $\delta^n a$ is separably algebraic over $K(a, \dots, \delta^{n-1}a)$. It follows that

$$\delta^{n+1}a = \frac{h(a, \delta a, \dots, \delta^n a)}{s_f(a)}$$

for some $h \in K[t_0, \dots, t_n]$. Let V be the localisation at $g \cdot s_f$ of the Zariski-locus of $(a, \delta a, \dots, \delta^n a)$ over K . From the assumptions (on existence of an algebraic solution in K), we see that V has a smooth K -rational point and that the morphism on V induced by

$$(t_0, t_1, \dots, t_n) \mapsto ((t_0, t_1, \dots, t_n), \left(t_1, t_2, \dots, t_n, \frac{h(t_0, t_1, \dots, t_n)}{s_f(t_0, t_1, \dots, t_n)}\right))$$

yields a regular algebraic map $s : V \rightarrow \tau V$. This equips V with a D-variety structure. Then, the assumption of (iii) yields $\alpha \in V(K)$ such that $s(\alpha) = (\alpha, \delta \alpha)$. But then α is the desired differential solution of $f(x) = 0$ & $g(x) \neq 0$ in K . ■

3 Power series in characteristic zero

In this section, we assume fields are of characteristic zero, and thus the results on differentially large fields from [LST24] may be deployed. We prove, in Corollary 3.5, two further characterizations of being differentially large.

For a differential field K (ordinary throughout), we endow $K((t))$ with its natural derivation extending the given derivation on K and satisfying $\delta(t) = 1$; that is,

$$\delta\left(\sum_{n \geq k} a_n t^n\right) = \sum_{n \geq k} \delta(a_n) t^n + \sum_{n \geq k} n a_n t^{n-1}.$$

In [LST24, 4.3] it is shown that (K, δ) is differentially large if and only if (K, δ) is e.c. in $(K((t)), \delta)$. We do not know if this characterization extends to positive characteristic, the proof relies on the existence of a *twisted version* of the Taylor morphism [LST24, 3.4], whose construction picks up rational denominators. Below

we prove that it suffices to ask for (K, δ) to be e.c. in the differential subfield of $(K((t)), \delta)$ consisting of differential algebraic elements (over K).

Definition 3.1 Let K be a differential field and let S be a differential K -algebra. We write S_{diffalg} for the differential subring of all $a \in S$ that are differentially algebraic over K .

Remark 3.2 Since $K((t))$ is the localization of $K[[t]]$ at t , the fraction field of $K[[t]]_{\text{diffalg}}$ is $K((t))_{\text{diffalg}}$.

Proposition 3.3 Let (K, δ) be a differential field (of characteristic zero) that is large as a field and let S be a differentially finitely generated K -algebra. If there is a K -algebra homomorphism $S \rightarrow L$ for some field extension L/K in which K is e.c. (as a field, there are no derivations on L given), then there is a differential K -algebra homomorphism $S \rightarrow K[[t]]_{\text{diffalg}}$.

Proof By [LST24, 3.5] there is a differential K -algebra homomorphism $\psi : S \rightarrow K[[t]]$. Applying 2.4 to $\psi(S)$, we may then find a finitely generated K -subalgebra B of $K((t))$, a derivation ∂ of B extending δ on K together with a differential K -algebra homomorphism $\varphi : \psi(S) \rightarrow (B, \partial)$. By [LST24, 3.5] applied to (B, ∂) and the inclusion map $B \hookrightarrow K((t))$ there is a differential K -algebra homomorphism $\gamma : B \rightarrow K[[t]]$. Since B is a finitely generated K -algebra, the image of γ is in $K[[t]]_{\text{diffalg}}$. Hence the map $\gamma \circ \varphi \circ \psi : S \rightarrow K[[t]]_{\text{diffalg}}$ has the required property. ■

A special case of 3.3 resembles an approximation statement over large and differential fields in the spirit of [DL84, Theorem 2.1].

Corollary 3.4 Let (K, δ) be a differential field of characteristic zero such that K is large as a field. Let Σ be a system of differential polynomials in finitely many differential variables over K . If the differential ideal generated by Σ has an algebraic solution in $K((t))$, then $\Sigma = 0$ has a differential solution in $K[[t]]_{\text{diffalg}}$.

Proof Apply 3.3 to the differential coordinate ring of Σ . ■

Corollary 3.5 Let K be a large field of characteristic 0 and let δ be a derivation of K . The following conditions are equivalent.

- (i) (K, δ) is differentially large.
- (ii) K is e.c. in $K[[t]]_{\text{diffalg}}$ as a differential field.
- (iii) For every K -irreducible algebraic D -variety (V, s) , if V has a K -point, then there is $a \in V(K)$ such that $s(a) = (a, \delta a)$.

Proof (i) \Rightarrow (ii) is a consequence of [LST24, 4.3(ii)], which says that K is e.c. in $K((t))$ as a differential field.

(ii) \Rightarrow (i). By 3.4 one verifies that K is e.c. in $K((t))$ as a differential field. Hence by [LST24, 4.3], (K, δ) is differentially large.

(iii) \Rightarrow (i) We verify 2.12(iii). Let (V, s) be a K -irreducible D -variety with a smooth K -point. Let $h \in K[V]$ nonzero. Then, there is an induced D -variety structure in the localization $K[V]_h$. Denote this D -variety by (W, t) . As K is large and V has a smooth K -point, we get that K is Zariski dense in V . Thus, W has a K -point. The assumption now yields a K -point b in W such that $s(b) = (b, \delta b)$. As h was arbitrary, it follows that the set of points $\{a \in V(K) : s(a) = (a, \delta a)\}$ is Zariski dense.

(i) \Rightarrow (iii) Let (V, s) be a K -irreducible D -variety with a K -point. Applying Proposition 3.3 with $S = K[V]$ and $L = K$, we find a $K((t))$ -rational point b of V such that $s(b) = (b, \delta b)$. As K is differentially large, it is e.c. in $K((t))$ as a differential field. Hence, we can find such a point in K . ■

We may now improve the construction of differentially large fields from [LST24, 5.2] in the ordinary case. A few preparations are necessary.

Proposition 3.6 *Let $(K_i, f_{ij})_{i,j \in I}$ be a directed system of differential fields and differential embeddings with the following properties.*

- (a) *All K_i are large as fields.*
- (b) *All embeddings $f_{ij} : K_i \rightarrow K_j$ are isomorphisms onto a subfield of K_j that is e.c. in K_j as a field.*
- (c) *For all $i \in I$ there exist $j \geq i$ and a differential homomorphism $K_i[[t]]_{\text{diffalg}} \rightarrow K_j$ extending f_{ij} .*

Then the direct limit L of the directed system is a differentially large field.

Proof The proof is identical to the proof of [LST24, 5.1], except we use Proposition 3.3 in that proof instead of [LST24, 3.5]. ■

Observation 3.7 *Let K be a differential field. Then $K[[t]]_{\text{diffalg}}$ is a Henselian valuation ring.*

Proof We write $S = K[[t]]_{\text{diffalg}}$. Since $S = K[[t]] \cap K((t))_{\text{diffalg}}$, it is a valuation ring. Clearly the maximal ideal of S is $t \cdot S$. To verify that S is Henselian it suffices to show that for all $\mu_2, \dots, \mu_n \in \mathfrak{m}$ there is some $f \in S$ with

$$1 + f + \mu_2 f^2 + \dots + \mu_n f^n = 0.$$

As $K[[t]]$ is Henselian, there is such an f in $K[[t]]$. Obviously, $f \in S$. ■

Theorem 3.8 *Let (K, δ) be any differential field of characteristic zero. Set $K_0 = K$ and let $K_{n+1} = K_n((t_n))_{\text{diffalg}}$. Then $\bigcup_{n \geq 0} K_n$ is differentially large.*

Proof By Observation 3.7, $K_n[[t_n]]_{\text{diffalg}}$ is a Henselian valuation ring. By [Pop10], $K_n((t_n))_{\text{diffalg}}$ is a large field. We see that all assumptions of Proposition 3.6 are satisfied for the K_n and the inclusion maps $K_n \hookrightarrow K_{n+k}$. Now the argument for [LST24, 5.2(i)] can be copied, where we use 3.6 instead of [LST24, 5.1]. ■

Remark 3.9 For every differential field K , there are power series in $K((t))$ that are differentially transcendental over K (here $K((t))$ is furnished with the natural derivation extending the given one on K and satisfying $\delta(t) = 1$); for example, the power series $f(t) = \sum_{n \geq 0} t^{2^n}$. Indeed, this power series, which lives in $\mathbb{Q}((t))$, is differentially transcendental over \mathbb{Q} by Mahler [Mah30] (also see [LR86]). Any such power series, viewed in $K((t))$, remains differentially transcendental over K because $\mathbb{Q}((t))$ and K are linearly disjoint over \mathbb{Q} in $K((t))$. This shows that, for any differential field K , the subfield $K((t))_{\text{diffalg}}$ of differentially algebraic elements is properly contained in $K((t))$.

We conclude this section by discussing possible improvements of Proposition 3.3.

Counter example 3.10 If K is algebraically closed in Proposition 3.3 then a stronger conclusion holds, namely there is a differential K -algebra homomorphism $S \rightarrow K[[t]]$ whose image is constrained. The reason is that there is a differential homomorphism $\varepsilon : S \rightarrow K^{\text{diff}}$ and then one can apply Proposition 3.3 to obtain a differential embedding of the image of ε into $K[[t]]$.

However, if K is not algebraically closed then in general there is no differential K -algebra homomorphism $S \rightarrow K[[t]]$ whose image is constrained. To see an example, consider the ordered field $\mathbb{R}(z)$ where $z > \mathbb{R}$ and let K be its real closure. We furnish K with the unique derivation extending the standard derivation $\frac{d}{dz}$ on $\mathbb{R}(z)$. Let x be a new transcendental element and let R be the real closure of the ordered field $K(x)$ with the ordering $x > K$. Extend the derivation of K to R by setting $\delta(x) = 0$. Let y be a square root of $x - z$ in R and let S be the differential K -subalgebra of R generated by y , hence $S = K[y, y^{-1}]$. Now if $\varphi : S \rightarrow K[[t]]$ is a differential K -algebra homomorphism, then $\varphi(x)' = \varphi(x') = 0$ and $\varphi(x) = \varphi(y^2 + z) = \varphi(y)^2 + z$, hence $\varphi(x)$ is a constant and $\varphi(x) - z$ is a square. As $z > \mathbb{R}$ and \mathbb{R} is the constant field of K , we see that $\varphi(x)$ cannot be in K . Hence $\varphi(x)$ is a new constant of $K[[t]]$ and therefore it is not constrained over K .

3.11 (On the canonicity of differentially algebraic solutions) Let K be a differential field. If S is a differentially finitely generated K -algebra and $\varphi : S \rightarrow K$ is a K -algebra homomorphism, then by [LST24, 3.5] one can explicitly construct a differential K -algebra homomorphism $\psi : S \rightarrow K[[t]]$, namely one can take ψ to be the twisted Taylor morphism T_φ^* associated with φ . Now, by Proposition 3.3 there is even a differential K -algebra homomorphism $\rho : S \rightarrow K[[t]]^{\text{diffalg}}$ and one might ask whether ρ can also be obtained in some canonical form out of φ . However Gabriel Ng has shown that this is not possible. We refer to [Ng23, Proposition 7.11] for details.

4 Expansions of large fields to a differentially large field

The main goal of this section is Theorem 4.3 which implies that any large field of characteristic zero of infinite transcendence degree over \mathbb{Q} can be expanded to a differentially large field. A further consequence of Theorem 4.3 is 6.2, which says that prime model extensions in CODF only exist in the trivial case. Throughout this section fields are assumed to be of characteristic zero.

Notation 4.1 Let K be a field (of characteristic zero). A **differentially large problem** of K is a pair (f, g) of polynomials from $K\{x\} = K[x_0, x_1, \dots]$ such that f is of order $n \geq 0$, the order of g is strictly less than n and for which there is an element $(c_0, \dots, c_n) \in K^{n+1}$ such that

$$f(c_0, \dots, c_n) = 0 \text{ \& \& } s_f(c_0, \dots, c_n) \neq 0.^2$$

We call \bar{c} an **algebraic solution of the differentially large problem**. Obviously a differentially large problem over K remains a differentially large problem over every field extension of K . If d is a derivation of K , then a **solution of a differentially large**

²Note that there is no condition on g here.

problem of K in a differential field (L, δ) extending (K, d) is an element $a \in L$ with $f(a) = 0$ & $g(a) \neq 0$, where polynomials are now evaluated as differential polynomials.

Proposition 4.2 Let L/K be a field extension, $n \in \mathbb{N}$ and assume that $\text{tr. deg}(L/K) \geq n$. Let (f, g) be a differentially large problem of K with $\text{ord}(f) = n$. Let d be a derivation of K and assume L is large.

Then there is a subfield K_1 of L that is finitely generated over K as a field, a derivation δ of K_1 extending d and a solution $a \in K_1$ of the differentially large problem (f, g) such that $a, \delta a, \dots, \delta^{n-1}a$ are algebraically independent over K .

Proof Let $\bar{x} = (x_0, \dots, x_n)$ and let Z be the solution set in L of the system

$$f(\bar{x}) = 0 \text{ \& } s_f(\bar{x}) \neq 0.$$

Claim There exists a point $(a_0, \dots, a_n) \in Z$ with $\text{tr. deg}(a_0, \dots, a_n/K) = n$. ■

Proof Let W be the variety defined by the two polynomials

$$f(\bar{x}), y \cdot s_f(\bar{x}) - 1 \in K[\bar{x}, y].$$

Write $h(\bar{x}, y) = y \cdot s_f(\bar{x}) - 1$. Then any common zero (\bar{a}, c) of f and h in the algebraic closure of L is a regular point of W , because $c \cdot s_f(\bar{a}) - 1 = 0$ implies $\frac{\partial f}{\partial x_n}(\bar{a}) \neq 0$ and obviously $\frac{\partial h}{\partial y} = s_f$ does not vanish at \bar{a} . Hence the determinant of the matrix

$$\begin{pmatrix} \frac{\partial f}{\partial x_n} & \frac{\partial f}{\partial y} \\ \frac{\partial h}{\partial x_n} & \frac{\partial h}{\partial y} \end{pmatrix} \text{ is not zero at } (\bar{a}, c). \text{ This shows that } W \text{ is smooth.}$$

Since (f, g) is a differentially large problem of K we know that W has a K -rational point. By [Fehl1, Theorem 1], using $\text{tr. deg}(L/K) \geq n = \dim(W)$, there is a K -embedding $K(W) \rightarrow L$. A generic point of W in $K(W)$ is then mapped to a point $(a_0, \dots, a_n) \in Z$ with $\text{tr. deg}(a_0, \dots, a_n/K) = n$. ■

As $s_f(a_0, \dots, a_n) \neq 0$, a_n is algebraic over $K(a_0, \dots, a_{n-1})$. But now we see that $g(a_0, \dots, a_n) \neq 0$ as the order of g is strictly less than n , and $K_1 := K(a_0, \dots, a_n)$ is isomorphic to the quotient field of $K\{x\}/\mathfrak{p}$, where $\mathfrak{p} = [f] : s_f^\infty$. This induces a derivation δ on K_1 and this derivation has the required properties: $a = a_0$ solves the given differentially large problem.

Theorem 4.3 Let L/K be an extension of fields of characteristic 0 and suppose L is a large field. Let d be a derivation of K . If $\text{tr. deg}(L/K) \geq \text{card}(K)$, then there is a derivation δ of L extending d such that (L, δ) is differentially large.

[Under necessary assumptions on the constant field C of K we will show in Theorem 5.8 that we may in addition find such a δ whose constant field is algebraic over C .]

Proof Let $\kappa = \text{card}(K)$. By extending K and d we may assume that $\text{tr. deg}(L/K) = \kappa$. Let $\{t_i \mid i < \kappa\}$ be a transcendence basis of L over K and let $(f_i, g_i)_{i \in \kappa}$ be a list of all differentially large problems of L ; so here $f_i, g_i \in L\{x\}$ in the terminology of Notation 4.1.

For $i < \kappa$ we define a subfield K_i of L and a derivation d_i of K_i such that

- (a) K_i contains t_i , $\text{tr. deg}(K_i/K)$ is finite for finite i and $\text{tr. deg}(K_i/K) \leq \text{card}(i)$ for $i \geq \omega$,

- (b) (K_i, d_i) extends (K_j, d_j) for $j < i$, and
- (c) (K_i, d_i) solves the differentially large problem (f_i, g_i) .

Suppose $i < \kappa$ and (K_j, d_j) has already been defined with properties (a)–(c); this also covers the case $i = 0$. Let $\bar{b} \subseteq L$ be finite with $f_i, g_i \in K(\bar{b})\{x\}$ such that there is an algebraic solution of the differentially large problem (f_i, g_i) in $K(\bar{b})$. Let K_* be the field generated by $K(t_i, \bar{b}) \cup \bigcup_{j < i} K_j$ and extend the derivation $\bigcup_{j < i} d_j$ to a derivation d_* of K_* arbitrarily. Obviously then $\text{tr. deg}(K_*/K)$ is finite if i is finite and $\leq \text{card}(i)$ otherwise.

Consequently $\text{tr. deg}(L/K_*)$ is infinite and we may apply Proposition 4.2 to the extension $K_* \subseteq L$, the derivation d_* and the differentially large problem (f_i, d_i) . We obtain an extension (K_i, d_i) of (K_*, d_*) such that K_i is a subfield of L that is finitely generated over K_* . Clearly (K_i, d_i) satisfies (a)–(c).

Then $L = \bigcup_{i < \kappa} K_i$ and by 2.8 the differential field (L, ∂) with $\partial = \bigcup_{i < \kappa} d_i$ is differentially large. ■

Remark 4.4 In characteristic $p > 0$ the conclusion in Theorem 4.3 fails even under the assumption that L/K is separable. For example L might be perfect (as a field), and hence any derivation on L is trivial.

Corollary 4.5 A large field L of characteristic zero is of infinite transcendence degree if and only if there is a derivation d of L such that (L, d) is differentially large.

Proof If L has infinite transcendence degree, then by Theorem 4.3 applied with $K = \mathbb{Q}$ shows that there is a derivation d of L such that (L, d) is differentially large. For the converse assume, there is a derivation d of L such that (L, d) is differentially large. By [LST24, 5.12], the algebraic closure \bar{L} of L is differentially closed. We may then replace L by the differential closure of \mathbb{Q} . By the non-minimality of the differential closure of \mathbb{Q} ([Ros74]), there is an embedding $L \rightarrow \bar{L}$ that is not surjective. Hence L cannot have finite transcendence degree. ■

5 Differentially large fields with prescribed constant field

We now aim to provide a version of Theorem 4.3 for all differentially large fields of characteristic 0 without extending the constants. More precisely, we prove in Theorem 5.8 below that for a field extension L/K with $\text{tr. deg}(L/K) \geq \text{card}(K)$, if d be a derivation of K whose constant field C_K is dense in L for the étale open topology of L , then there is an extension δ of d on L such that (L, δ) is differentially large and C_L is algebraic over C_K . Hence, under a density assumption of the constants C_K , if C_K is algebraically closed in L , then the construction of the derivation in Theorem 4.3 can be performed without introducing new constants. We conclude this section with an application to dense pairs of large fields in 5.12.

We first (briefly) introduce the notion of *L-prime ideals* in the context of a fixed field extension L/K . For a differential ring S we write $\text{Sped}(S)$ for the subspace of $\text{Spec}(S)$ consisting of the differential prime ideals of S .

Definition 5.1 Let L/K be an extension of fields and let d be a derivation of K . Let S be a differential K -algebra. We call a prime ideal \mathfrak{p} of S a **differential L -prime ideal** if it is differential and S/\mathfrak{p} can be embedded into L as a K -algebra; observe that there is

no derivation given on L . We write $\text{Sped}_L(S)$ for the subspace of $\text{Sped}(S)$ consisting of differential L -prime ideals. We say that a point $\mathfrak{p} \in \text{Sped}(S)$ is **L -locally closed** if it is a locally closed point of $\text{Sped}_L(S)$. If S is finitely generated as a K -algebra and (0) is the unique point of $\text{Sped}_L(S)$, we say that S is **L -simple**. Note that L -simplicity implies that there is a K -algebra embedding $S \rightarrow L$.

Examples 5.2

- (i) If L is an algebraically closed field of infinite transcendence degree over K and S is a differentially finitely generated K -algebra, then $\text{Sped}_L(S) = \text{Sped}(S)$, and L -locally closed is the same as being **constrained** in the sense of Kolchin [Kol74].
- (ii) If K is real closed and L is an $|S|^+$ -saturated real closed field, then $\text{Sped}_L(S)$ is the subspace of differential prime ideals \mathfrak{p} that are real, i.e., -1 is not a sum of squares in S/\mathfrak{p} . When we are in this example we will say **real constrained** instead of L -locally closed.
- (iii) Clearly, being constrained and real implies real-constrained. However, the converse does not always hold. For instance, consider the real closure $K = \mathbb{Q}(t)^{\text{rcl}}$, where $\mathbb{Q}(t)$ is equipped with the unique ordering such that $t > \mathbb{Q}$ and with the unique derivation extending $\frac{d}{dt}$ on $\mathbb{Q}(t)$. Let α_2 be a transcendental over K . In the formally real field $K(\alpha_2)$ define a derivation δ that extends the one on K such that $\delta(\alpha_2) = \frac{-1}{2\alpha_2}$. Let $\alpha_1 = t + \alpha_2^2$. Then, $\delta(\alpha_1) = 0$ and α_1 is transcendental over K (as $\alpha_1 > \mathbb{Q}^{\text{rcl}}$). Now consider $\alpha = (\alpha_1, \alpha_2)$. Clearly α is not constrained over K (as α_1 is a constant which is not algebraic over $C_K = \mathbb{Q}^{\text{rcl}}$). But α is real-constrained over K . Indeed, for any differential specialization $\beta = (\beta_1, \beta_2)$ of α over K with $K\langle\beta\rangle$ formally real, we see that β_1 is transcendental over K (for the same reason that α_1 was); thus, the map $K(\alpha) \rightarrow K(\beta)$ fixing K and mapping $(\alpha_1, \alpha_2) \mapsto (\beta_1, \beta_2)$ is a differential isomorphism (not necessarily preserving the orderings).

Observation 5.3 Let L be a differentially large field and let $K \subseteq L$ be a differential subfield. Then for every L -simple K -algebra S there is a differential K -embedding $S \rightarrow L$.

Proof As S is L -simple, there is a K -algebra embedding $\varphi: S \rightarrow L$. Now $S \otimes_K L$ is a finitely generated differential L -algebra and φ extends to an L -algebra homomorphism $S \otimes_K L \rightarrow L$. By differential largeness, [LST24, 4.3(iv)] says that there is a differential L -algebra homomorphism $\psi: S \otimes_K L \rightarrow L$. Composing ψ with the natural map $S \rightarrow S \otimes_K L$ gives a differential K -algebra homomorphism $S \rightarrow L$. Since S is L -simple, this map is an embedding. ■

Proposition 5.4 Let L/K be a field extension and let d be a derivation of K . Let S be a differentially finitely generated K -algebra. If $\text{Sped}_L(S)$ is nonempty, then there are L -locally closed points \mathfrak{p} of S and for each such point there is some $q \in S$ such that $(S/\mathfrak{p})_q$ is L -simple.

Proof By Noetherianity of $\text{Sped}(S)$, there are points of $\text{Sped}_L(S)$ that are maximal for inclusion in $\text{Sped}_L(S)$. These points are even closed in $\text{Sped}_L(S)$.

Now take an L -locally closed points \mathfrak{p} of S . Hence there is some $q \in S$ such that the prime ideal \mathfrak{p} is maximal for inclusion in $D(q) \cap \text{Sped}_L(S)$. In other words the zero ideal is the unique element of $\text{Sped}_L(A)$, where $A = (S/\mathfrak{p})_q$; in particular there is a K -algebra embedding $A \rightarrow L$.

By 2.4 there are $a \in A$ and a (not necessary differential) finitely generated K -subalgebra B of A_a and a derivation ∂ of B together with a differential K -algebra homomorphism $f : A \rightarrow (B, \partial)$. In particular the kernel of f is in $\text{Sped}(A)$. Since B is a K -subalgebra of A_a , the kernel of f is even in $\text{Sped}_L(S)$ and so the kernel of f is 0. Thus f is an embedding and $\text{qf}(A)$ is a finitely generated field extension of K . Since A is a differentially finitely generated K -algebra we may then localize it at some element of A and see that A is finitely generated as a K -algebra. ■

In what follows we will talk about the *étale open topology* on K -rational points of K -varieties for a field K , cf. [JTWY24, p. 4034]. Explicitly we will only need a few basic properties of the étale open topology of K itself and we only record what we need later on.

5.5 (The étale open topology) *Let K be a field.*

- (i) *We call a subset U of K **standard étale open** if it is the image of the projection $K^2 \rightarrow K$ onto the first coordinate of a set of the form $\{(a, b) \in K^2 \mid P(a, b) = 0 \text{ \& } Q(a, b) \neq 0\}$, where $P, Q \in K[x, y]$ such that $\frac{\partial}{\partial y}P$ is invertible in the localization of $K[x, y]/(P)$ at Q . In the terminology of [Pool17, Definition 3.5.38] these sets are precisely the images of K -rational points of **standard étale morphisms** defined over K with codomain \mathbb{A}^1 . The standard étale open sets form a basis of a topology on K which is the **étale open topology**, cf. [Pool17, Definition 3.5.38], [JTWY24, p. 4037].*
- (ii) [JTWY24] *The field K is large if and only if the étale open topology is not discrete. If K is algebraically closed, then the étale open topology is the Zariski topology. If K is real closed, then the étale open topology is the order topology. If K possesses a nontrivial henselian valuation v , then the étale open topology is the open ball topology of v .*

Proposition 5.6 *Let L/K be a field extension such that L is large and let d be a derivation of K . Let S be an L -simple differential K -algebra. If K is algebraically closed in L and C_K is dense in L for the étale open topology of L , then the constant field of $\text{qf}(S)$ is C_K .*

Proof By L -simplicity, we may assume that S is a K -subalgebra of L . We write $F = \text{qf}(S)$ and δ for the derivation of F . Since S is L -simple, there is $g \in S$ such that the localisation $\text{Sped}_L(S_g)$ only consist of the zero ideal.

Now suppose $f = \frac{p}{q} \in F$ is a constant, thus $\delta(f) = 0$. We aim to show that f is in C_K . We work in the localization $S_{g \cdot q}$, and we view it as the coordinate ring of an affine variety V defined over K . Then f yields an algebraic map $f : V(L) \rightarrow L$. The image $W = f(V(L))$ ³ is K -definable in L in the language of rings.

Case 1. W is infinite.

Since W is an existentially L -definable set it must have nonempty interior for the étale open topology of L by [WY23, Corollary A, p.613]. Since C_K is dense in L , there is $\varepsilon \in V(L)$ —hence a K -algebra homomorphism $S_{g \cdot q} \rightarrow L$ —such that $c := \varepsilon(f) \in C_K$ and we claim that $f = c$.

³Formally: W is the set of all $\varepsilon(f)$, where $\varepsilon : S_{g \cdot q} \rightarrow L$ is a K -algebra homomorphism.

Since $\delta(f - c) = 0$, the ideal $\mathfrak{a} := (f - c)$ of $S_{g,q}$ is differential and contained in the kernel \mathfrak{q} of ε . Choose any extension of the derivation of K to L and let $T_\varepsilon^* : S_{g,q} \rightarrow L[[t]]$ be the twisted Taylor morphism of ε and that derivation. We write $\mathfrak{q}^\#$ for the kernel of T_ε^* and obtain a differential K -algebra embedding $S_{g,q}/\mathfrak{q}^\# \hookrightarrow L[[t]]^{[4]}$. It follows that the K -variety V_1 defined by $S_{g,q}/\mathfrak{q}^\#$ has a smooth $L[[t]]$ -rational point. Since L is a large field, V_1 also has a smooth L -rational point. Since S is a K -subalgebra of L we see that $\text{tr. deg}(L/K) \geq \text{tr. deg}(S/K) \geq \text{tr. deg}((S_{g,q}/\mathfrak{q}^\#)/K)$. By [Feh11, Theorem 1] we know that there is a K -algebra embedding $S_{g,q}/\mathfrak{q}^\# \hookrightarrow L$. We have shown that $\mathfrak{q}^\#$ is in $\text{Sped}_L(A)$. But (0) is the only differential L -prime ideal of $S_{g,q}$, thus $\mathfrak{q}^\# = (0)$. On the other hand, \mathfrak{a} is a differential ideal and contained in the kernel \mathfrak{q} of ε . This implies $\mathfrak{a} \subseteq \mathfrak{q}^\# = (0)$, showing that $f - c = 0$ as required.

Case 2. W is finite.

Since V is irreducible and defined over K the assumption that K is algebraically closed in L , implies that the variety $V \times_K L$ is also irreducible. Consequently $V(L)$ is an irreducible subset for the Zariski topology of the L -rational points of V . It follows that the image W of f seen as a map $V(L) \rightarrow L$ is also irreducible. As W is finite, W is a singleton set. Hence f is a constant algebraic map and thus $f = c$. ■

Corollary 5.7 *Let L/K be an extension of differential fields and suppose L is large as a field. Suppose K is algebraically closed in L and C_K is dense in L for the étale open topology of L . Let $a \in L^n$ and $g \in K\{x\}$ with $g(a) \neq 0$.*

- (i) *There is an L -simple differential K -algebra S such that the fraction field of S has constant field C_K , together with a differential K -algebra homomorphism $\varphi : K\{a\} \rightarrow S$ satisfying $g(\varphi(a)) \neq 0$.*
- (ii) *If L is differentially large, then there is some $b \in L^n$ with $g(b) \neq 0$ such that $C_{K\langle b \rangle} = C_K$ together with a differential K -algebra homomorphism $K\{a\} \rightarrow K\{b\}$ mapping a to b .*

Proof (i) Since (0) is a differential L -prime ideal, there is a differential L -prime ideal \mathfrak{p} of $K\{a\}$ with $g \notin \mathfrak{p}$ and some $s \in K\{a\} \setminus \mathfrak{p}$ such that $S = (K\{a\}/\mathfrak{p})_{g(a) \cdot s}$ is L -simple. In particular, S can be embedded as a K -algebra into L . By Proposition 5.6, the fraction field of S has constant field C_K .

(ii) follows from (i) and Observation 5.3. ■

Theorem 5.8 *Let L/K be a field extension such that L is large with $\text{tr. deg}(L/K) \geq \text{card}(K)$. Let d be a derivation of K whose constant field C is dense in L for the étale open topology of L . Then there is an extension δ of d on L such that (L, δ) is differentially large whose constant field is algebraic over C .*

Proof Using Corollary 5.7 we adapt the strategy of the proof of Theorem 4.3. We may replace K by its algebraic closure of K in L , hence we need to find a derivation on L extending d with constant field C . Let $\kappa = \text{card}(K)$. By extending K with sufficiently many differentially algebraic independent elements, we may assume that $\text{tr. deg}(L/K) = \kappa$; this will not extend the constants as one verifies without difficulty.

⁴We only need the derivation on L to obtain some K -algebra embedding as asserted, the classical Taylor morphism would not deliver this.

Let $(f_i, g_i)_{i \in \kappa}$ be a list of all differentially large problems of L , where $f_0 = x, g_0 = 1$; so here $f_i, g_i \in L\{x\}$ in the terminology of 4.1.

For $i < \kappa$, starting with $K_0 = K$, we define a subfield K_i of L that is algebraically closed in L and a derivation d_i of K_i such that

- (a) (K_i, d_i) extends (K_j, d_j) for $j < i$.
- (b) $\text{tr. deg}(K_i/K) \leq \max\{\aleph_0, \text{card}(i)\}$.
- (c) $\text{tr. deg}(L/K_i)$ is infinite (which is implied by (b) when L is uncountable).
- (d) $f_i, g_i \in K_i\{x\}$ and (K_i, d_i) solves the differentially large problem (f_i, g_i) .
- (e) (K_i, d_i) has constant field C .

Suppose $0 < i < \kappa$ and (K_j, d_j) has already been defined for $j < i$ with properties (a)–(e). Let $K_* = \bigcup_{j < i} K_j$ with derivation $d_* = \bigcup_{j < i} d_j$. Obviously then $\text{tr. deg}(K_*/K) \leq \max\{\aleph_0, \text{card}(i)\}$. If i is infinite, then L is uncountable and of size $> \text{card}(i)$, hence $\text{tr. deg}(L/K_*)$ is infinite. If i is finite, then $K_* = K_{i-1}$ and $\text{tr. deg}(L/K_*)$ is infinite as well.

Since $\text{tr. deg}(L/K_*)$ is infinite, there is a countable infinite set $T \subseteq L$ that is algebraically independent over K_* such that $\text{tr. deg}(L/K_*(T))$ is infinite, such that $f_i, g_i \in K_*(T)^{\text{alg}}\{x\}$ and such that there is an algebraic solution of the differentially large problem (f_i, g_i) in $K_*(T)^{\text{alg}}$ (here the superscript “alg” stands for the algebraic closure in L).

It follows that $\text{tr. deg}(L/K_*(T)) \geq \text{card}(K_*(T))$ in either case. Now the field $K_*(T)$ is isomorphic to the fraction field $K_*\langle x \rangle$ of the differential polynomial ring $K_*\{x\}$ and therefore there is a derivation ∂ of $K_*(T)$ extending d_* such that $(K_*(T), \partial)$ is K_* -isomorphic to $K_*\langle x \rangle$ with its natural derivation. It follows that the constant field of ∂ is the constant field of K_* , which is C by property (e) in the induction hypothesis. Hence we may extend ∂ to the algebraic closure K_+ of $K_*(T)$ in L without extending the constants.

Since $\text{tr. deg}(L/K_+) \geq \text{card}(K_+)$ we may now apply Theorem 4.3 and extend ∂ to a derivation on L such that (L, ∂) is differentially large. Since C is dense in L it is also dense in K_+ . Since (f_i, g_i) is a differentially large problem of K_+ by choice of T , we may apply Corollary 5.7(ii), which shows that (f_i, g_i) has a differential solution a in (L, ∂) such that $(K_+\langle a \rangle, \partial)$ has constant field C . We may then define K_i to be the algebraic closure of $(K_+\langle a \rangle, \partial)$ in L and see that all conditions (a)–(e) are satisfied.

Finally, $L = \bigcup_{i < \kappa} K_i$ because for each $b \in L$, the differentially large problem $(x - b, 1)$ is solved in the union. Hence the theorem follows. ■

From the description of the étale open topology in 5.5, we see that every differentially large field has a dense constant field for that topology, hence the density assumption in Theorem 5.8 is necessary.

Corollary 5.9

- (i) For every real closed subfield K of the field \mathbb{R} with $\text{tr. deg}(\mathbb{R}/K) \geq 2^{\aleph_0}$ and any derivation d of K there is a derivation δ on \mathbb{R} extending d such that (\mathbb{R}, δ) is a CODF whose constant field is the constant field of (K, d) . Recall that the étale open topology of \mathbb{R} is the Euclidean topology of \mathbb{R} .
- (ii) For every p -adically closed subfield K of the field \mathbb{Q}_p with $\text{tr. deg}(\mathbb{Q}_p/K) \geq 2^{\aleph_0}$ and any derivation d of K there is a derivation δ on \mathbb{Q}_p extending d such that (\mathbb{Q}_p, δ)

- is differentially large, whose constant field is the constant field of (K, d) . Recall that the étale open topology of \mathbb{Q}_p is the valuation topology of \mathbb{Q}_p and \mathbb{Q} is dense in \mathbb{Q}_p .
- (iii) For every pair $K \subseteq L$ of countable fields, if L is large, K is algebraically closed in L and dense in L for the étale open topology of L with $\text{tr. deg}(L/K)$ infinite, then there is a derivation δ on L such that (L, δ) is differentially large with constant field K .

Example 5.10 Dense pairs of fields $C \subseteq L$ as required in 5.8 – hence L is large, C is dense for the étale open topology of L with $\text{tr. deg}(L/C) \geq \text{card}(C)$ —also occur naturally in power series fields: Let k be any field of characteristic 0. We work in the Henselian valued field $k((t^{\mathbb{Q}}))$ of generalized power series of k . Let C be the algebraic closure of $k(t)$ in $k((t^{\mathbb{Q}}))$ and let M be the completion of C for the valued field C , hence M is the subfield of $k((t^{\mathbb{Q}}))$ consisting of power series whose support is cofinal in \mathbb{Q} and of order type ω .

Now for any \mathbb{Q} -linearly independent set $\Lambda \subseteq k$ of cardinality $\text{card}(k)$ (a baby version of) Ax's solution to the functional Schanuel conjecture implies that the series $\exp(\lambda \cdot t)$ with $\lambda \in \Lambda$ are algebraically independent over k .

So if L is a large subfield of M (for example, M itself), containing all the $\exp(\lambda \cdot t)$, then the pair L/C has the required properties.

A further consequence of Theorem 5.8 is an application to dense pairs of large fields.

Theorem 5.11 Let T be a theory of large fields of characteristic 0 in the language \mathcal{L} of rings. Let T_{pair} be the $\mathcal{L}(P)$ theory of proper pairs $K \subsetneq L$ of models of T for which K is algebraically closed in L and K is dense in the étale open topology of L ; here P is a new unary predicate.⁵ Let T^δ be the theory T together with the theory of differentially large fields in the language $\mathcal{L}(\delta)$ of ordinary differential rings.

If T^δ is a complete theory, then the theory T_{pair} is also complete.

Proof If $(L, K) \models T_{\text{pair}}$, then a standard compactness argument shows that there is an elementary extension $(L', K') \succ (L, K)$, such that $\text{tr. deg}(L'/K')$ is infinite. By the downwards Skolem–Löwenheim theorem there is a countable elementary restriction $(L'', K'') \prec (L', K')$ such that $\text{tr. deg}(L''/K'')$ is infinite.

Now take $(L_1, K_1), (L_2, K_2) \models T_{\text{pair}}$. In order to show that the pairs (L_1, K_1) and (L_2, K_2) are elementary equivalent we may apply the argument above to each pair and assume that L_i is countable and of infinite transcendence degree over K_i . By Corollary 5.9, we may expand L_i to a differentially large field (L_i, δ_i) with constant field K_i . Hence the completeness of T^δ implies that (L_1, δ_1) and (L_2, δ_2) are elementary equivalent as differential fields and therefore the pairs (L_1, K_1) and (L_2, K_2) are elementary equivalent as well. ■

Corollary 5.12 Let T be a complete and model complete theory of large fields of characteristic 0 in the language \mathcal{L} of rings, possibly expanded by constants. Let T_{pair} be the theory of proper (étale-)dense pairs of models of T . Then T_{pair} is complete and the following neostability theoretic properties transfer from T to T_{pair} :

⁵Notice that all these conditions are first order in the language $\mathcal{L}(P)$, even when the étale open topology is not a definable field topology. Density of the subfield is preserved by elementary equivalence, as one sees by using the description of the open sets in 5.5.

- (i) if T is stable, then T_{pair} is the theory of (beautiful) pairs of ACF_0 ,
- (ii) if T is simple, then T_{pair} is simple,
- (iii) if T is NSOP_1 , then so is T_{pair} , and
- (iv) if T is NIP, then so is T_{pair} .

Proof By [LST24, Corollary 4.8(iii)], T^δ is complete and model complete (recall that T^δ is the theory T together with the theory of differentially large fields). In particular, Theorem 5.11 yields that T_{pair} is complete. Now, (i) follows from the fact that a large field of characteristic zero with a stable theory must be algebraically closed [JTWY24]. For (ii), by the argument in the proof of Theorem 5.11, it suffices to observe that T^δ is simple whenever T is simple. This appears in [ML24, Corollary 3.6(i)]. Similarly, from [ML24, Corollary 3.6(ii)] we get that T^δ is NSOP_1 whenever T is NSOP_1 , and so (iii) follows. By [Moh24, Corollary 5.1], T^δ has NIP if T has NIP, hence we obtain (iv). ■

Remark 5.13 For the case of real closed fields, the completeness result of Corollary 5.12 is A. Robinson's theorem saying that the theory of proper pairs of dense real closed fields is complete. More generally, under the assumptions of Corollary 5.12, the theory T_{pair} coincides with the theory of geometrically-dense pairs of models of T in the sense of Fornasiero [For11, Section 8]. Indeed, since T is a complete and model-complete theory of large fields (in the ring language), then T is geometric (T is very slim by [JK10, Theorem 5.4] and very slim implies geometric by [JY23, Corollary 2.9]); and so it suffices to show that (in this context) étale-denseness coincides with geometric-denseness. Recall that K is geometrically-dense in L iff K intersects every infinite definable subset of L (see [For11, Example 7.2]). It is then clear that geometric-denseness implies étale-denseness. But then completeness of T_{pair} (by Corollary 5.12) implies that the two notion of denseness must coincide.

6 Prime models in CODF are algebraic

We now apply Theorem 4.3 to answer a question about prime model extensions for CODF. Recall that a CODF in the sense of Singer (cf. [Sin78]) is the same as a differentially large field that is real closed as a pure field. In [Sin78], Singer shows that CODF has no prime model, i.e., there is no CODF that embeds into all other CODF s.⁶ We now show that in fact no differential and formally real field (i.e., it possesses an ordering) has a prime model extension for CODF⁷, unless its real closure is already a CODE. In particular, no formally real field equipped with the trivial derivation has a prime model extension in CODF. The proof is essentially an application of Theorem 4.3 together with the following purely field theoretic fact.

Proposition 6.1 *Let R be a real closed field and let κ be its cardinality. Then, there are real closed fields M, N containing R of transcendence degree κ over R with the following property: If $S \supseteq R$ is a real closed field then S can be embedded over R into M and into N if and only if $\text{tr. deg}(S/R) \leq 1$ and R is Dedekind complete in S .*

⁶Note that CODF is model complete in the language of differential rings, i.e., every embedding of CODF s is elementary.

⁷A prime model extension of K for CODF is a model \hat{K} of CODF having K as a differential subfield such that \hat{K} embeds over K as a differential field into any other CODF that has K as a differential subfield.

Proof We take $M \supseteq R$ by successively adjoining infinitely large elements a_α for $\alpha < \kappa$. Hence $a_\alpha > R(a_\beta \mid \beta < \alpha)$ in M and M is algebraic over $R(a_\alpha \mid \alpha < \kappa)$. Then R is Dedekind complete in M and M has transcendence degree κ over R .

For N we may take any real closed subfield of $R((t^{\mathbb{R}}))$ of transcendence degree κ over R . Such fields exist because of the following reason: Let Λ be a basis of the \mathbb{Q} -vector space R . Since R is real closed, the cardinality of Λ is κ . Then the set $\{\exp(\lambda \cdot t) \mid \lambda \in \Lambda\} \subseteq R[[t]]$ is an algebraically independent subset of $R((t^{\mathbb{R}}))$ over R : this is a baby case of Ax's positive solution to the functional Schanuel conjecture, but is not difficult to prove directly. Hence we may take N as the real closure of $R(\exp(\lambda \cdot t) \mid \lambda \in \Lambda)$ in $R((t^{\mathbb{R}}))$. Clearly N has transcendence degree κ over R .

Since R is Dedekind complete in M and in N , any real closed field S containing R with $\text{tr. deg}(S/R) \leq 1$ in which R is Dedekind complete, can be embedded into M and into N . It remains to show that any real closed subfield S of M containing R that can be embedded into N over R is of transcendence degree at most 1 over R ; note that R is Dedekind complete in S because R is Dedekind complete in M (and in N).

For a contradiction, suppose S has transcendence degree 2 over R . We furnish M with the valuation whose valuation ring is the convex hull of R in M . Real closures are now taken in M throughout and this is indicated by the superscript rcl . Take $\bar{a} = (a_{\alpha_1}, \dots, a_{\alpha_n})$, $\alpha_1 < \dots < \alpha_n$ such that $S \subseteq R(\bar{a})^{\text{rcl}}$. Then by choice of the a_α the chain $R \subseteq R(a_{\alpha_1})^{\text{rcl}} \subseteq \dots \subseteq R(a_{\alpha_1}, \dots, a_{\alpha_n})^{\text{rcl}}$ witnesses that the value group of $R(a_{\alpha_1}, \dots, a_{\alpha_n})^{\text{rcl}}$ has height n , where height stands for the number of convex subgroups of the value group. Since $\text{tr. deg}(S/R) = 2$, there are $n - 2$ elements b_1, \dots, b_{n-2} from $\{a_{\alpha_1}, \dots, a_{\alpha_n}\}$ that are algebraically independent over S . Since S can be embedded into $R((t^{\mathbb{R}}))$ we know that S has height 1: Crucially we use here that any such embedding preserves the valuations because the natural valuation on $R((t^{\mathbb{R}}))$ again has the convex hull of R in $R((t^{\mathbb{R}}))$ as its valuation ring. But now the chain $R \subseteq S \subseteq S(b_1)^{\text{rcl}} \subseteq \dots \subseteq S(b_1, \dots, b_{n-2})^{\text{rcl}} = R(a_{\alpha_1}, \dots, a_{\alpha_n})^{\text{rcl}}$ witnesses that the value group of $R(a_{\alpha_1}, \dots, a_{\alpha_n})^{\text{rcl}}$ has height at most $n - 1$, which gives the desired contradiction. ■

Theorem 6.2 *Let K be a differential and formally real field. If K has a prime model extension \hat{K} for CODF, then \hat{K} is algebraic over K .*

Proof Suppose there is a prime model extension \hat{K} of K for CODF but \hat{K} is not algebraic over K . Let R be the algebraic closure of K in \hat{K} . Then R is a differential subfield of \hat{K} and \hat{K} is also a prime model of R for CODF: If $R \subseteq M \models \text{CODF}$, then any K -embedding $\hat{K} \rightarrow M$ must be the identity on R . Hence we may assume that K is real closed all along.

Choose real closed fields M, N for K as in Proposition 6.1. By Theorem 4.3 there are extensions of the derivation of K to M, N respectively such that M, N furnished with these extensions are CODF s. Since \hat{K} can be embedded into M and into N by assumption, Proposition 6.1 implies that \hat{K} must be of transcendence degree ≤ 1 over K and K is Dedekind complete in \hat{K} . As $K \neq \hat{K}$, we know that $\text{tr. deg}(\hat{K}/K) = 1$.

Since \hat{K} is a CODF it follows that \hat{K} has a positive infinitesimal element t with respect to K such that $t' = 1$ (in particular $t \notin K$). Then \hat{K} is a differential subfield of $K((t^{\mathbb{Q}}))$ (endowed with the derivation extending the one on K and satisfying $t' = 1$).

By [LST24, end of 5.3] we know that t^{-1} has no integral in $K((t^{\mathbb{Q}}))$. This contradicts the fact that t^{-1} has an integral in the CODF \hat{K} . ■

Remark 6.3 The proofs of Proposition 6.1 and Theorem 6.2 can be adapted to get the analogous statements about differential and formally p-adic fields and the class of p-adically closed differentially large fields. One possible task for future work is to extend Theorem 6.2 (or rather Proposition 6.1) to topological differential fields in the sense of [GP10]. We do not know if there is a version of Theorem 6.2 outside of that context. For example, if K is a subfield of a pseudo-finite field and d is a derivation of K , it is unclear whether there is a prime model over (K, d) in the class of differentially large and pseudo-finite fields (all of characteristic zero).

Acknowledgements We thank Antongiulio Fornasiero for pointing out 5.13 and the anonymous referee for thoroughly proof-reading the paper as well as for providing 3.9.

References

- [DL84] J. Denef and L. Lipshitz, *Power series solutions of algebraic differential equations pdf*. Math. Ann. 267(1984), no. 2, 213–238.
- [Feh11] A. Fehm, *Embeddings of function fields into ample fields pdf*. Manuscripta Math. 134(2011), nos. 3–4, 533–544.
- [For11] A. Fornasiero, *Dimensions, matroids, and dense pairs of first-order structures*. Ann. Pure Appl. Logic 162(2011), no. 7, 514–543.
- [GP10] N. Guzy and F. Point, *Topological differential fields pdf*. Ann. Pure Appl. Logic 161(2010), no. 4, 570–598.
- [IL23] K. Ino and O. L. Sánchez, *Separably differentially closed fields*. Preprint, 2023, arXiv e-prints. <https://arxiv.org/abs/2302.11319>.
- [Jac64] N. Jacobson. *Lectures in abstract algebra*. In: F. W. Gehring, P. R. Halmos, and C. C. Moore (eds.), *Theory of fields and Galois theory*. Vol. 3, Van Nostrand, Princeton, NJ, 1964, 332 pp.
- [Jar11] M. Jarden. *Algebraic patching pdf*, Monographs in Mathematics, Springer, Heidelberg, 2011.
- [JK10] M. Junker and J. Koenigsmann, *Schlanke Körper (slim fields)*. J. Symb. Logic 75(2010), no. 2, 481–500.
- [JRR95] J. Johnson, G. M. Reinhart, and L. A. Rubel, *Some counterexamples to separation of variables pdf*. J. Differ. Equ. 121(1995), no. 1, 42–66.
- [JTWY24] W. Johnson, C.-M. Tran, E. Walsberg, and J. Ye, *The étale-open topology and the stable fields conjecture pdf*. J. Eur. Math. Soc. 26(2024), no. 10, 4033–4070.
- [JY23] W. Johnson and J. Ye, *A note on geometric theories of fields*. Model Theory 2(2023), no. 1, 121–132.
- [Kol73] E. R. Kolchin, *Differential algebra and algebraic groups pdf*. Vol. 54, Pure and Applied Mathematics, Academic Press, New York, 1973.
- [Kol74] E. R. Kolchin, *Constrained extensions of differential fields*. Adv. Math. 12(1974), 141–170.
- [KP05] P. Kowalsi and A. Pillay, *Quantifier elimination for algebraic d-groups pdf*. Trans. Amer. Math. Soc. 358(2005), no. 1, 167–181.
- [LR86] L. Lipshitz and L. A. Rubel, *A gap theorem for power series solutions of algebraic differential equations pdf*. Amer. J. Math. 108(1986), no. 5, 1193–1213.
- [LST24] O. L. Sánchez and M. Tressl, *Differentially large fields pdf*. Algebra Number Theory 18(2024), no. 2, 249–280.
- [Mah30] K. Mahler, *Arithmetische eigenschaften einer klasse transzendental-transzendenter funktionen*. Mathematische Zeitschrift 32(1930), 545–585.
- [ML24] S. Mohamed and O. L. Sánchez, *Neostability transfers in derivation-like theories*. Preprint, 2024. arXiv e-prints. <https://arxiv.org/abs/2409.11248>.
- [Moh24] S. Mohamed, *The uniform companion for fields with free operators in characteristic zero*, Preprint, 2024. arXiv e-prints. <https://arxiv.org/abs/2311.01856>.

- [Ng23] G. Ng, *Taylor morphisms*. Preprint, 2023. <https://arxiv.org/abs/2308.11731.pdf>.
- [Pool17] B. Poonen, *Rational points on varieties pdf*. Vol. 186, Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2017.
- [Pop10] F. Pop, *Henselian implies large pdf*. Ann. of Math. (2) 172(2010), no. 3, 2183–2195.
- [PP98] D. Pierce and A. Pillay, *A note on the axioms for differentially closed fields of characteristic zero pdf*. J. Algebra 204(1998), no. 1, 108–115.
- [Ros74] M. Rosenlicht, *The nonminimality of the differential closure pdf*. Pacific J. Math. 52(1974), 529–537.
- [Sin78] M. F. Singer, *The model theory of ordered differential fields pdf*. J. Symb. Logic 43(1978), no. 1, 82–91.
- [Tre05] M. Tressl, *The uniform companion for large differential fields of characteristic 0 pdf*. Trans. Amer. Math. Soc. 357(2005), no. 10, 3933–3951.
- [Woo73] C. Wood, *The model theory of differential fields of characteristic $p \neq 0$ pdf*. Proc. Amer. Math. Soc. 40(1973), 577–584.
- [WY23] E. Walsberg and J. Ye, *Éz fields pdf*. J. Algebra 614(2023), 611–649.

The University of Manchester, Department of Mathematics, Oxford Road, Manchester, M13 9PL, UK
e-mail: marcus.tressl@manchester.ac.uk omar.sanchez@manchester.ac.uk