Research

Are you online?

ECP0014

Internet and psychosocial interventions: What is the evidence?

H. Riper

Clinical, Neuro And Developmental Psychology, Vrije Universiteit, Amsterdam, Netherlands
doi: 10.1192/j.eurpsy.2021.207

Abstract Body: Digital mental health, before, during and after the COVID-19 pandemic Over the last two decades the digital landscape of mental health care research and service innovation has gained momentum. This period is characterized by many successes' stories but brilliant failures as well. Today, e-mental health is like a two-headed Janus. One side of his face illustrates the birth of innovative technologies that entered mental health service provision. In parallel, the evidence-base for the application of these new technologies, such as internet-based treatments for depression, has been established with effect sizes comparable to those of face-to-face treatments. The other side of his face shows, however, that eMental-health has not yet lived up to its' full potential as its actual delivery, evaluation and implementation in routine care has proven to be a much longer and bumpier road than expected. The question addressed in this presentation will be 'what does the future hold'? Acknowledging that futures are difficult to predict Heleen Riper nevertheless provides insights into how we may overcome some of these bumps by combining the best of two worlds, i.e. blending digital and face-to-face components into one integrated treatment approach. She will illustrate these new developments by virtue of the results of the H2020 European Comparative Effectiveness Study for Major Depression and current experiences with especially videoconferencing during the Covid-19 pandemic.

Disclosure: No significant relationships.

ECP0011

Social isolation and physical distance: Experiences from a phone pal

M. Pinto Da Costa

Unit For Social And Community Psychiatry (who Collaborating Centre For Mental Health Services Development), Queen Mary University of London, London, United Kingdom
doi: 10.1192/j.eurpsy.2021.208

Abstract Body: People with psychosis are commonly socially isolated, both due to their condition, and the stigma towards them. Remote volunteering over smart-phone can be a way to overcome social isolation and physical distance, promoting social inclusion. This talk will present the qualitative findings from a feasibility study – the Phone Pal – which connected in the United Kingdom patients with psychosis with community volunteers, to communicate with each other for up to 12 weeks via smart-phone (through texts, WhatsApp messages, e-mails, audio or video calls). Participants described at the end of the study their experiences of communicating

with their match over the smart-phone in terms of frequency, duration and timing of communication, their communication method, content and style, and the changes of communication over time. Several participants reported a positive impact of being connected with someone, meeting a new person, feeling supported and feeling better, and a few described challenges, such as disappointment, guilt and burden. These interview findings show that some matches were able to develop a positive and friendly relationship, and were willing to continue to be in contact with each other beyond the study duration. It is hoped that this talk will generate a lively discussion, gathering further understanding about the potential benefits and challenges of remote volunteering over smart-phone for patients and volunteers, and its potential usefulness in the current pandemic times.

Disclosure: No significant relationships.

Mental Health Policy

Trauma and public mental health

ECP0013

Trauma and the role of the wounded healer

A. Hankir^{*1}, R. Zaman² and F. Carrick³

¹Institute Of Psychiatry, Psychology And Neuroscience, King's College London, London, United Kingdom; ²Psychiatry, University of Cambridge, Cambridge, United Kingdom and ³Neurology, University of Central Florida, Orlando, United States of America

*Corresponding Author.

doi: 10.1192/j.eurpsy.2021.209

Abstract Body: Carl Jung used the term, 'The Wounded Healer' as an archetypal dynamic to describe a phenomenon which may take place in the relationship between analyst and analysand. Jung discovered the Wounded Healer archetype in relation to himself. For Jung, 'It is our own hurt that gives the measure of our power to heal'. Indeed, recurrent themes in the autobiographical narratives of Wounded Healers is that their experiences living with trauma inspired them to become more empathetic, driven and insightful. Many report that debilitating though the symptoms of mental illness are, the stigma is far worse. In this paper we describe the inception of an innovative anti-stigma programme, 'The Wounded Healer' that blends the power of storytelling and the performing arts with psychiatry and how The Wounded Healer helps to heal the wounds that were afflicted by the trauma of stigma.

Disclosure: No significant relationships.

ECP0014

The impact of new psychosocial stressors on the mental health of young people: Results from a national multicentric study in Italy

G. Sampogna

Department Of Psychiatry, University of Campania "Luigi Vanvitelli", Naples, Italy

doi: 10.1192/j.eurpsy.2021.210

Abstract Body: The COVID-19 pandemic with the related containment measures is having a negative impact on the mental health of the general population worldwide. This event has been described as a new form of trauma, which is influencing not only physical and mental health, but also the society as a whole. Among Western countries, Italy has been one of the first severely hit by the pandemic in terms of number of cases and mortality rates. In March, 2020, strictly restrictive measures have been issued in order to contain the spread of the disease. This period has been known as “Phase one” of the national health emergency, where all not necessary activities were closed, almost 30,000 people died and almost 100,000 people were home-isolated. In this context, the COvid Mental hEalth Trial (COMET) network, including ten university Italian sites and the National Institute of Health, has been established in order to promote a national online survey for assessing the impact of lockdown measures on the mental health of the Italian general population. In the COMET survey, it has been included a specific focus on young people, which are expected to be the most vulnerable to the consequences of the pandemic and of the strict containment measures. Findings from this study can be useful to inform national and international associations on the importance to provide adequate support to the mental health of the young people.

Disclosure: No significant relationships.

ECP0015

Building resilience to early life trauma in belarus and Ukraine

O. Kazakova^{*1}, F. Thomas² and O. Suvalo³

¹Public Health, Lund University, Malmö, Sweden; ²Wellcome Centre For Cultures And Environments Of Health, University of Exeter, Exeter, United Kingdom and ³The Institute Of Mental Health, The Ukrainian Catholic University, Lviv, Ukraine

*Corresponding Author.

doi: 10.1192/j.eurpsy.2021.211

Abstract Body: Early life trauma (ELT) refers to various types of adversity that occur during the early years (usually defined as the first 5 years) of a person's life. It is a key determinant of mental health and well-being throughout the life course. A series of three workshops on early life trauma and mental health care were conducted in Belarus and Ukraine in 2018-2019 to support stakeholders and service providers to better understand and respond to ELT, and to support the development of a network of ELT specialists dedicated to finding common goals, pooling cross-disciplinary data and sharing experiences and good practice across countries. The workshops found that different attitudes, expectations and experiences amongst stakeholders and service providers could hinder the development of consistent, effective and empowering care in Belarus and Ukraine. However, opportunities for more protective and health-enhancing responses were also identified, including the need for: evidence-based education and training; clear roles and communication pathways across sectors; and inter-sectoral partnerships and networks to leverage resources, mitigate practitioner burnout, and build a continuum of support within communities. Findings have been disseminated through a directory of resources in Belarus, a project webpage (www.earlylifetrauma.info) and a report on ELT in Belarus and Ukraine published by WHO Europe.

Disclosure: No significant relationships.

ECP0016

Improving health care response to domestic violence

A. Dowrick

Nuffield Department Of Primary Care Health Sciences, University of Oxford, Oxford, United Kingdom

doi: 10.1192/j.eurpsy.2021.212

Abstract Body: 1 in 4 women in Europe will experience domestic violence and abuse (DVA) in their lifetime. Abuse can take many forms, including, psychological, physical, sexual, financial, and emotional abuse. There is growing recognition of the health consequences of DVA in public policy and academic research across Europe and the globe. These include but are not limited to: a negative impact on mental health; increased substance misuse; increased presentation in emergency departments; increased rates of abortion or miscarriage; and increased presence of any sexual health problem. International and national policy guidance indicates that healthcare professionals have important roles in responding to patients experiencing DVA, usually with regard to identifying abuse and referring for specialist support. Barriers to engaging in DVA relate to lack of confidence in recognising abuse or initiating conversations about it, stigma surrounding DVA, fears of exacerbating violence, limited awareness of the resources available to support patients, and limited training in providing trauma-informed medical care. This session will give insight into efforts to improve the healthcare response to DVA. It will draw together examples of healthcare DVA programmes from across the globe and bring together lessons learnt about what works and what does not work in addressing DVA in clinical settings, with specific insights for professionals working in mental health services.

Disclosure: No significant relationships.

Educational

What should early career psychiatrists know?

ECP0017

Training as a psychiatrist during a pandemic

C. Migchels

University Psychiatric Centre, KULeuven, Kortenberg, Belgium

doi: 10.1192/j.eurpsy.2021.213

Abstract Body: Training is a crucial time in any psychiatrist's career. It's a time when you are faced with new responsibilities and challenges, both professionally and personally, and senior and peer support are indispensable. The Covid-19 pandemic has had an impact on all the essential needs of trainees: supervision, theoretical training, clinical care and contact with both patients and colleagues have been put under pressure and are no longer the self-evident part of training that they were before. Trainees are facing an unprecedented balancing act between their duties as physicians, their training needs and their dedication to the often vulnerable patients under their care, all whilst dealing with the personal consequences of the pandemic and the restrictions that have been put in place.