

Genetic diversity analyses of *Brassica napus* accessions using SRAP molecular markers

Riaz Ahmad¹, Farhatullah^{2*}, Carlos F. Quiros¹, Hidayatur Rahman² and Zahoor Ahmad Swati³

¹Department of Plant Sciences, University of California, Davis, CA 95616, USA,

²Department of Plant Breeding and Genetics, The University of Agriculture, Peshawar,

Pakistan and ³Institute of Biotechnology and Genetic Engineering, The University of Agriculture, Peshawar, Pakistan

Received 4 February 2013; Accepted 6 June 2013 – First published online 26 July 2013

Abstract

Knowledge about genetic diversity among *Brassica napus* cultivars developed for many growing regions and their possible use as potential inbred lines for hybrid seed production is limited. We studied the genetic diversity and relationships among *B. napus* accessions using Sequence Related Amplified Polymorphism (SRAP) markers, which preferentially amplify open reading frames. A total of 60 spring-type *B. napus* accessions were screened using 20 SRAP primers, which revealed 162 polymorphic fragments with an average of eight markers per primer combination. Genetic similarity estimates ranged from 40 to 100, which indicated sufficient diversity among the accessions. The majority of the accessions were uniquely identified by the markers with the exception of near-isogenic inbred lines. Cluster analysis displayed five major groups. The first major cluster comprised 23 accessions mostly of Australian origin, whereas the second cluster included 13 accessions mostly of Canadian origin. The accessions in the first and second clusters were identified as maintainers of cytoplasmic male sterility. The two restorer lines R-111 and R-101 along with their corresponding backcross progeny constituted the third cluster. Scandinavian cultivars made the fourth separate cluster. One cultivar Salam and its respective inbred line were the most divergent lines. Variations in the number of markers between open-pollinated cultivars and their respective selfed inbred lines were also observed. The clustering pattern mostly supported their respective pedigree and characteristic traits. Genetic diversity in genetically distinct groups in the tested maintainer and restorer lines can be exploited for hybrid development in *B. napus*.

Keywords: cytoplasmic male sterility; fertility restoration; genetic diversity; oilseed rape; SRAP markers

Introduction

Oilseed rape (*Brassica napus* L.; genome AACC, $2n = 38$) is originated through spontaneous interspecific hybridizations between turnip rape (*Brassica rapa* L., syn. *campestris*; AA, $2n = 20$) and cabbage (*Brassica oleracea* L.; CC, $2n = 18$) (Palmer *et al.*, 1983; Iñiguez Luy and Federico, 2010). The oilseed rape gene pool is

subdivided into winter and spring types (Becker *et al.*, 1995; Bus *et al.*, 2011). Within winter oilseed rape types, a decay of genetic diversity with reduced levels of erucic acid and glucosinolates has been observed (Bus *et al.*, 2011). Spring type is grown extensively in Canada, Australia, Southeast Asia and Scandinavia. Canola-type cultivars (low levels of erucic acid (<2%) and glucosinolate (<30 μ M)) of *B. napus* and *B. rapa* (Downey and Rimmer, 1993) are widely grown commercially.

The prominent breeding system in Australia since 1970 has been the open-pollinated (OP) pedigree selection

*Corresponding author. E-mail: drfarhat@aup.edu.pk

(Cowling, 2007). For genetic improvement of *B. napus*, development of hybrids (having high levels of heterosis) is being preferred over OP varieties (Sernyk and Stefansson, 1983; Grant and Beversdorf, 1985; Ahmad *et al.*, 2011). Canola hybrids are getting increased market share worldwide due to a high seed yield under normal, dry and nutrient-poor conditions (Gehring *et al.*, 2007; Malhi *et al.*, 2007). In hybrid brassica breeding programmes, cytoplasmic male sterility (CMS) lines are crossed with a large number of promising breeding lines. Parents that give complete pollen sterility are identified as maintainers (M) and used for further backcrossing to CMS sources, while those parents that give 100% pollen fertility are identified as restorers (R) of sterility and are further tested for combining ability.

The knowledge of genetic diversity is indispensable in the development of commercial hybrids. In addition to morphological characterizations, molecular characterization is essential for elucidating the genetic relationship among accessions. Genetic diversity and cultivar relatedness in *Brassica* can be assessed using various tools including, isozyme and DNA-based markers such as restriction fragment length polymorphism (RFLPs), Random Amplification of Polymorphic DNA (RAPDs), simple sequence repeats and Amplified fragment length polymorphism (AFLPs) (Figdore *et al.*, 1988; Hu and Quiros, 1991; Song and Osborn, 1992; Hasan *et al.*, 2006; Chen *et al.*, 2008; Turi *et al.*, 2012).

For the present study, we selected a marker system called sequence-related amplified polymorphism (SRAP) because of its simplicity, reproducibility and disclosure of multiple markers from a single two-primer reaction (Li and Quiros, 2001; Ahmad *et al.*, 2004; Wu *et al.*, 2009; Ahmad *et al.*, 2013). SRAP molecular markers are more concordant with the morphological variability and evolutionary history of morphotypes than AFLP markers (Ferriol *et al.*, 2003).

The objectives of the present study were to assess and characterize the pattern of genetic diversity in *B. napus* OP cultivars, M and R lines, and to identify potential parents for the development of commercial hybrids using SRAP molecular markers. We also studied the extent of molecular marker variations between OP cultivars and their respective selfed inbred lines. This information would help breeders and geneticists understand the structure of *B. napus* germplasm and help them predict which parents have the best chance for cross-combination to give more heterosis.

Materials and methods

Plant material

A set of 60 *B. napus* accessions including 11 maintainers, 23 restorers and 26 OP cultivars were included in the

present study (Supplementary Table S1, available online). The OP cultivars were mostly named cultivars from Australia, Canada, Europe and Pakistan and included spring types either with or without canola quality (Supplementary Table S1, available online). The M lines were confirmed by crossing with CMS lines and then selfed for at least five to seven generations. Similarly, the R lines were confirmed after 100% restoration of CMS lines and then selfed for at least five to seven generations.

DNA extraction

The plant material was grown in pots at the green house facility at the University of California, Davis, USA. DNA was extracted from fresh leaves by the cetyltrimethylammonium bromide (CTAB) method of Doyle and Doyle (1990) with some modifications. A 300–400 mg sample was added into a 1.5 ml microcentrifuge tube and crushed into powder under liquid nitrogen. Preheated extraction buffer (500 μ l of 2% CTAB, 100 mM Tris–HCl, pH 8.0, 1.4 M NaCl, 20 mM EDTA, 1% PVP-40 and 10 mM DTT) and 10 μ l proteinase K (50 μ g/ml) were then added and the tube was incubated at 65°C for 90 min in a water bath. The tube was cooled to room temperature and 500 μ l of chloroform–isoamyl alcohol (24:1) were added. The solution was mixed gently and centrifuged at 13,000 \times g for 5 min. The supernatant was transferred to a fresh tube, 0.6 volumes of isopropanol were added and mixed gently to precipitate the DNA. The mixture was centrifuged at 13,000 \times g for 5 min, the supernatant was removed and the DNA pellet was washed with 70% (v/v) ethanol. The pellet was then air-dried and resuspended in 100 μ l of Tris–EDTA buffer. DNA quality and quantity was determined by electrophoresis in an ethidium bromide-stained 1.0% (w/v) agarose gel.

Molecular markers

A PCR-based technique, SRAP, which preferentially amplifies each open reading frame, was tested on the *B. napus* accessions (Li and Quiros, 2001). Initially, 45 SRAP primer combinations were tested on eight diverse *B. napus* accessions from different origins. Of these, 20 primer combinations were selected, as they provided the greatest reproducibility and the number of markers per gel (Table 1). Eight forward primers, with the one labelled with IRD800 or IRD700 to be detected in a Li-Cor DNA sequencer (Long Readir 4200), and 11 reverse primers were used (Table 2).

At the beginning of the PCR reaction, the annealing temperature was set at 35°C and run for five cycles.

Table 1. Sequence-related amplified polymorphism primers used for the analysis of *Brassica napus* accessions and the number of polymorphic fragments revealed by each primer combination

Primer combinations	Polymorphic bands	Total number of bands
ME1 + EM3	9	32
TE1 + EM3	10	23
ME2 + EM17	11	37
ME4 + EM17	8	24
ME3 + EM17	4	16
ME3 + EM6	7	15
ME4 + EM6	7	14
ME3 + EM18	11	25
TE1 + EM18	7	29
ME4 + EM18	7	22
ME2 + EM18	15	27
ME1 + EM18	9	21
ME3 + EM19	8	27
TE1 + EM20	5	14
ME1 + EM20	7	27
ME2 + EM4	9	27
ME3 + EM2	3	22
ME2 + EM1	11	21
DC1 + OD34	9	15
ME2 + SA7	5	14
Total	162	425
Average	8.1	21.3

Then, the annealing temperature was raised to 50°C for another 35 cycles. Denaturing was done at 94°C for 1 min, while extension was at 72°C for 1 min in both the first five cycles and the last 35 cycles. Thereafter, 1 µl of the reaction products was mixed with 6 µl of formamide dye, denatured at 94°C for 3 min and rapidly cooled on ice for 5 min. Amplicons were separated in denaturing 5.5 % (w/v) polyacrylamide gels (0.25 mm spacer thickness) and detected by a DNA sequencer (Gene Reader 4200; Li-Cor). Only bright, clearly distinguishable bands were used in the genetic analysis. Apparent polymorphisms appearing with band sizes below 50 bp or above 600–700 bp were rejected due to

the inconsistency of amplification and the weakness of bands. To test the reproducibility of the SRAP banding patterns, two independent PCR amplifications were performed for each of the same samples and run side by side.

Data collection and analysis

Only fragments that were polymorphic among the accessions and could be clearly scored were used in the data analysis. Each of these fragments was scored independently. The fragment data were entered into a spreadsheet to form a 0–1 matrix, with 1 representing the presence of a fragment and 0 representing the absence of a fragment for each fragment accession combination. Genetic similarity (GS) among the accessions was calculated according to the method of Nei and Li (1979):

$$GS_{ij} = 2N_{ij}/(N_i + N_j),$$

where N_{ij} is the number of bands in common between lines i and j , and N_i and N_j are the total number of bands in lines i and j , respectively. Thus, GS_{ij} reflects the proportion of bands in common between two accessions and may range from 0 (no common bands) to 1 (identical profiles of two lines). Cluster analysis based on the unweighted pair-group method with arithmetic average (UPGMA; Sneath and Sokal, 1973) was performed on the matrix of GS estimates and a dendrogram constructed using NTSYS pc 2.0. Bootstrap values of 1000 were calculated and only numbers above 500 were indicated on the tree branches.

Results

B. napus accessions including 11 maintainers, 23 restorers and 26 OP cultivars (Supplementary Table S1,

Table 2. List of eight forward primers, with the one labelled with IRD800 or IRD700 to be detected in a Li-Cor DNA sequencer, and 11 reverse primers

Forward primers (5'–3')		Reverse primers (5'–3')	
ME1	TGAGTCCAAACCGGATA	EM1	GACTGCGTACGAATTCAAT
ME2	TGAGTCCAAACCGGAGC	EM2	GACTGCGTACGAATTCTGC
ME3	TGAGTCCAAACCGGAAT	EM3	GACTGCGTACGAATTCTGC
ME4	TGAGTCCAAACCGGACC	EM4	GACTGCGTACGAATTCTGA
ME8	TGAGTCCAAACCGGTGC	EM6	GACTGCGTACGAATTCTGA
DC1	TAAACAATGGCTACTCAAG	EM17	GACTGCGTACGAATTCTGA
TE1	TGTGTGGTTAATATGAGC	EM18	GACTGCGTACGAATTCTGA
		EM19	GACTGCGTACGAATTCTGA
		EM20	GACTGCGTACGAATTCTCC
		SA7	CGCAAGACCCACCACAA
		OD30	GCGATCACAGAAGGAAGGT

available online) were evaluated for genetic diversity study using 20 SRAP primer combinations (Table 1). These primers were selected because they revealed polymorphisms among the eight *B. napus* accessions in a preliminary screening. The 20 SRAP primer combinations produced sufficient polymorphism in the tested 60 accessions, which was used for calculating similarity coefficient. A total of 425 bands ranging in size from 50 to 650 bp, with an average of 21 bands per individual per primer combination were produced, of which 162 were polymorphic with an average of eight polymorphic bands per primer combination (Table 1). The primer combination Me2 + Em18 was highly effective in generating polymorphism (15 polymorphic bands in the 60 lines tested), whereas the combination Me3 + Em2 produced only three polymorphic bands. The majority of the accessions were uniquely identified by the markers with the exception of near-isogenic M lines and their corresponding OP cultivars. No marker difference was observed between the two R inbred lines R-159 and R-16. The percentage of polymorphic fragments was greater in the M lines since these were mostly originated from diverse sources.

The molecular data of 162 markers were used to calculate genetic similarity coefficient for all pairs of accessions and a phenogram was constructed from these values with the UPGMA (Fig. 1). Genetic similarity estimates calculated among the accessions varied from 0.40 to 0.97 for all the accessions except for two identical R inbred lines R-159 and R-161. Cluster analysis placed all the accessions into five main groups (Fig. 1). The first group (cluster 1) included mainly Australian OP cultivars and some of their respective selfed inbred lines. This cluster was further divided into three sub-clusters (I-a, I-b and I-c). Sub-cluster I-a included six canola-type M lines (Dunkeld, Dunkeld-S6, Bulbul, Bulbul-S6, Rainbow and Rainbow-S6) and one R inbred line (R-115). Sub-cluster I-b included two restorer lines (R-128 and R-117) of the same origin and ten accessions of both canola and non-canola types (Oscar, Oscar-S6, Maluka, Maluka-S6, Eureka, Marnoo, Taparoo, Altex, Altex-S6 and Galaxy) of Australian origin and identified as the M of CMS. Sub-cluster I-c included only four canola-type M lines (Shiralee, Shiralee-S5, Range and Narendra). The second cluster mainly included accessions of Canadian origin and was further divided into two sub-clusters: II-a (Siren, Tower, Tower-S5, Triton, Siren-S6) and II-b (Wester, Wester-S7, Regent, Golden and Excel) and two R lines (R-110 and R-121). The third cluster mainly included R lines and was further sub-divided into two sub-clusters: III-a (R-111, R-111-1, R-111-2, R-111-3, R-111-4, R-111-10, R-111-5, R-111-7 and R-111-9) and III-b (R-101, R-101-1, R-101-2, R-101-3, R-101-4 and R-101-9). The fourth cluster included mostly OP cultivars

of Scandinavian origin from Europe (Topas, Global, Crystal, Stallion and Delta). The last two accessions 'Salam' and its respective inbred line are the most divergent and are separate from the rest of the samples.

Analyses of a set of 11 OP cultivars and their respective inbreds selfed for five to seven generations indicated a range of marker differences between the cultivars and inbreds (Table 3). Most of the inbreds clustered with their respective cultivars. The maximum number of markers was lost in the cultivar Siren followed by Salam.

Discussion

SRAP molecular markers resulted in eight markers per primer combination, which is more than the RAPD markers of four per gel as observed by Mailer *et al.* (1994) in rapeseed cultivars and less than 19.1 per gel of the AFLP markers as reported by Lombard *et al.* (2000) in spring- and winter-type accessions. The large number of markers in the AFLP study may be in part due to testing of highly divergent spring- and winter-type *Brassica* accessions. Our results are in accordance with other researchers using SRAP molecular markers (Budak *et al.*, 2004; Guo and Luo, 2006; Sun *et al.*, 2006), where they also found a moderate level of polymorphism using SRAP molecular markers.

The SRAP molecular markers were quite useful in the discrimination of *B. napus* accessions. In general, most of the lines closely related by pedigree and/or derived from germplasm having specific traits clustered together (Wu *et al.*, 2009; Ahmad *et al.*, 2013). The cluster analyses separated the accessions into five major groups. The accessions from Australia, Canada and Europe were grouped separately. The genetic diversity within the cluster was very narrow. This may be due to the increased selection pressure resulting in a number of high oil and protein content of canola-type varieties (e.g. Dunkeld, Rainbow and Oscar). Second, by looking to the pedigree of Australian accessions, they mainly contain genetic material from few cultivars such as Zephyr, Bronowski, and Chisaya and in some cases Norin20 from Japan and Cresus-o-Precose. This has made the genetic background of these cultivars very narrow and molecular markers have grouped these accessions into a single large cluster. All of the Australian and Canadian OP cultivars were M of the CMS and thus most probably carry the same cytoplasm. Diers and Osborn (1994) showed that the Australian cultivars Taparoo, Galaxy, Maluka and Marnoo clustered together as was observed in our study.

The University of Manitoba, Canada, *B. napus* cultivars 'Tower' and 'Regent', clustered together and have the same pedigree as noted by Sernyk (1990). Similarly, Mailer *et al.* (1994) noted that the Canadian cultivars




Fig. 1. (colour online) Dendrogram showing the relationships among the 60 *B. napus* maintainers, restorers and OP cultivars using SRAP molecular markers. Numbers on the branches indicate bootstrap values above 500.

Table 3. Number of marker differences between open-pollinated cultivars and their respective selfed lines (for five to seven generations)

Cultivar	S5-6 inbred	Fragments unique to cultivar ^a	Fragments unique to inbred ^b
Dunkeld	Dunkeld-S6	4	0
Bulbul	Bulbul-S6	3	0
Rainbow	Rainbow-S6	4	1
Oscar	Oscar-S6	3	1
Maluka	Maluka-S6	6	3
Altex	Altex-S6	7	1
Shiralee	Shiralee-S5	5	2
Siren	Siren-S6	10	4
Tower	Tower-S5	2	1
Wester	Wester-S7	5	0
Salam	Salam	8	5

^aNumber of markers present in the cultivar that were absent in the inbred line. ^bNumber of markers present in the inbred line that were absent in the cultivar.

Westar, Excel and Regent also clustered together using RAPD markers. Results of 43 RFLP markers of Diers and Osborn (1994) also indicated the close association of the Canadian cultivars Regent, Wester, Triton and Tower. However, their study could not separate the cultivars 'Triton' and 'Tower' from each other.

The four Scandinavian cultivars Crystal and Stallion (Sweden, Sval6f Seed Ltd), Global (Denmark, Sval6f Seed Ltd), Delta (Sweden, Pioneer Hibred International) and Topas clustered closely. These are distant from Australian accessions since none of the Australian or Canadian accessions has genetic material from these lines. Mailer *et al.* (1994) also reached the same conclusion of clustering of the Sweden cultivars, Delta, Stallion, Crystal and Global using RAPD markers and concluded that the relationships among some of the cultivars are closer for those from the same breeding programme than for those from different programmes. Similarly, the most distant accessions were the white-flowered cultivar 'Salam' and its corresponding inbred line (obtained by crossing *B. oleracea* and *B. napus*; Quazi, 1988).

The two R lines R-111 and R-101 along with their corresponding backcross and F₅₋₆ progeny made the third cluster and further divided into two sub-clusters. The accessions within these clusters display poor genetic diversity because most of them have been derived locally and one or both of their parents are common in their pedigrees. In sub-cluster III-a, one of the common parents is R-111, while for another sub-cluster (III-b), the R line R-101 is one of the parent in their pedigree. The fact that these R lines are separated from the M inbred lines could be useful for a possible hybrid combination. Restorer lines such as R-115 and R-128 grouped within the same cluster with maintainer lines and having comparatively

more genetic similarity indicated their limited utility for hybrid development within the same cluster. Restorer, exhibiting high genetic dissimilarity with most of the M lines, could be promising in hybrid development.

A set of 11 OP cultivars and their respective inbreds selfed for five to seven generations were also analysed in the present study. The inbreds were developed to be used as M or R for hybrid seed production. We used molecular markers to determine the extent of similarity between OP cultivars and their respective selfed inbreds. In addition to the morphological differences such as reduced height and loss of vigour, the marker analysis also revealed a range of marker differences between the cultivars and inbreds (Table 3). In the phylogenetic tree, most of the inbreds were near to their respective cultivars except for the inbred line from the cultivar 'Siren'. The maximum number of extra fragments was present in the cultivar 'Siren' but was not present in the inbred line. This scenario was observed for most of the inbred lines where an extra marker was present in the cultivar, but not present in the inbred line. The same observation was also reported by Diers and Osborn (1994). According to these authors, loss of fragments resulted from inbreeding when inbreds were made from heterozygous or heterogeneous cultivars. We also had a similar situation where inbreds had an extra fragment but the cultivar did not possess it. The extra marker in the inbred line might be explained by the fact that the plants selected for inbreeding contained fragments from the cultivar that were not present in those selected for SRAP analysis. The number of fragments present in the cultivar that were absent in the inbred line and that of fragments present in the inbred line that were absent in the cultivar are possible because the cultivars are somewhat heterogeneous (Diers and Osborn, 1994).

Conclusions

The present study showed that SRAP molecular markers functioned well in exploring the genetic diversity among *B. napus* accessions. The study revealed that considerable genetic diversity was detected between the maintainer and restorer inbred lines, but not within the same cluster. The SRAP molecular markers were more concordant with the morphological variability and pedigree of the lines. The recurrent parent-backcross line pairs and the cultivar inbred pairs were each clustered closely together as expected. The R lines in different clusters could be potential pollinators for the elite CMS/M lines in different clusters. Furthermore, the diversity assessed can be manipulated to broaden the genetic base of elite maintainer and CMS lines for the development of commercial hybrid varieties.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S147926211300021X>

Acknowledgements

The authors thank Victor Wutor, Lethbridge Research Center, Agri-Food Canada and Janssens Marc, University of Bonn, Germany, for technical review of the manuscript. They also thank Genyi Li Assistant Professor, Department of Plant Sciences, University of Manitoba, Canada, for his help in the SRAP molecular marker system. The authors are indebted to Doug Walker, Vince D. Antonio, University of California at Davis, USA for maintaining the plant material in the greenhouse and Yasir Khan for his valuable help in formatting the manuscript. The authors are grateful to two anonymous reviewers for their technical inputs.

References

- Ahmad R, Potter D and Southwick SM (2004) Genotyping of peach and nectarine cultivars with SSR and SRAP molecular markers. *Journal of American Society of Horticulture Science* 129: 204–210.
- Ahmad R, Farhatullah and Carlos FQ (2011) Inter- and intra-cluster heterosis in spring type oilseed rape (*Brassica napus* L.) hybrids and prediction of heterosis using SRAP molecular markers. *SABRAO Journal of Breeding and Genetics* 43: 27–43.
- Ahmad R, Farhatullah, Khan RS and Carlos FQ (2013) Inheritance of fertility restorer gene for cytoplasmic male-sterility in *B. napus* and identification of closely linked molecular markers to it. *Euphytica* (May 2013) Advance online publication. DOI: 10.1007/s10681-013-0942-y
- Becker HC, Engqvist GM and Karlsson B (1995) Comparison of rapeseed cultivars and resynthesised lines based on allozyme and RFLP markers. *Theoretical and Applied Genetics* 91: 62–67.
- Budak H, Shearman RC, Parmaksiz I, Gaussoin RE, Riordan TP and Dweikat I (2004) Molecular characterization of buffalograss germplasm using sequence-related amplified polymorphism markers. *Theoretical and Applied Genetics* 108: 328–334.
- Bus A, Körber N, Snowdon RJ and Stich B (2011) Patterns of molecular variation in a species-wide germplasm set of *Brassica napus*. *Theoretical and Applied Genetics* 123: 1413–1423.
- Chen S, Nelson MN, Ghamkhar K, Fu T and Cowling WA (2008) Divergent patterns of allelic diversity from similar origins: the case of oilseed rape (*Brassica napus* L.) in China and Australia. *Genome* 51: 1–10.
- Cowling WA (2007) Genetic diversity in Australian canola and implications for crop breeding for changing future environments. *Field Crops Research* 104: 103–111.
- Diers BW and Osborn TC (1994) Genetic diversity of oilseed *Brassica napus* germplasm based on restriction fragment length polymorphisms. *Theoretical and Applied Genetics* 88: 662–668.
- Downey RK and Rimmer SR (1993) Agronomic improvement in oilseed Brassicas. *Advances in Agronomy* 50: 1–66.
- Doyle JJ and Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12: 13–15.
- Ferriol M, Pico B and Nuez F (2003) Genetic diversity of a germplasm collection of *Cucurbita pepo* using SRAP and AFLP markers. *Theoretical and Applied Genetics* 107: 271–282.
- Figdore SS, Kennard WC, Song KM, Slocum MK and Osborn TC (1988) Assessment of the degree of restriction fragment length polymorphism in brassica. *Theoretical and Applied Genetics* 75: 833–840.
- Grant I and Beversdorf WD (1985) Heterosis and combining ability estimates in spring oilseed rape (*Brassica napus* L.). *Canadian Journal of Genetics and Cytology* 27: 472–478.
- Gehring A, Snowdon R, Spiller T, Basunanda P and Friedt W (2007) New oilseed rape (*Brassica napus*) hybrids with high levels of heterosis for seed yield under nutrient poor conditions. *Breeding Science* 57: 315–320.
- Guo DL and Luo ZR (2006) Genetic relationships of some PCNA persimmons (*Diospyros kaki* Thunb.) from China and Japan revealed by SRAP analysis. *Genetic Resources and Crop Evolution* 53: 1597–1603.
- Hasan M, Seyis F, Badani AG, Pons-Kuhnemann J, Friedt W, Lühs W and Snowdon RJ (2006) Analysis of genetic diversity in the *Brassica napus* L. gene pool using SSR markers. *Genetic Resource and Crop Evolution* 53: 793–802.
- Hu J and Quiros CF (1991) Identification of broccoli and cauliflower cultivars with RAPD markers. *Plant Cell Reports* 10: 505–511.
- Iñiguez Luy FL and Federico ML (2010) The genetics of *Brassica napus* L. In: Bancroft I and Schmidt R (eds) *Plant Genetics and Genomics Series: Genetics and Genomics of the Brassicaceae*. New York, NY: Springer Science.
- Li G and Quiros CF (2001) Sequence-related amplified polymorphism (SRAP) a new marker system based on a simple, PCR reaction: its application to mapping, and gene tagging in Brassica. *Theoretical and Applied Genetics* 103: 455–461.
- Lombard V, Baril CP, Dubreuil P, Blouet F and Zhang D (2000) Genetic relationship and fingerprinting of rapeseed cultivars by AFLP: consequences for varietal registration. *Crop Science* 40: 1417–1425.
- Mailer RJ, Scarth R and Fristensky B (1994) Discrimination among cultivars of rapeseed (*Brassica napus* L.) using DNA polymorphisms amplified from arbitrary primers. *Theoretical and Applied Genetics* 87: 697–704.
- Malhi SS, Brandt S, Ulrich D, Lafond GP, Johnston AM and Zentner RP (2007) Comparative nitrogen response and economic evaluation for optimum yield of hybrid and open-pollinated canola. *Canadian Journal of Plant Science* 87: 449–460.
- Nei M and Li W (1979) Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences USA* 79: 5269–5273.
- Palmer JD, Shields CR, Cohen DB and Orton TJ (1983) Chloroplast DNA evolution and the origin of amphidiploid *Brassica* species. *Theoretical and Applied Genetics* 65: 181–189.
- Quazi MH (1988) Interspecific hybrids between *Brassica napus* L. and *B. oleracea* L. developed by embryo culture. *Theoretical and Applied Genetics* 75: 309–318.

- Sernyk JL (1990) *Catalogue of Oilseed Rape Cultivars: 1990 Edition*. Madison, WI: Agrigenetics Company.
- Sernyk JL and Stefansson BR (1983) Heterosis in summer rape (*Brassica napus* L.). *Canadian Journal of Plant Science* 63: 407–413.
- Sneath PHA and Sokal RR (1973) *Numerical Taxonomy: The Principles and Practice of Numerical Classification*. San Francisco, CA: Freeman.
- Song K and Osborn TC (1992) Polyphyletic origins of *Brassica napus*: New evidence based on organelle and nuclear RFLP analyses. *Genome* 35: 992–1001.
- Sun SJ, Gao W, Lin SQ, Zhu J, Xie BG and Lin ZB (2006) Analysis of genetic diversity in *Ganoderma* population with a novel molecular marker SRAP. *Applied Microbiology and Biotechnology* 72: 537–543.
- Turi NA, Farhatullah, Malik AR and Zabta KS (2012) Genetic diversity in the locally collected *Brassica* species of Pakistan based on microsatellite markers. *Pakistan Journal of Botany* 44: 1029–1035.
- Wu X, Chen B, Lu G, Wang H, Xu K and Guizhan G (2009) Genetic diversity in oil and vegetable mustard (*Brassica juncea*) landraces revealed by SRAP markers. *Genetic Resources and Crop Evolution* 56: 1011–1022.