

The IACOB spectroscopic database of galactic OB stars

Sergio Simón-Díaz^{1,2}, Norberto Castro^{1,2}, Miriam Garcia^{1,2} and
Artemio Herrero^{1,2}

¹Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

²Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain

Abstract. We present the IACOB spectroscopic database, the largest homogeneous database of high-resolution, high signal-to-noise ratio spectra of Northern Galactic OB-type stars compiled up to date. The spectra were obtained with the FIES spectrograph attached to the Nordic Optical Telescope. We briefly summarize the main characteristics and present status of the IACOB, first scientific results, and some future plans for its extension and scientific exploitation.

Keywords. stars: early-type, techniques: spectroscopic, catalogs, astronomical data bases: miscellaneous

1. Introduction

In an epoch in which we count with a new powerful generation of stellar atmosphere codes including all the important physics for the modelling of massive OB stars, with (clusters of) high efficiency computers allowing the computation of large grids of stellar models in more than reasonable computational times, and with the possibility to obtain good quality, medium resolution spectra of hundreds O and B-type stars in clusters outside the Milky way in just one snapshot (see e.g. the *FLAMES I & II Surveys of Massive Stars*, Evans *et al.* 2008, 2010; see also Lennon *et al.*, these proceedings), the compilation of medium and high-resolution spectroscopic databases of OB stars in our Galaxy is becoming more and more important. With this idea in mind, two years ago we began to compile the IACOB spectroscopic database, aiming at constructing the largest database of multi-epoch, high resolution, high signal-to-noise ratio (S/N) spectra of Galactic Northern OB-type stars. The IACOB perfectly complements the efforts also devoted in the last years by the GOSSS (P.I. Maíz-Apellániz) and the OWN (P.I's Barbá & Gamen, leading a multi-epoch, high-resolution spectroscopic survey of Galactic O and WR stars in the Southern hemisphere; see Barbá *et al.* 2010) teams.

2. Characteristics of the IACOB and present status

We are using the FIES spectrograph at the 2.56 m Nordic Optical Telescope (NOT) in the Roque de los Muchachos observatory (La Palma, Spain) to compile spectra for the IACOB (see <http://www.not.iac.es>). A summary of the instrumental configuration and observing dates (before Sept. 2010) is presented in Table 1. 720 spectra of 105 stars with spectral types O4-B2 and luminosity classes ranging from I (Supergiants) to V (Dwarfs) have already been compiled. The O-type targets were selected among those stars with $V \leq 8$ included in the GOS catalogue (GOSC, Maíz-Apellániz *et al.* 2004). The main part of the B-type stars sample correspond to the works presented in Simón-Díaz (2010) and Simón-Díaz *et al.* (2010). The final spectra normally have $S/N \geq 200$.

Table 1. General characteristics of the IACOB v1.0 spectroscopic database.

Instrumental configuration		Observing run & Dates
Telescope: NOT2.56 m	Spect. range: 3800 - 7000 Å	08 A-D: 2008/11/05-08, 10 D: 2010/06/22
Instrument: FIES	Res. power: 46000	09 A-D: 2009/11/09-12, 10 E: 2010/07/15
Mode: med-res	Sampling: 0.03 Å/pix	10 A-C: 2010/06/05-07, 10 F: 2010/08/07

3. Some scientific results using spectra from the IACOB

There are already 2 published papers using data from the IACOB (and more in prep.). In Simón-Díaz (2010), we used the stellar atmosphere code FASTWIND (Puls *et al.* 2005) to perform a thorough self-consistent spectroscopic analysis of 13 early B-type stars from the various subgroups comprising the Orion OB1 association; this study showed that the dispersion of O and Si abundances between stars in the various subgroups found in previous analyses (e.g. Cunha & Lambert 1992) was a spurious result, being the consequence of a bad characterization of the abundance errors propagated from the uncertainties in the stellar parameter determination. In Simón-Díaz *et al.* (2010) we showed the first observational evidence for a correlation between macroturbulent broadening and line-profile variations in OB Supergiants using spectroscopic timeseries for a sample of 13 OB Sgs; this may support the hypothesis that macroturbulent broadening in this type of stars is likely a result of the collective effect of stellar pulsations. A subsample of IACOB spectra has been used within the *FLAMES-II Survey of Massive Stars: the Tarantula Survey* consortium to construct an atlas of medium resolution spectra of Galactic OB-type stars (Sana *et al.* in prep.). We plan to use this atlas for the spectral classification of the massive stars in 30 Dor. Finally, the scientific exploitation of the IACOB spectra concerning the quantitative spectroscopic analysis of the stars has already began and a series of papers with results will be published soon (see details in Simón-Díaz *et al.*, in prep.).

4. Future plans for the IACOB

In the next semesters, we will continue with the compilation of spectra for the IACOB, observing stars with $V \leq 8$ in at least three epochs (more in the case of known or newly detected binaries). Our idea is to make public the database via the Virtual Observatory in the next year. In the meantime, interested people can have access to the database under request to the author (ssimon@iac.es). The complete list of stars will be published in Simón-Díaz *et al.*, in prep.. We will acknowledge any observer who having obtained FIES spectra will like to add the spectra to the IACOB database after scientific exploitation.

References

- Barbá, R. H., Gamen, R., Arias, J. I., Morrell, N. *et al.* 2010, in: T. Rivinius & M. Curé (eds.), *The Interferometric View on Hot Stars*, Rev. Mexicana AyA Conference Series 38, p. 30
- Cunha, K. & Lambert, D. L. 1992, *ApJ*, 399, 586
- Evans, C., Hunter, I., Smartt, S., Lennon, D. *et al.* 2008, *The Messenger*, 131, 25
- Evans, C. J., Bastian, N., Beletsky, Y., Brott, I. *et al.* 2010, in: R. de Grijs & J. R. D. Lépine (eds.), *Star clusters: basic galactic building blocks throughout time and space*, IAU Symposium 266, p. 35
- Maíz-Apellániz, J., Walborn, N. R., Galué, H. Á. & Wei, L. H. 2004, *ApJS*, 151, 103
- Puls, J., Urbaneja, M. A., Venero, R., Repolust, T. *et al.* 2005, *A&A*, 435, 669
- Simón-Díaz, S. 2010, *A&A* 510, 22
- Simón-Díaz, S., Herrero, A., Uytterhoeven, K., Castro, N. *et al.* 2010, *ApJ (Letters)*, 720, L174



Speakers in action. From left to right and top to bottom: Alex Fulerton, Jose Groh, Sylvia Ekstrom, Thomas Rivinius, Stéphane Mathis, Hideyuki Saio, Maria-Fernanda Nieva, and Véronique Petit,