

As stated in the Preface, this book assumes some rudimentary knowledge of string theory, but it is a good idea to recall the basics. The field is notoriously vast and complex, so this chapter should not be understood as a replacement for serious study on one of the many great introductions [1–11]. In most of the book, we will approximate string theory by *supergravity*, an effective theory of gravitons and other fields; the presentation will be biased toward that.

In this chapter, we also assume knowledge of general relativity (GR) and some acquaintance with spinors, but we will try to keep mathematical sophistication at a minimum. We will develop some ideas, such as spinors and differential geometry, in much greater detail in the next few chapters before we return to physics. Still, already in this chapter we will pepper our presentation with occasional forward references to those mathematically more advanced treatments, to whet the reader's appetite.

1.1 Perturbative strings

A quantum field collects creation (and annihilation) operators for a representation of the Poincaré group. Once one fixes the value of the momentum p of the created state, the remaining degrees of freedom are a representation of the *little group*, or *stabilizer*, of p , namely the subgroup $\text{Stab}(p) \subset \text{SO}(d)$ of elements that leave p invariant. This is

$$\text{Stab}(p) = \text{SO}(d-1) \quad (p^2 < 0), \quad \text{Stab}(p) = \text{SO}(d-2) \ltimes \mathbb{R}^{d-2} \quad (p^2 = 0), \quad (1.1.1)$$

for the massive or massless case. (We will review the $p^2 = 0$ case in Section 3.3.6.) In the massless case, we would also have the possibility of selecting an infinite-dimensional representation, but this is usually regarded as exotic; so we select a finite-dimensional representation, ignoring the \mathbb{R}^{d-2} factor. Ordinary fields then represent objects with finitely many degrees of freedom, which we call spin and helicity for $m^2 > 0$ and $= 0$, respectively. Moreover, we usually take these objects to interact via terms of the type $\int d^d x \phi_1(x) \dots \phi_2(x)$: these allow the value of a field to influence directly that of another only at the same point.

All these reasons make us think of the quanta of a field as point particles. To describe a quantum theory of interacting extended objects, we need to change this picture somehow. First of all, a string can have infinitely many vibration modes, so a field that creates a string must be somehow a collection of infinitely many ordinary fields. Second, extended objects can interact when their centers of mass are not superimposed. So the interaction terms should be nonlocal.

Such a *string field theory* (SFT) is fascinating but also just as complicated as our description suggests. So in fact most studies of interacting strings focus on an approach that is *first-quantized*: one first decides the Feynman diagram one wants to consider, and then computes the amplitude associated with it. (A similar approach is used sometimes in quantum field theory too, under the name of *world-line formalism*.)

In this section, we will review quickly some aspects of this perturbative treatment of string interactions. There are five possible consistent string models:

- Type IIA
- Type IIB
- Heterotic with gauge group $E_8 \times E_8$
- Heterotic with gauge group $SO(32)$
- Type I

All these select $d = 10$ as spacetime dimension, in a sense we will clarify later in this chapter. The last case, type I, can be viewed as a certain quotient procedure from IIB strings, which we will introduce in [Section 1.4.4](#). So in this section we will discuss the other four. We will actually start our discussion from a model that has a *tachyon*, namely a scalar with a negative mass, but whose discussion is simpler: the *bosonic* string.

1.1.1 Bosonic strings

The action for a particle moving in a curved background is proportional to its “length in spacetime,” namely, to the proper time measured along its world-line (its trajectory γ in spacetime):

$$S_{\text{part}} = -m \int_{\gamma} d\sigma_0 \sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}, \quad (1.1.2)$$

where $x^\mu(\sigma_0)$ are the coordinates of the point in spacetime as a function of the world-line coordinate σ_0 , and $\dot{x}^\mu \equiv \partial_0 x^\mu$. In flat space, this is indeed minimized on straight lines in spacetime, which maximize proper time. For curved $g_{\mu\nu}$, (1.1.2) is minimized on geodesics. If we also have a Maxwell field and our particle is charged, we have to add a term

$$S_{\text{part,EM}} = q \int_{\gamma} d\sigma_0 A_\mu \partial_0 x^\mu, \quad (1.1.3)$$

where q is the charge, and A_μ is the vector potential. In [Section 4.1.4](#), we will see that the integrand is an example of a natural operation called *pull-back*.

String action

By analogy with (1.1.2), the natural action for a string would seem to be the volume of its two-dimensional world-sheet in spacetime. However, it is classically equivalent to the *Polyakov action*, which is easier to quantize:

$$S_{\text{F1,g}} = -\frac{1}{2} T_{\text{F1}} \int_{\Sigma} d^2\sigma h^{\alpha\beta} \sqrt{-h} g_{MN} \partial_\alpha x^M \partial_\beta x^N. \quad (1.1.4)$$

This type of action is also called a *sigma model*, for reasons going back to four-dimensional models of mesons, or sometimes *nonlinear sigma model* when g_{MN} is not flat. The $x^M(\sigma_0, \sigma_1)$, $M = 0, \dots, d-1$, describe the embedding of Σ in physical spacetime (often called *target space*), and h is a metric on Σ . The mass m in (1.1.2) has been replaced by the mass/length ratio, or *tension*:

$$T_{F1} = \frac{1}{2\pi l_s^2}. \quad (1.1.5)$$

“F” stands for *fundamental*, to distinguish this string from other extended objects that will appear later; 1 denotes the space extension of the string. The constant l_s is called *string length*. (We will always keep it explicit in this chapter, but later we will often work in string units and set $l_s = 1$.)

In this section, we are going to focus on strings that are *closed* or, in other words, that have no boundary. A generic¹ time slice is then a collection of several copies of the circle S^1 . The time evolution of each of these for a finite time will be a cylinder; then σ^1 is a periodic coordinate, $\sigma^1 \sim \sigma^1 + \pi$. These cylinders are then glued together at some values of σ^0 to obtain a general Σ .

Spectrum in flat space

Quantizing (1.1.4) is challenging for general g_{MN} but relatively easy in Minkowski space $g_{MN} = \eta_{MN}$: superficially (1.1.4) then becomes a collection of free bosons, with equations of motion $\partial^2 x^M = 0$. For a closed string, the slice at $\sigma^0 = \text{constant}$ is an S^1 ; there are then discrete Fourier modes for each x^M . Since the equation of motion is of second order, the states are in correspondence to the values of these Fourier modes and their derivatives. Alternatively, we can write a solution of the world-sheet equations of motion as $x^M = x^M(\sigma^+) + x^M(\sigma^-)$, where $\sigma^\pm = \sigma^1 \pm \sigma^0$, and introduce Fourier modes $\alpha_i^M, \tilde{\alpha}_i^M$ for the left- and right-movers $x^M(\sigma^\pm)$. The only subtlety is that the world-sheet metric $h_{\alpha\beta}$ is a Lagrange multiplier, which gives a constraint. This can be taken care of in many ways: by solving the constraint, or by introducing Faddeev–Popov ghosts and the Becchi–Rouet–Stora–Tyutin (BRST) method (the so-called covariant quantization). Skipping many interesting details, here we will just give the results.

Even for a fixed momentum, the spectrum has infinitely many states, of the form

$$\alpha_{-i_1}^{N_1} \dots \alpha_{-i_n}^{N_n} \tilde{\alpha}_{-j_1}^{\tilde{N}_1} \dots \tilde{\alpha}_{-j_n}^{\tilde{N}_n} |0\rangle, \quad (1.1.6)$$

where $|0\rangle$ is the world-sheet vacuum, and $i_k, j_k \geq 0$ (possibly repeated). As we mentioned, these correspond to the vibration modes of the string, and in a spacetime picture they would require infinitely many ordinary quantum fields to create them. Their masses are

$$m^2 = \frac{4}{l_s^2} \left(\frac{2-d}{24} + N \right), \quad (1.1.7)$$

where $N = \sum i_k = \sum j_k$ is a nonnegative integer. The identity between these two expressions is called *level matching* and is the link between the left- and right-moving sectors, which otherwise proceed on parallel tracks. If $d > 2$, we see that the lowest

¹ The mathematical meaning of the word *generically*, which we will use in this book, is “for any choice except for a set of measure zero.”

value of m^2 , for $N = 0$, is actually *negative*. Such a mode is usually called a *tachyon* and signals an instability. For this reason, the bosonic string we are discussing in this section is usually only considered a toy model.

Nevertheless, it already displays a very interesting feature. For the *critical dimension* $d = 26$, the modes with $n = 1$ in (1.1.6) and (1.1.7) are massless. They read

$$\alpha_{-1}^M \tilde{\alpha}_{-1}^N |0\rangle, \tag{1.1.8}$$

and so they correspond to fields with two indices. Among these we thus find a *massless spin-two field* $h_{MN} = \delta_{gMN}$. The action (1.1.4) can then be thought of as a string moving in a condensate of such a field. This is a bit similar to expanding a quantum field theory (QFT) around a vacuum where a field has acquired a nonzero expectation value.

So we have found that in $d = 26$, the string modes include those that would normally be associated with a graviton. Remarkably, the scattering amplitudes one obtains with this formalism are *finite*. The string tension acts as a regulator: Taking the limit $l_s \rightarrow 0$, the scattering amplitudes become divergent again. In this limit, the theory becomes a local QFT model again, and a local theory of gravity has divergent amplitudes.

Coupling to condensates of other fields

Among (1.1.8), we find other massless modes. Following (1.1.1), we need to consider only the components of (1.1.8) in the $d - 2 = 24$ dimensions transverse to the momentum p , which are 24^2 . The physical components h_{MN} of a graviton are represented by a traceless 24×24 matrix; this is the generalization of the transverse traceless (TT) gauge familiar from the treatment of gravitational waves in four dimensions. The remaining modes are thus the antisymmetric part of (1.1.8) and its trace. The fields that create these states are an antisymmetric *Kalb–Ramond* field $B_{MN} = -B_{NM}$, and a scalar field ϕ called *dilaton*. So in total the massless fields of the bosonic string are

$$g_{MN}, \quad B_{MN}, \quad \phi. \tag{1.1.9}$$

We can consider condensates of B_{MN} and ϕ , too; this leads to the extra terms in the action:

$$S_{F1,B,\phi} = -\frac{1}{2} T_{F1} \int_{\Sigma} d^2\sigma \left[\epsilon^{\alpha\beta} B_{MN} \partial_{\alpha} x^M \partial_{\beta} x^N + l_s^2 \sqrt{-h} R_{(2)} \phi \right]. \tag{1.1.10}$$

Here $R_{(2)}$ is the scalar curvature of the world-sheet metric $h_{\alpha\beta}$, and $\epsilon = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. The coupling with B is the natural generalization of the coupling (1.1.3). The coupling with the dilaton is peculiar in that

$$\frac{1}{4\pi} \int_{\Sigma} \sqrt{-h} R_{(2)} = 2 - 2g, \tag{1.1.11}$$

where g is the *genus* of the world-sheet Σ . This is the stringy analogue of the number of loops, and can be intuitively described (when Σ has no boundary) as the *number*

of handles; a more formal definition will be given in Section 4.1.10. Because of this, the computation of all scattering amplitudes is organized in powers:

$$g_s^{2g-2}, \quad g_s \equiv e^\phi. \quad (1.1.12)$$

We can think of g_s as a *string coupling constant*: when it is small, the powers (1.1.12) are smaller for Riemann surfaces Σ of increasing g , which can be thought of as the stringy analogue of Feynman diagrams of increasing complexity.

The action

$$S_{\text{bos}} = S_{\text{F1},g} + S_{\text{F1},B,\phi} \quad (1.1.13)$$

is classically invariant under general coordinate transformation $\sigma_\alpha \rightarrow \sigma'_\alpha(\sigma_0, \sigma_1)$, if we also take care to transform the world-sheet metric $h_{\alpha\beta}$. This is a gauge invariance, in that it doesn't affect the physical configuration, the image of the world-sheet embedding $x^\mu(\sigma)$, but only how we parameterize it. Equation (1.1.13) is also invariant under Weyl rescaling $h_{\alpha\beta} \rightarrow e^f h_{\alpha\beta}$. In two dimensions, one can fix the coordinate-change freedom by taking, for example, $h_{\alpha\beta}$ to have constant scalar curvature. Even so, a residual invariance remains: coordinate transformations that leave the metric invariant up to a Weyl transformation. These are called *conformal transformations*.

Conformal invariance and effective action

It is crucial that this residual gauge invariance remains at the quantum level. It decouples potentially harmful negative-norm states that would come from the fact that x^0 in (1.1.4) has a wrong-sign kinetic term. This is similar to what happens in the quantization of the electromagnetic field, for example. Conformal invariance is also behind the absence of high-energy divergences. Usually scattering amplitudes become problematic when two particles collide at a small impact parameter. The world-sheet of a string scattering is a non-compact Riemann surface with several spikes s_i corresponding to the incoming and outgoing strings. Conformal invariance means that the distance between two points on the world-sheet has no intrinsic meaning: only ratios of distances do. So a small impact parameter might seem to correspond to two such spikes s_1 and s_2 getting close, but that only means that they are close relative to their distance from other external strings s_i . This corresponds to a Riemann surface that develops a long neck, where the two s_i are both attached, far from the others.

The Noether current associated to dilatations in a field theory is $T^{\mu\nu} x_\nu$, where $T_{\mu\nu}$ is the stress–energy tensor. This is conserved if $0 = \partial_\mu (T^{\mu\nu} x_\nu) = T^{\mu\nu} g_{\mu\nu} = T^\mu_\mu$. Evaluating the expectation value $\langle T^\mu_\mu \rangle$ of this trace is thus a way to check if there is a *Weyl anomaly*.

From the point of view of the world-sheet, the spacetime fields (1.1.9) are really couplings for the action of the fields $x^M(\sigma)$. So a Weyl anomaly can also be detected by computing the beta functions of the action (1.1.13) for the couplings (1.1.9). This can be obtained by the usual perturbative methods; the coupling for this computation is given by l_s^2 , or rather the dimensionless combination $l_s^2 \times$ (spacetime curvature). This results in the following three conditions:

$$R_{MN} + 2\nabla_M \partial_N \phi - \frac{1}{4} H_{MPQ} H_N{}^{PQ} + O(l_s^2) = 0, \quad (1.1.14a)$$

$$\nabla_M (e^{-2\phi} H^M{}_{NP}) + O(l_s^2) = 0, \quad (1.1.14b)$$

$$\frac{2}{3l_s^2} (26 - d) + R - \frac{1}{2} |H|^2 - 4e^\phi \nabla^2 e^{-\phi} + O(l_s^2) = 0. \quad (1.1.14c)$$

We have introduced

$$H_{MNP} = \partial_M B_{NP} + \partial_N B_{PM} + \partial_P B_{MN}, \quad |H|^2 \equiv \frac{1}{6} H_{MNP} H^{MNP}. \quad (1.1.15)$$

This can be considered as a *field-strength* for the *potential* B_{MN} , similar to the relation between $F_{MN} = \partial_M A_N - \partial_N A_M$ and A_M in electromagnetism. Indeed, there is also a gauge transformation

$$B_{MN} \rightarrow B_{MN} + \partial_M \hat{\lambda}_N - \partial_N \hat{\lambda}_M, \quad (1.1.16)$$

under which (1.1.15) is invariant. The world-sheet action (1.1.10) is invariant too under this, because the transformation adds a total derivative term.

From spacetime point of view, where (1.1.9) are fields, (1.1.14) are to be interpreted as equations of motion. They can be obtained by extremizing²

$$S_{\text{bos}} = \frac{1}{2\kappa_b^2} \int d^d x \sqrt{-g} e^{-2\phi} \left(\frac{2}{3l_s^2} (26 - d) + R + 4\partial_M \phi \partial^M \phi - \frac{1}{2} |H|^2 + O(l_s^2) \right) \quad (1.1.17)$$

with respect to (1.1.9). By dimensional reasons, κ_b has dimension l_s^{12} . (The metric coefficients have no dimension, while R contains two derivatives and has mass dimension two.) In general, the *Planck mass* m_P is defined as the mass scale entering the Einstein–Hilbert action; the *Planck length* l_P is its inverse, and (1.1.17) tells us that it is proportional to l_s .

As a consistency check, we see that flat space is a solution of (1.1.14) only if we set $d = 26$, which is the value where we found the massless fields (1.1.9) in the first place. More generally, to trust (1.1.14) we have to make sure that the expansion parameter $l_s^2 \times$ (curvature) is small, so we better solve those equations of motion separately at every order. This leads again to taking

$$d = 26. \quad (1.1.18)$$

It is conceptually possible to consider solutions where $d \neq 26$, and the first term in (1.1.14c) competes with the others, but in that case we have to worry that the other terms in the l_s expansion become relevant too, and we have not given them in (1.1.14). If, on the other hand, one is able to prove that a certain world-sheet model is conformal exactly, without using the l_s expansion at all, then $d = 26$ is not necessary. There are not many such cases: one is the *linear dilaton* background, where ϕ is linear in one of the coordinates. This leads to *noncritical string theories*, which historically have been important toy models.

Another point of view on the critical dimension is this. We observed that (1.1.13) has conformal invariance. Conformal transformations form a group; for flat space it is $\text{SO}(d - 2, 2)$ for $d > 2$, but for $d = 2$ it becomes infinite dimensional. Indeed,

² This variation is a little more involved than the usual Einstein–Hilbert action variation because of the prefactor $e^{-2\phi}$. More details will be given in Section 10.1.2.

any transformation $x^\pm \rightarrow x^{\pm'}(x^\pm)$ is conformal for any metric of the type $ds^2 = e^f dx^+ dx^-$. The generators L_m , $m \in \mathbb{Z}$, of such transformations on the x^+ obey the Lie algebra

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}m(m^2 - 1)\delta_{n+m,0}, \quad (1.1.19)$$

called *Virasoro algebra*. The $L_0, L_{\pm 1}$ form an $SO(1,2)$ subalgebra where c does not appear. As usual, spacetime transformations are generated by the stress–energy tensor, so these L_m are related to it. After a Wick rotation, $x_+ \rightarrow z = \sigma^1 + i\sigma^0$, and we can collect all the generators in

$$T_{zz}(z) = \sum_n L_n z^{-n-2}. \quad (1.1.20)$$

In a Lie algebra, the commutation relations should always be linear, so we need to think of the second term in (1.1.19) as containing a new generator c , which commutes with all the others, and thus lies in the center of the algebra; so c in (1.1.19) is called *central charge*. The \tilde{L}_m on the x_- variable generate a second copy of the same algebra (1.1.19), and they are collected in $T_{\bar{z}\bar{z}}$.

This c is also a measure of the Weyl anomaly: for any QFT model that is conformal on a flat (world-sheet) metric $h_{\alpha\beta} = \eta_{\alpha\beta}$, a nonzero c tells us that conformal invariance is broken for more general $h_{\alpha\beta} \neq \eta_{\alpha\beta}$. A free boson contributes $c = 1$, while the ghosts give -26 . Thus if we quantize *around flat space*, where the $x^M(\sigma)$ bosons are free, for quantum conformal invariance we need to take $d = 26$.

The fact that the action (1.1.17) exists at all is nontrivial from the point of view of the world-sheet derivation we described. We can think of it as being an approximation to the string field theory action S_{SFT} , which would also contain the massive fields creating all the states (1.1.6). We can call it an effective action, in the usual quantum field theory sense: It reproduces the results one would obtain from S_{SFT} , at energies that are *low*, namely much smaller than l_s^{-1} . Indeed, another way to compute (1.1.17) is to compute string scattering amplitudes using the world-sheet approach, and then guessing what spacetime action would reproduce them.

The diagrams leading to (1.1.17) have $g = 0$ in (1.1.12), leading to $g_s^{-2} = e^{-2\phi}$, thus explaining the presence of that exponential. The higher powers of l_s hidden in (1.1.17) also receive contributions from higher values of g (and thus from more complicated Feynman diagrams). So the effective action will have a double expansion in powers of both:

$$S = \sum_{j,k} S_{j,k} l_s^j e^{k\phi}. \quad (1.1.21)$$

These higher-order corrections can in principle be computed; we will see some examples for superstrings. When we first discover that GR is non-renormalizable and needs (curvature)² counterterms [25], we might perhaps hope that by adding more and more such counterterms, with arbitrary powers (curvature)^k, we might eventually find a theory that has no divergence. Finding such a renormalizable theory of gravity would be very hard without some sort of guidance: not only would we have to find a fixed point of the renormalization group (RG) flow by going backward in energy, but we would also have to worry about modes with wrong kinetic energy, which in such theories generically abound. (Adding operators with higher numbers

of derivatives to a Lagrangian also adds propagating modes, each of which might be a ghost.) String theory is renormalizable, and in principle we can reexpress it precisely as such a sum of infinitely many corrections to (1.1.17).

This discussion seems, however, to assume that the effective action is analytic in the parameters l_s, e^ϕ , or, in other words, that it coincides with its Taylor expansion (1.1.21). In mathematics, we know many functions that are not analytic, and they might also appear here. This is the reason we have put the word “perturbative” in the title of this section; we will make amends in Section 1.4.

Some critics of string theory complain that the theory has not been proven to be *background independent*. What they mean is that in the world-sheet approach based on (1.1.13), we first have to fix a background configuration for the spacetime fields (1.1.9), and then we can compute an action for the small fluctuations around it. A priori, it might even be unclear if this procedure is describing a single theory or a collection of theories that have nothing to do with each other. The emergence of (1.1.17) should be reassuring in this respect: that effective action can be expanded around any background, and matches the result of the world-sheet method around it. A more satisfactory rebuttal is the proof at the level of string field theory in [26].

Torus compactification

Finally, let us have a first taste of string compactifications, by supposing that the theory lives on $\mathbb{R}^{25} \times S^1$. Thus we declare one of the coordinates to be periodically identified, say $x^{25} \equiv x^{25} + 2\pi R$. Now $x^{25}(\sigma^0, \sigma^1)$ is no longer necessarily periodic as a function of σ^1 , even for a closed string: rather, if we take $\sigma^1 \sim \sigma^1 + \pi$, we demand

$$x^{25}(\sigma^0, \sigma^1 + \pi) = x^{25}(\sigma^0, \sigma^1) + 2\pi wR. \tag{1.1.22}$$

This represents a string that winds $w \in \mathbb{Z}$ times around the S^1 . Another new effect is familiar from quantum mechanics: the overall momentum of the string in the S^1 direction is now not continuous but quantized: $p^{25} = \frac{q}{R}, q \in \mathbb{Z}$.

The mass spectrum in \mathbb{R}^{25} is now modified from (1.1.7) to

$$m^2 = \frac{4}{l_s^2}(-1 + N) + \left(\frac{q}{R} - w\frac{R}{l_s^2}\right)^2 = \frac{4}{l_s^2}(-1 + \tilde{N}) + \left(\frac{q}{R} + w\frac{R}{l_s^2}\right)^2, \tag{1.1.23}$$

where now $N = \sum i_k$ and $\tilde{N} = \sum_k \tilde{i}_k$ are no longer necessarily equal (as they were in (1.1.7)); comparing the two expressions, we have $N - \tilde{N} = wq$.

For a generic value of R , the massless spectrum is still (1.1.8) and (1.1.9); but now it should be reinterpreted. The components

$$g_{M25}, \quad B_{M25} \tag{1.1.24}$$

are now two vector fields in \mathbb{R}^{25} ; g_{2525} is a scalar. The remaining components of (1.1.9) then give a metric, a Kalb–Ramond field, and a scalar in \mathbb{R}^{25} .

From (1.1.23), however, we also see another option: if $\frac{q}{R} - w\frac{R}{l_s^2} = \pm\frac{2}{l_s}$, then we have a new massless state for $N = 0$. This is possible for

$$R = l_s, \tag{1.1.25}$$

taking $q = -w = \pm 1$; then $\tilde{N} = \sum_k \tilde{i}_k = 1$. This state $\tilde{\alpha}_{-1}^M|0\rangle$ has a single index, and so it is created by a vector field. At this value of R , we also have the possibility of using the same trick with the other expression in (1.1.23), this time leading to

$q = w = \pm 1, \tilde{N} = 0, N = 1$. So we have a total of four more vector fields in \mathbb{R}^{25} . It turns out that these combine with the previous two (1.1.24) to give a nonabelian gauge group

$$SU(2) \times SU(2). \tag{1.1.26}$$

This compactification was rather nice in that the string could be quantized exactly, at least perturbatively in g_s . In more complicated cases, we won't be so lucky, and we will have to limit ourselves to the less powerful effective field theory methods, potentially missing phenomena such as this non-abelian gauge group enhancement.

1.1.2 Type II superstrings

Supersymmetric world-sheet action

The world-sheet action (1.1.13) can be made supersymmetric. At the most basic level, this means that we promote the $x^M(\sigma)$ to a function of σ and of new formal coordinates θ^\pm that anticommute: $\theta^+\theta^- = -\theta^-\theta^+, (\theta^\pm)^2 = 0$. The Taylor expansion in the new coordinates truncates:

$$X^M = x^M + \theta^+\psi_+^M + \theta^-\psi_-^M + \theta^+\theta^-F^M. \tag{1.1.27}$$

We can also introduce the derivative operators

$$D_\pm = \partial_{\theta^\pm} + i\theta^\pm\partial_\pm, \quad \partial_\pm \equiv \partial_{\sigma^\pm}. \tag{1.1.28}$$

Then, (1.1.4), for example, is replaced by

$$S_{F1,g}^{1,1} = -\frac{1}{2}T_{F1} \int_\Sigma d^2\sigma d^2\theta (g + B)_{MN}(X) D_+X^M D_-X^N, \tag{1.1.29}$$

with the integration rule $\int d\theta^\pm\theta^\pm = 1, \int d\theta^\pm 1 = \int d\theta^\pm\theta^\mp = 0$. We also added the contribution from B . The terms (1.1.10) can also be supersymmetrized in this way. The final result is quite messy for a general background where g_{MN} and B_{MN} are arbitrary; it can be found, for example, in [27, sec. 6.3.1]. For example, it contains a kinetic term

$$g_{MN}(\psi_+^M\partial_-\psi_+^N + \psi_-^M\partial_+\psi_-^N) \tag{1.1.30}$$

for the world-sheet fermions ψ_\pm^M .³ The F^M in (1.1.27) are auxiliary fields: they have no kinetic term, and can be replaced with the solutions of their equations of motion.

Since we have introduced a single θ^+ and a single θ^- , the resulting model is said to have $\mathcal{N} = (1, 1)$ supersymmetry. Any two-dimensional bosonic model can be promoted to such a model. In the context of compactifications, one often needs to separate external and internal dimensions, and the supersymmetrization of the world-sheet model in the latter has more supercharges; a common case one needs is $\mathcal{N} = (2, 2)$. This is more challenging to achieve, because such *extended* supersymmetry requires that one combine the x^M with each other in pairs. Such a pairing is reminiscent of the idea of complex coordinates, and is at the root of why differential geometry is useful for compactifications. This idea will return in Chapter 9.

³ The bosonic world-sheet indices \pm are conceptually not the same as the \pm on the fermions, denoting chirality. To emphasize the difference, some authors change the world-sheet indices to $+$ \rightarrow $\#$ and $-$ \rightarrow \ominus . This also has the benefit that every term in a world-sheet will then have an equal number of pluses and minuses; see for example [28].

Equation (1.1.29) is called the *Neveu–Schwarz–Ramond* (NSR) model. While we introduced it by supersymmetrizing the world-sheet action, we will see later that the resulting spacetime theory also has the much more nontrivial property of *spacetime* supersymmetry.

Spectrum

Even around flat space, the spectrum of (1.1.29) is now more complicated because it depends on what we impose on the fermionic ψ_{\pm}^M . Since a fermion should only get back to itself after a 4π rotation, under 2π we can impose either periodic or antiperiodic boundary conditions, called Neveu–Schwarz (NS) and Ramond (R) respectively. These can be imposed independently on the ψ_{\pm}^M , leading to four sectors: NSNS, NSR, RNS, and RR. The spectrum has to be analyzed in each sector separately, because the Fourier modes for the ψ_{\pm}^M behave differently in each.

In the NS sector, the fermionic Fourier modes are $b_{-i-1/2}^M$, $i \geq 0$. The two lowest-lying states are

$$|0\rangle_{\text{NS}}, \quad \mathbf{1}, \quad m^2 = \frac{1}{8l_s^2}(2-d); \quad (1.1.31a)$$

$$b_{-1/2}^M|0\rangle_{\text{NS}}, \quad \mathbf{8}_V, \quad m^2 = \frac{1}{l_s^2}\left(\frac{(2-d)}{8} + 1\right). \quad (1.1.31b)$$

We have also indicated what representation these states form under the compact part $\text{SO}(d-2) = \text{SO}(8)$ of the massless little group (1.1.1). For (1.1.31b), the subscript “V” is because there are two more dimension-eight representations of $\text{SO}(8)$, which will soon play a role too.

In the R sector, the fermionic Fourier modes are d_{-i}^M , $i \geq 0$. In this case, the vacuum has already $m^2 = 0$, but in fact it is not unique: the modes d_0^M now don’t raise the energy, and they act on the space of vacua. These d_0^M satisfy a *Clifford algebra* $\{d_0^M, d_0^N\} = 2g^{MN}1$, and as a consequence the space of R vacua transforms as a spinor under spacetime symmetries. In Section 2.1, we will attack Clifford algebras and spinors systematically in every dimension; for now, we only state the main features we need, which are quite similar to the properties of gamma matrices in four dimensions.

- Gamma matrices Γ^M can be defined in every dimension as matrices that satisfy $\{\Gamma^M, \Gamma^N\} = 2g^{MN}1$.
- In $d = 10$ dimensions, they are 32×32 matrices; in $d = 8$, they are 16×16 .
- The space of spinors on which the Γ^M act is a representation for the Lorentz group; in d even, it decomposes in two chiralities, for which we introduce indices $\alpha, \dot{\alpha}$. Multiplication by a single Γ^M changes chirality, so the nonzero blocks are $\Gamma_{\alpha\dot{\beta}}^M$ and $\Gamma_{\dot{\alpha}\beta}^M$.
- In both $d = 10$ with Lorentzian signature and $d = 8$ with Euclidean signature, there is a choice of Γ^M that are all real. (This aspect will be treated more specifically in Sections 2.2.3 and 2.3.)

As a representation of the transverse $\text{SO}(8)$ in the little group (1.1.1), the R states then form a reducible representation of dimension 16, which further splits in two

representation of dimension eight, traditionally called $\mathbf{8}_S$ and $\mathbf{8}_C$. We summarize all this by writing

$$|0, \alpha\rangle_R, \quad \mathbf{8}_S, \quad m^2 = 0; \tag{1.1.32a}$$

$$|0, \dot{\alpha}\rangle_R, \quad \mathbf{8}_C, \quad m^2 = 0. \tag{1.1.32b}$$

The closed-string spectrum is obtained by taking tensor products of these two sectors. For example, if we impose NS conditions for both ψ_{\pm}^M , we have the NSNS sector, whose low-lying states are obtained by taking tensor products of two copies of (1.1.31). Here we see once again the presence of a tachyon, $|0\rangle_{NS} \otimes |0\rangle_{NS}$. In the critical dimension

$$d = 10, \tag{1.1.33}$$

this sector contains massless states $b_{-1/2}^M \tilde{b}_{-1/2}^M |0\rangle_{NS} \otimes |0\rangle_{NS}$, which would be created by the fields (1.1.9) we encountered for the bosonic string, so we are going to stick to $d = 10$ from now on.

Quite nontrivially, just like in the bosonic string, this is also the dimension where the *Weyl anomaly vanishes*. This can be checked in any of the methods we saw for the bosonic string. In terms of the central charge of the Virasoro algebra, for example, there are now ghosts and superghosts, which give a total contribution -15 . Quantizing around flat space, each boson gives a contribution $+1$, but now each fermion gives an extra $+1/2$. Since (1.1.29) has an equal number of bosons of fermions, we get $(1 + 1/2)d = 15$, which leads again to $d = 10$.

A less welcome similarity with the bosonic string is the presence of a tachyon. Fortunately, this can be eliminated by the so-called *Gliozzi–Scherk–Olive (GSO) projection* on the spectrum. Initially an ad hoc prescription, it later emerged to be required by consistency when one takes the world-sheet Σ to be a torus, an invariance under coordinate changes called *modular invariance*. It projects out $|0\rangle_{NS}$ in the NS sector, and also eliminates one of the two sets in (1.1.32). For the left-moving ψ_+^M , it is immaterial which one we choose; we keep $\mathbf{8}_S$. However, now the choice for the right-moving ψ_-^M does matter; this leads to two different theories.

If we keep $\mathbf{8}_C$ for the ψ_-^M , we obtain the theory called *IIA string theory*; its spectrum is given in Table 1.1. We use $SO(8)$ group theory to decompose the tensor products of representations as direct sums. In the last column, we have named the corresponding spacetime fields.

- In the NSNS sector, the decomposition is that of an 8×8 matrix in its symmetric traceless, antisymmetric and trace, with the same logic that led us to the fields (1.1.9) for the bosonic string.

NSNS	$\mathbf{8}_V \otimes \mathbf{8}_V = \mathbf{35}_V \oplus \mathbf{28}_V \oplus \mathbf{1}_V$	g_{MN}, B_{MN}, ϕ
RNS	$\mathbf{8}_S \otimes \mathbf{8}_V = \mathbf{56}_S \oplus \mathbf{8}_C$	$\psi_{M\dot{\alpha}}^1, \lambda_{\dot{\alpha}}^1$
NSR	$\mathbf{8}_V \otimes \mathbf{8}_C = \mathbf{56}_C \oplus \mathbf{8}_S$	$\psi_{M\dot{\alpha}}^2, \lambda_{\dot{\alpha}}^2$
RR	$\mathbf{8}_S \otimes \mathbf{8}_C = \mathbf{56}_V \oplus \mathbf{8}_V$	C_{MNP}, C_M

- In the RNS sector, the $\mathbf{8}_S \otimes \mathbf{8}_V$ representation has a vector and a spinor index, leading to an object $\Psi_{M\alpha}$. This is not irreducible: we can use an eight-dimensional gamma matrix to extract a sort of trace $\lambda_{\dot{\alpha}} \equiv (\Gamma^M)_{\alpha}^{\dot{\beta}} \Psi_{M\dot{\beta}}$, transforming as $\mathbf{8}_C$; the remaining traceless part of $\Psi_{M\alpha}$ is the $\mathbf{56}_S$. Both have one spinor index, and thus are spacetime fermions. Since they only have an index of one kind (either α or $\dot{\alpha}$), they are chiral, or *Weyl*. In the aforementioned real basis for the Γ^M , these fermions are also all real. The basis-independent notion is called the *Majorana* property (Section 2.3.1).
- The NSR sector is similar, but with $\mathbf{8}_S \leftrightarrow \mathbf{8}_C$. These are also spacetime fermions. The $\psi_{\alpha M}^1, \psi_{\dot{\alpha} M}^2$ are related by supersymmetry to the metric field, and hence are called *gravitinos*. The $\lambda_{\dot{\alpha}}^1, \psi_{\alpha}^2$ are called *dilatinos*.
- In the RR sector, we have an object with two spinorial indices:

$$C_{\alpha\dot{\beta}}. \tag{1.1.34}$$

These have two spinor indices, and thus are sometimes called *bispinors*; they are spacetime *bosons*.

A more familiar description for the RR (1.1.34) is obtained by expanding them in a basis for bispinors. This is familiar from QFT in $d = 4$: it consists of antisymmetrized products

$$\Gamma^{MN} \equiv \frac{1}{2}(\Gamma^M \Gamma^N - \Gamma^N \Gamma^M), \quad \Gamma^{MNP} = \frac{1}{6}(\Gamma^M \Gamma^N \Gamma^P \pm \text{perm.}), \tag{1.1.35}$$

and so on. The expansion on this basis is called a *Fierz identity* and will be analyzed systematically in Section 3.4; for now, we sketch the result. If we contract (1.1.34) with a single eight-dimensional gamma matrix, we obtain the vector

$$C_M = (\Gamma_M)^{\alpha\dot{\beta}} C_{\alpha\dot{\beta}}, \tag{1.1.36}$$

which is the $\mathbf{8}_V$ there. One can further contract with products of gamma matrices; with two of them, we obtain $\Gamma_{\alpha\dot{\beta}}^{MN}$ or $\Gamma_{\dot{\alpha}\beta}^{MN}$, which cannot be contracted with (1.1.34), but with three we do obtain $C_{MNP} = (\Gamma_{MNP})^{\alpha\dot{\beta}} C_{\alpha\dot{\beta}}$. This explains the entries in the bottom-right corner of Table 1.1. By construction, this is completely antisymmetric:

$$C_{MNP} = -C_{NMP} = -C_{PNM} = -C_{MPN}. \tag{1.1.37}$$

The number of independent components of a completely antisymmetric tensor with k indices in d (transverse) dimensions is $\binom{d}{k}$; for (1.1.37) this gives $\binom{8}{3} = 56$, confirming the last row of Table 1.1. One might think that we also need to consider $\Gamma^{M_1 \dots M_k}$ with higher odd k , but these are actually related to the ones already present in Table 1.1, as shown by the matching dimension count in the representation. Sometimes it is convenient to work with a redundant set of RR potentials; this *democratic* formalism will be presented in Section 10.1.

The second choice is to keep $\mathbf{8}_S$ for the ψ_-^M . The resulting theory is now called *IIB string*, and its spectrum is shown in Table 1.2. Most differences with Table 1.1 are straightforward; the one worthy of most attention is that now the RR sector consists of a bispinor

$$C_{\alpha\beta}. \tag{1.1.38}$$

Table 1.2. Massless IIB spectrum.

NSNS	$\mathbf{8}_V \otimes \mathbf{8}_V = \mathbf{35}_V \oplus \mathbf{28}_V \oplus \mathbf{1}_V$	g_{MN}, B_{MN}, ϕ
RNS	$\mathbf{8}_S \otimes \mathbf{8}_V = \mathbf{56}_S \oplus \mathbf{8}_C$	$\psi_{M\alpha}^1, \lambda_{\dot{\alpha}}^1$
NSR	$\mathbf{8}_V \otimes \mathbf{8}_S = \mathbf{56}_S \oplus \mathbf{8}_C$	$\psi_{M\alpha}^2, \lambda_{\dot{\alpha}}^2$
RR	$\mathbf{8}_S \otimes \mathbf{8}_S = \mathbf{35}_C \oplus \mathbf{28}_C \oplus \mathbf{1}_C$	C_{MNPQ}^+, C_{MN}, C_0

In the Fierz expansion, now the Γ^{MN} has the correct index structure, leading to $C_{MN} = (\Gamma_{MN})^{\alpha\beta} C_{\alpha\beta}$; a similar projection can be defined for all the products $\Gamma^{M_1 \dots M_k}$ with even k , but in fact $k > 6$ are redundant, and even the one for $k = 4$ has a “self-duality property” that halves its degrees of freedom, and whose consequences we will see soon. This is the reason of the superscript + on C_{MNPQ}^+ . Just like for (1.1.37), by construction these tensors are all completely antisymmetric.

Spacetime supersymmetry

In both IIA and IIB, the spacetime bosons arise from the NSNS and RR sectors, while the fermions arise from the NSR and RNS sectors. They have the same total number (128) of degrees of freedom. This is a symptom of the aforementioned *spacetime* supersymmetry. These mix the bosonic fields (NSNS, RR) with the fermionic ones (NSR, RNS). They are rather complicated, and at this stage their expression would not look very informative. We will have a first look at them in Section 8.2 for the case where only the metric is present, and in Section 10.1 in full. For now, we just comment about their infinitesimal parameters, which are *two* Majorana–Weyl spinors:

$$\epsilon^1, \quad \epsilon^2. \quad (1.1.39)$$

(This is the original reason these theories are called “type II.”) They have the same chirality as the gravitino: thus in IIA $\epsilon_{\dot{\alpha}}^1$ has positive chirality, and $\epsilon_{\dot{\alpha}}^2$ has negative chirality, while in IIB both $\epsilon_{\dot{\alpha}}^a$ have positive chirality. Altogether this gives 32 supercharges, which is the highest number for any supersymmetric theory.

This property is made manifest in the alternative *Green–Schwarz* model. A more recent formulation is the *Berkovits*, or *pure spinor*, model [29, 30]. (Pure spinors will play an important role in this book, but for somewhat different reasons.) These alternative formulations are more complicated, but are better at describing strings in condensates of RR fields, while with the NSR model (1.1.29) this is difficult.

1.1.3 Heterotic strings

Definition

In both the bosonic string and the superstring, quantization of the left- and right-movers seems to proceed almost independently. This is because they are almost free as two-dimensional field theories, apart from the constraint associated to the Lagrange multiplier $h_{\alpha\beta}$. For example, in the bosonic string spectrum (1.1.6), the constraint is only visible in the level matching condition mentioned after (1.1.7).

This creates an opportunity: we can try to define a hybrid, or *heterotic* theory which looks like the bosonic string for the left-movers, and like the superstring for the

right-movers [31]. This might look impossible: is the spacetime dimension going to be 26 or 10? The dimension of spacetime, however, is a macroscopic concept, irrelevant at length scales below l_p , where quantum gravity sets in. So a solution to the puzzling mismatch in dimension is to compactify 16 of the 26 bosonic coordinates. Provided we manage to satisfy the level matching condition relating the left- and right-moving sectors, all should be well.

The easiest way to compactify the 16 left-moving bosons x_+^I is to take them to belong to a *torus*. In Section 1.1.1, we saw a circle compactification of the bosonic string: we took one of the coordinates $x^{25} \sim x^{25} + 2\pi R$. The simplest torus would be obtained as $T^{16} \equiv (S^1)^{16}$, with the same identification for all the x_+^I . A straightforward generalization is to take a basis $\{R_a^I\}$, $a = 1, \dots, 16$ of \mathbb{R}^{16} , and to introduce equivalence relations

$$x^I = x^I + \pi R_a^I \quad \forall a. \tag{1.1.40}$$

The set of integer multiples of the $\{R_a^I\}$ is called a *lattice* Γ . The space defined by (1.1.40) is topologically still T^{16} ; the choice of Γ affects its size and shape. The momenta are quantized according to

$$p^I R_a^I \in \mathbb{Z}. \tag{1.1.41}$$

The solutions to this equation are the elements of a *colattice*: there is a dual set of vectors p_a^I of which all solutions to (1.1.41) are integer multiples.

Spectrum

As usual, now the spectrum is computed independently for the left-movers and for the right-movers, imposing level matching. For the left-movers, $(\partial_0 - \partial_1)x^I = 0$ relates p^I to the R_a^I : this generalizes setting $q/R = wR/l_s^2$ in (1.1.23). So in fact the lattice and colattice coincide,⁴ and Γ is said to be *self-dual*. Moreover, the analogue of (1.1.23) now reads

$$m^2 = \frac{4}{l_s^2} \left(-1 + N + \frac{1}{2} \hat{p}^2 \right) = \frac{4}{l_s^2} \tilde{N}. \tag{1.1.42}$$

From this we see that $\frac{1}{2} \hat{p}^2 \in \mathbb{Z}$: the lattice is said to be *even*.

The massless spectrum is again obtained as a tensor product of left- and right-movers. The latter have both an NS and an R sector, just as in the superstring (recall (1.1.31) and (1.1.32)). For the left-movers, (1.1.42) gives us two possibilities. We can take $\hat{p}^2 = 0$ and $N = 1$: a single Fourier mode acting on the vacuum (compare with the left-moving part of (1.1.8)). This can be α_{-1}^M , with $M = 0, \dots, 9$, or α_{-1}^I , with $I = 1, \dots, 16$ one of the extra 16 bosonic directions. So far, this gives us the massless states

$$\alpha_{-1}^M |0\rangle_{\text{bos}} \otimes b_{-1/2}^M |0\rangle_{\text{NS}}, \quad \alpha_{-1}^M |0\rangle_{\text{bos}} \otimes |0, \alpha\rangle_{\text{R}}; \tag{1.1.43a}$$

$$\alpha_{-1}^I |0\rangle_{\text{bos}} \otimes b_{-1/2}^M |0\rangle_{\text{NS}}, \quad \alpha_{-1}^I |0\rangle_{\text{bos}} \otimes |0, \alpha\rangle_{\text{R}}. \tag{1.1.43b}$$

The first in (1.1.43a) gives the familiar (g_{MN}, B_{MN}, ϕ) in the NSNS sector of the superstring; the second is identical to the RNS sector of the superstring. These are

⁴ At this point, one might also take more generally the lattice to be a sublattice of the colattice, but modular invariance (already mentioned for the superstring) eliminates this possibility.

Table 1.3. Massless heterotic spectrum.

NS	$\mathbf{8}_V \otimes \mathbf{8}_V = \mathbf{35}_V \oplus \mathbf{28}_V \oplus \mathbf{1}_V$	g_{MN}, B_{MN}, ϕ
R	$\mathbf{8}_V \otimes \mathbf{8}_S = \mathbf{56}_S \oplus \mathbf{8}_C$	$\psi_{M\alpha}, \lambda_{\dot{\alpha}}$
NS	$\mathbf{8}_V$	A_M^a
R	$\mathbf{8}_S$	$\chi_{\dot{\alpha}}^a$

the first two lines in Table 1.3. The (1.1.43b) are 16 abelian vector fields A_M^I and spinors $\chi_{\dot{\alpha}}^I$; they are the Cartan subalgebra part of the second two lines in Table 1.3.

Indeed, we are not done yet, because (1.1.42) allows us a second possibility: we can take $N = 0$ and look for $\hat{p}^2 = 2$, namely vectors in Γ of length two. There exist only two lattices in \mathbb{R}^{16} that are both self-dual and even; both are the root lattices $\Gamma_{\mathfrak{g}}$ for a Lie algebra \mathfrak{g} of a Lie group that can be either

$$\text{SO}(32) \quad \text{or} \quad E_8 \times E_8. \quad (1.1.44)$$

The elements of Γ of length two are the nonzero roots of \mathfrak{g} . A tensor product of all these states with $b_{-1/2}^M |0\rangle_{\text{NS}}$ and $|0, \alpha\rangle_{\text{R}}$, as in (1.1.43b), produces one vector field and one spinor for each nonzero root of \mathfrak{g} . Both (1.1.44) have rank 16, which is just the number of the A_M^I we found previously. So in total we have $A_M^a, \chi_{\dot{\alpha}}^a$ with $a = 1, \dots, \dim(\mathfrak{g})$, and we reproduce the missing part of the last two lines in Table 1.3. The A_M^a are the gauge vectors of a nonabelian gauge algebra \mathfrak{g} . This enhancement is similar to (1.1.26) for the bosonic string.

The heterotic string has the advantage of having a built-in nonabelian gauge symmetry. We will see in Section 1.3 that D-branes give an alternative way of obtaining nonabelian gauge groups G in type II, but only of the type $G = U(N)$; later in Section 1.4.4 we will see $G = \text{SO}(N)$ and $\text{Sp}(N)$. Obtaining a more interesting gauge group such as E_8 is in fact possible in IIB, but requires more sophisticated techniques that we will only study much later, in Section 9.4.⁵

1.2 Supergravity

An effective spacetime action for superstrings can be found using the same methods that led us to (1.1.17). Because of its spacetime supersymmetry, it is called *10-dimensional supergravity*.

1.2.1 RR fields

In this book, we will actually only see the action for the *bosonic* fields. The superstring NSNS fields are the same as those (1.1.9) of the bosonic string. The new

⁵ It might seem that a unitary gauge group is quite enough to accommodate the Standard Model's $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$. Even if these three factors unify at higher energies, the gauge groups that seem most promising are $G = \text{SU}(5)$ and $\text{SO}(10)$. However, the particular representations that one needs for those *grand unified theories* (GUT) are easier to obtain by starting from a group such as E_8 or its subgroup E_6 .

bosonic fields are those in the RR sector. In this subsection, we focus on them before writing the effective actions for IIA and IIB in the next subsections.

Completely antisymmetric tensors

The RR fields are all completely antisymmetric tensors with k -indices. Such tensors are also called k -forms; we give here a minimal introduction to their properties, leaving a deeper treatment to Sections 3.1 and 4.1.

The *antisymmetrized derivative* or *exterior differential*

$$(dA)_{M_1\dots M_k} \equiv k\partial_{[M_1}C_{M_2\dots M_k]} \equiv \partial_{M_1}C_{M_2\dots M_k} - \partial_{M_2}C_{M_1\dots M_k} \pm \dots, \quad (1.2.1)$$

takes a $(k-1)$ -form to a k -form. The index antisymmetrizer $[M_1\dots M_k]$ sums over all permutations with a ± 1 equal to the sign $\sigma(i_1\dots i_k)$ of the permutation taking $1\dots k$ to $i_1\dots i_k$, and divides by a $k!$. So, for example, $(dA)_{MN} = \partial_M C_N - \partial_N C_M$ is the Maxwell field-strength of C_M . As another example, (1.1.15) can be written as $H_{MNP} = (dB)_{MNP} = 3\partial_{[M}B_{NP]}$, with only three terms instead of six because of antisymmetry of B . To avoid index proliferation, often we will write $C_{M_1\dots M_k}$ symbolically simply as C_k , and denote (1.2.1) by dC_k . An important property of (1.2.1) is

$$dd\alpha_k = 0. \quad (1.2.2)$$

Explicitly,

$$(dd\alpha)_{M_1\dots M_k} = k(k-1)\partial_{[M_1}\partial_{M_2}\alpha_{M_3\dots M_k]} = 0, \quad (1.2.3)$$

because $\partial_{[M_1}\partial_{M_2]} = 0$ on smooth functions. For an electromagnetic potential A_M , this gives $\partial_{[M}F_{NP]} = \partial_{[M}\partial_N A_P] = 0$, one of the Maxwell equations. As another example, from (1.1.15) it now follows $\partial_{[M}H_{NPQ]} = 0$, or in other words,

$$dH = 0. \quad (1.2.4)$$

The antisymmetrized, or *wedge*, product of two forms $\alpha_k, \alpha_{k'}$ is defined as

$$(\alpha \wedge \alpha')_{M_1\dots M_k N_1\dots N_{k'}} \equiv \frac{(k+k')!}{k!k'!} \alpha_{[M_1\dots M_k} \alpha'_{N_1\dots N_{k'}]}. \quad (1.2.5)$$

This satisfies $\alpha_k \wedge \alpha_{k'} = (-1)^{kk'} \alpha_{k'} \wedge \alpha_k$. In particular,

$$H \wedge H = 0. \quad (1.2.6)$$

We now also have the Leibniz identity:

$$d(\alpha_k \wedge \alpha_{k'}) = d\alpha_k \wedge \alpha_{k'} + (-1)^k \alpha_k \wedge d\alpha_{k'}. \quad (1.2.7)$$

In $d = 10$, the largest possible number of indices of a k -form is $k = 10$. Moreover, a ten-form α_{10} is unique up to rescaling: it must be proportional to the Levi-Civita tensor $\epsilon_{M_1\dots M_{10}}^{(0)} \equiv \sigma(M_1\dots M_{10})$, the sign of the permutation taking $M_1\dots M_{10}$ to $0\dots 9$. (The (0) label is used here because a more mathematically sophisticated version of this tensor will enter the scene later.) Writing then $\alpha_{0\dots 9} = f$, we define

$$\int \alpha_{10} \equiv \int d^{10}x f. \quad (1.2.8)$$

A similar definition holds in any dimension d .

Twisted field-strengths

RR forms usually appear in the action through a *twisted* field-strength⁶

$$F_k \equiv dC_{k-1} - H \wedge C_{k-3}. \quad (1.2.9)$$

For example, we will soon encounter $F_{MNPQ} = 4\partial_{[M}C_{NPQ]} - 4H_{[MNP}C_{Q]}$. (The factors come from (1.2.1) and (1.2.5).) For $k < 3$, the second term is absent; for example, $F_2 \equiv dC_1$. The F_k satisfy a *Bianchi identity*:

$$dF_k = H \wedge F_{k-2}, \quad (1.2.10)$$

as one sees using (1.2.2), (1.2.4), and (1.2.6). In a sense, this is more fundamental than (1.2.9): sometimes we will modify (1.2.9), with (1.2.10) remaining true. Later in this chapter, we will consider sources, which will violate (1.2.10) on some spacetime defects. A form field H or F_k satisfying (1.2.10) with no source is often called a *flux*. By an abuse of language, one sometimes calls by this name any form field, sourced or not.

Gauge transformations

In electromagnetism, the field-strength $F_{MN} = \partial_M A_N - \partial_N A_M$ is invariant under the gauge transformation $A_M \rightarrow \partial_M \lambda_0$. We already observed after (1.1.15) that the Kalb–Ramond field has a gauge transformation $B \rightarrow B + d\hat{\lambda}_1$, which leaves invariant its field-strength $H = dB$. For the RR fields, (1.2.9) are invariant under the gauge transformations:

$$\delta C_k = d\lambda_{k-1} - H \wedge \lambda_{k-3}. \quad (1.2.11)$$

So, for example, under λ_1 we have $\delta C_2 = d\lambda_1$, $\delta C_4 = -H \wedge \lambda_1$, while under λ_3 we have $\delta C_4 = d\lambda_3$, $\delta C_2 = 0$.

In electromagnetism, λ_0 need not be a function: it can be multivalued, as long as

$$e^{iq_e \lambda_0} \quad (1.2.12)$$

is single-valued, where q_e is the elementary electric charge. Multivalued λ_0 are called a *large* gauge transformation. They exist because the gauge group is compact, which in turn comes from the requirement that the wave function should be single-valued. This is relevant when there is a nontrivial loop in spacetime, either because of an excluded region (as in the Aharonov–Bohm effect) or because spacetime has a nontrivial topology. Consider for example an S^1 with a periodic coordinate $x \sim x + L$. Single-valuedness of (1.2.12) requires $\lambda_0 \rightarrow \lambda_0 + 2\pi N/q_e$, $N \in \mathbb{Z}$. The gauge transformation $A_M \rightarrow A_M + \partial_M \lambda_0$ is more correctly rewritten as

$$A_M \rightarrow A_M + \Lambda_M, \quad (1.2.13)$$

where Λ_M is required to have $\int dx \Lambda_x = 2\pi N/q_e$.

Suppose we want to give a *constant* expectation value to a component, $A_x = A_x^0$. This has no physical meaning if the direction x is noncompact, because we can gauge it away with a gauge transformation $\lambda = x A_x^0$. But if $x \sim x + L$ is a coordinate on a

⁶ Many authors use the symbol F_k for dC_{k-1} alone, and then put a tilde on our (1.2.9).

circle, then with a periodic λ we cannot gauge away A_x^0 any more; with a large gauge transformation, (1.2.13) gives us the identification

$$A_x^0 \cong A_x^0 + \frac{2\pi}{Lq_e}. \quad (1.2.14)$$

In the limit $L \rightarrow \infty$ we see again that all constant values can be gauged away. Similar large generalizations of (1.2.11) exist; we will discuss them after we introduce the analogue of the elementary electric charge for the C_k in Section 1.3.

1.2.2 IIA supergravity

We now come to the effective action for IIA string theory. As we anticipated, we will only show the action for the bosonic fields.

Bosonic action of IIA supergravity

The leading bosonic action for type IIA superstrings is called type IIA supergravity. It reads

$$S_{\text{IIA}} = S_{\text{kNS}} + \frac{1}{4\kappa^2} \left[- \int d^{10}x \sqrt{-g} (|F_2|^2 + |F_4|^2) + \int B \wedge dC_3 \wedge dC_3 \right], \quad (1.2.15)$$

where $|F_k|^2 \equiv \frac{1}{k!} F_{M_1 \dots M_k} F^{M_1 \dots M_k}$, extending the definition in (1.1.15);

$$2\kappa^2 = (2\pi)^7 l_s^8; \quad (1.2.16)$$

and the kinetic NSNS term

$$S_{\text{kNS}} = \frac{1}{2\kappa^2} \int d^{10}x \sqrt{-g} e^{-2\phi} \left(R + 4\partial_\mu \phi \partial^\mu \phi - \frac{1}{2} |H|^2 \right) \quad (1.2.17)$$

is the same Lagrangian as in (1.1.17), only now integrated in $d = 10$ dimensions rather than 26. This term will also appear in all the supergravity effective actions we will see later. The $e^{-2\phi}$ prefactor signals its origin from $g = 0$ string diagrams, as we remarked following (1.1.17). In a region where the dilaton is constant, the Planck length is now

$$l_P = g_s^{1/4} l_s. \quad (1.2.18)$$

The string scattering amplitudes originally give a prefactor $e^{-2\phi}$ for the rest of (1.2.15) as well, but the action happens to look nicer if one rescales it away by redefining the C_k .

RR terms

The first parenthesis in (1.2.15) can be regarded as the kinetic term for $F_2 = dC_1$, and for $F_4 = dC_3 - H \wedge C_1$ from (1.2.9). The last term is more peculiar in (1.2.15). Unpacking the form notation (1.2.1), (1.2.5), and (1.2.8), one obtains:

$$\int B \wedge dC_3 \wedge dC_3 = \frac{10!}{2 \cdot (3!)^2} \int d^{10}x B_{[01} \partial_2 C_{345} \partial_6 C_{789]}. \quad (1.2.19)$$

It does not involve the metric, and it contains a potential B_2 without an exterior derivative. For this reason, it is called the *Chern–Simons* term, after the Chern–Simons action in three dimensions $S_{CS} = \int_{M_3} CS_A$, where

$$CS_A \equiv \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \quad (1.2.20)$$

is a three-form with the property $dCS_A = \text{Tr}(F \wedge F)$; the two-form $F = dA + A \wedge A$ is the nonabelian field-strength in the form notation of Section 1.2.1. Alternative expressions for (1.2.19), such as $-\int H \wedge C_3 \wedge dC_3$, can be obtained by integration by parts. Note that if we worked in $d = 11$ rather than $d = 10$, we could compute⁷

$$d(B \wedge dC_3 \wedge dC_3) = H \wedge dC_3 \wedge dC_3 \stackrel{(1.2.9),(1.2.6)}{=} H \wedge F_4 \wedge F_4. \quad (1.2.21)$$

Frame change

Superficially, the sign of the kinetic term for ϕ in (1.2.17) looks wrong: it is not of the usual form “kinetic energy minus potential energy,” since $\dot{\phi}^2$ appears with a minus sign. (This issue of course was already present in (1.1.17).) But ϕ also appears multiplying the Einstein–Hilbert term R ; so its dynamics is less simple than it looks. To make it more transparent, one can define an alternative metric

$$g_{MN}^E \equiv e^{-\phi/2} g_{MN}. \quad (1.2.22)$$

This is called *Einstein frame* metric, because in terms of it (1.2.17) becomes

$$S_{E,\text{KNS}} = \frac{1}{2\kappa^2} \int d^{10}x \sqrt{-g_E} \left(R_E - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} e^{-\phi} |H|^2 \right), \quad (1.2.23)$$

with the Einstein–Hilbert term now appearing without the dilaton prefactor; indices are also now contracted with the Einstein frame metric. Here the dilaton’s kinetic term has the conventional sign. In most of the book, we will use the original *string frame* metric g_{MN} .

Supersymmetry

As we mentioned in Section 1.1.2, type II superstrings are symmetric under the 32 supercharges ϵ^a ; so the fermionic completion of the supergravity action (1.2.15) will enjoy such symmetry too. Theories with 32 supercharges are relatively rare: besides IIB, an important example we will see later in this chapter is eleven-dimensional supergravity, of which IIA is a *dimensional reduction*.⁸ This large amount of supersymmetry has consequences on the structure of the l_s and g_s corrections; for example, it fixes the two-derivative action completely.

This is a symmetry of the *theory*: not all supercharges will leave invariant a particular field configuration. The generic field configuration breaks supersymmetry

⁷ Equation (1.2.21), together with a generalization of Stokes’s theorem that we will study in Section 4.1.10, allows us to reinterpret the CS term as an integral over an eleven-dimensional space of which the ten-dimensional one is a boundary.

⁸ Most other theories with 32 supercharges are further dimensional reductions of IIB or eleven-dimensional supergravity, some of which we’ll see in Chapter 11; but there now exist examples which are not thought to arise this way [32].

completely, just like the generic metric g_{MN} has no Killing vectors. One of the topics of interest for this book will be the study of how many supercharges leave invariant given field configurations, or vice versa of which field configurations are invariant under a certain number of supercharges.

String corrections

Corrections to the supergravity approximation first occur at the eight-derivative level. The curvature terms read

$$S_{\text{IIA},R^4} = \frac{1}{2\kappa^2} \frac{l_s^6 \zeta(3)}{3 \cdot 2^{11}} \int d^{10}x \sqrt{-g} e^{-2\phi} \left(t_{M_1 \dots M_8} t^{N_1 \dots N_8} + \frac{1}{8} \epsilon_{PQM_1 \dots M_8} \epsilon^{PQN_1 \dots N_8} \right) \cdot R^{M_1 M_2}_{N_1 N_2} R^{M_3 M_4}_{N_3 N_4} R^{M_5 M_6}_{N_5 N_6} R^{M_7 M_8}_{N_7 N_8} \tag{1.2.24}$$

at the leading order in g_s . Here ζ is Riemann’s zeta function, and the tensor t is defined by

$$t_{M_1 \dots M_8} M^{M_1 M_2} M^{M_3 M_4} M^{M_4 M_5} M^{M_7 M_8} = 24 \text{Tr} M^4 - 6(\text{Tr} M^2)^2. \tag{1.2.25}$$

Just like for supergravity, (1.2.24) can be obtained either by computing string amplitudes [33], or by computing the world-sheet beta functions beyond the leading approximation [34]. At $g = 1$, or in other words $e^{0 \cdot \phi}$ according to (1.1.12), there is a similar term, where the combination $t_8 t_8 - \epsilon \epsilon / 8$ appears.

Even at this l_s^8 level, the complete structure of the action is not completely established beyond (1.2.24). For example, one expects couplings to the form field strengths. There is a famous coupling of the type $\int B_2 R^4$, related to anomalies [35, 36]. One can try to infer the remaining terms by using dualities [37] or supersymmetry [38]. A complementary approach is via string amplitudes in the four-field approximation (which gives a contribution to (1.2.24), when one linearizes it in the metric fluctuation $g_{MN} \sim \eta_{MN} + h_{MN}$); with this restriction, one can go to arbitrary precision in l_s in the aforementioned Berkovits formalism [39].

Massive IIA

IIA supergravity has a deformation by a parameter F_0 called *Romans mass* [40], still preserving 32 supercharges.⁹ This theory was originally obtained by realizing a *Stückelberg mechanism*, by which the Kalb–Ramond field B acquires a mass, at the price of absorbing (or “eating”) the degrees of freedom of C_1 .

The original Stückelberg mechanism [43] is a variant of (and predates) the more familiar Brout–Englert–Higgs (BEH) mechanism. In both cases, a vector field A_μ acquires a mass by eating the degrees of freedom of a scalar a . In BEH, we gauge a rotation in field space, leading to a covariant derivative schematically of the form $\partial_\mu a + A_\mu a$. In the Stückelberg case, we instead introduce

$$D_\mu a \equiv \partial_\mu a + mA_\mu, \tag{1.2.26}$$

where m is a mass. This is invariant under a translation: $a \rightarrow a - m\lambda$, if also $A_\mu \rightarrow A_\mu + \partial_\mu \lambda$. This transformation can of course be used to set $a = 0$; in this sense, a

⁹ There is in fact also a second deformation of IIA with 32 supercharges [41]; this arises as a dimensional reduction of eleven-dimensional supergravity where one identifies spacetime with itself up to an overall rescaling. These are the only maximally supersymmetric deformations [42].

is “eaten” by A_μ . The covariant kinetic term $D_\mu a D^\mu a$, when expanded, is now seen to contain a mass term $m^2 A_\mu A^\mu$ for the vector field. Equation (1.2.26) appears often in string compactifications with fluxes. We will comment further on the difference between BEH and Stückelberg, and generalize both, in Section 4.2.2.

The variant of (1.2.26) we need in IIA is obtained by adding an index to both participant fields, thus introducing

$$F_2 \equiv dC_1 + F_0 B. \quad (1.2.27)$$

This modifies (1.2.9), which would have given $F_2 = dC_1$. Now C_1 and B plays the role of a and A_μ in (1.2.26), respectively. We have now used the symbol F_0 for the mass parameter. The reason for this name becomes apparent when we notice that

$$dF_2 = HF_0. \quad (1.2.28)$$

This is of the form (1.2.10), suggesting that F_0 should be regarded as a new, non-dynamical RR “field-strength” with zero indices. We will see later more cogent reasons for this interpretation.

We also have to change F_4 with respect to (1.2.9):

$$F_4 = dC_3 - H \wedge C_1 + \frac{1}{2} F_0 B \wedge B; \quad (1.2.29)$$

(1.2.10) still holds. The gauge transformations (1.2.11) still leave these invariant, but that of B does not, unless we modify it:

$$\delta B = d\hat{\lambda}_1, \quad \delta C_1 = -F_0 \hat{\lambda}_1, \quad \delta C_3 = -F_0 \hat{\lambda}_1 \wedge B. \quad (1.2.30)$$

With these new definitions for the F_k , the IIA action is now modified as

$$S_{\text{IIA}} = S_{\text{KNS}} + \frac{1}{4\kappa^2} \left[- \int d^{10}x \sqrt{-g} (F_0^2 + |F_2|^2 + |F_4|^2) + \int \text{mCS}_{10} \right], \quad (1.2.31)$$

where mCS_{10} is a ten-form that again has the formal property

$$d(\text{mCS}_{10}) = H \wedge F_4 \wedge F_4 \quad (1.2.32)$$

as in (1.2.21). Viewed in this way the modification with respect to (1.2.15) is minimal, although an explicit expression is less nice:

$$\text{mCS}_{10} = B \wedge \left(dC_3^2 + \frac{1}{3} F_0 dC_3 \wedge B^2 + \frac{1}{20} F_0^2 B^4 \right), \quad (1.2.33)$$

where $B^k \equiv B \wedge \dots \wedge B$.

We will see a more elegant way of understanding both (1.2.15) and (1.2.31) in Section 10.1, at the cost of a slightly more sophisticated mathematical apparatus. On that occasion, we will also see the supersymmetry transformations.

1.2.3 IIB supergravity

IIB supergravity is very similar to IIA [44–46]. The most important difference is that it is a *chiral* theory: in Table 1.2, we see that the two ψ^a and the two λ^a have the same chirality. Usually the presence of chiral spinors threatens an anomaly.

While there are no gauge fields, in a gravitational theory we have to worry about potential anomalies for *diffeomorphisms*, which are just as lethal as their gauge counterparts. The chiral ψ^a and λ_a indeed give a nonvanishing contribution to such a diffeomorphism anomaly. Fortunately, however, the RR field C_{MNPQ}^+ also gives a contribution, which exactly cancels the fermionic ones [47], avoiding a potential catastrophe in a nontrivial way.

C_{MNPQ}^+ gives a contribution to an anomaly in spite of being bosonic because of its self-duality property:

$$F_{M_1\dots M_5} = \frac{1}{5!} \sqrt{-g} \epsilon_{M_1\dots M_5}^{(0)}{}^{M_6\dots M_{10}} F_{M_6\dots M_{10}}, \tag{1.2.34a}$$

where again $\epsilon^{(0)}$ is the completely antisymmetric tensor defined later in (1.2.6). In the condensed notation of the previous subsection, (1.2.34a) is written as

$$F_5 = *F_5. \tag{1.2.34b}$$

Just like in (1.2.9), we can take $F_5 = dC_4^+ - H \wedge C_2$. There are alternatives, such as $F'_5 = dC_4^+ + \frac{1}{2}(B \wedge dC_2 - H \wedge C_2)$. Both F_5 and F'_5 satisfy the Bianchi identities (1.2.10); one can bring one into the other by redefining C_4^+ (with no ill effect on the action (1.2.36)).

Pseudoaction

While self-duality of F_5 saves the theory, it also makes it hard to write down an action. A self-duality property similar to (1.2.34) can be introduced for any k -form potential a_k with self-dual field-strength $da_k = *da_k$ in $d = 2(k + 1)$ dimensions. The simplest example is for $k = 0$: a_0 is then a scalar in $d = 2$ dimensions, and self-duality implies

$$\partial_M a_0 \partial^M a_0 = \sqrt{-g} \epsilon_{(0)}^{MN} \partial_M a_0 \partial_N a_0 = 0, \tag{1.2.35}$$

so the usual Lagrangian density vanishes. More generally, this happens for any even k , and in particular for our (1.2.34): the naive Lagrangian density $|F_5|^2 = 0$. There are strategies to cope with this issue; see [48, 49] for a recent proposal. In the following, we will simply write a *pseudoaction*: all equations of motion can be derived by varying it, *except* for the constraint (1.2.34). From now on, we will drop the superscript and call $C_4^+ \rightarrow C_4$, for uniformity with the other potentials in both type II theories.

A pseudoaction for IIB is then

$$S_{\text{IIB}} = S_{\text{KNS}} + \frac{1}{4\kappa^2} \left[- \int d^{10}x \sqrt{-g} \left(|F_1|^2 + |F_3|^2 + \frac{1}{2} |F_5|^2 \right) + \int B \wedge dC_2 \wedge dC_4 \right], \tag{1.2.36}$$

where S_{KNS} was given in (1.2.17). As in IIA, the last term is of Chern–Simons type, and has the formal property that in eleven dimensions $d(B \wedge dC_2 \wedge dC_4) = H \wedge F_3 \wedge F_5$, similar to (1.2.21). Alternative expressions can be obtained by integration by parts, such as $-\int H \wedge C_2 \wedge dC_4$. The fermionic completion of (1.2.36) is supersymmetric with 32 supercharges, just like IIA.

The leading l_s corrections are the same as in IIA, (1.2.24). At the $g = 1$ level, the same combination $t_8 t_8 - \epsilon\epsilon/8$ is present. Knowledge of the full eight-derivative corrections is as incomplete as in IIA, but in this case dualities are more powerful and determine the dependence on ϕ of the terms in (1.2.24) beyond the $g = 0$ and $g = 1$ terms [50]. This is based on an important additional symmetry of IIB that is already present at the level of supergravity, to which we now turn.

SL(2, \mathbb{R}) symmetry

Given a 2×2 matrix with unit determinant,

$$m = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{R}), \quad (1.2.37)$$

we define the transformation law

$$\begin{aligned} \tau &\rightarrow m \cdot \tau \equiv \frac{a\tau + b}{c\tau + d}, & g_{MN} &\rightarrow |c\tau + d| g_{MN} \\ F_5 &\rightarrow F_5, & \begin{pmatrix} C_2 \\ B \end{pmatrix} &\rightarrow m \begin{pmatrix} C_2 \\ B \end{pmatrix}, \end{aligned} \quad (1.2.38)$$

where

$$\tau \equiv C_0 + ie^{-\phi} \quad (1.2.39)$$

is called the *axiodilaton*.

Equation (1.2.38) is a symmetry of the action (1.2.36). This type of nonlinear action on τ is called a *Möbius transformation*; it will appear in several other contexts. The string-frame metric transforms, but the Einstein frame metric (1.2.22) does not:

$$g_{MN}^E \rightarrow g_{MN}^E. \quad (1.2.40)$$

There is also an alternative expression for the transformation law of the two-form potentials, in terms of

$$G_3 \equiv dC_2 - \tau H = dC_2 - HC_0 - ie^{-\phi} H = F_3 - ie^{-\phi} H. \quad (1.2.41)$$

From (1.2.38):

$$G_3 \rightarrow (a dC_2 + bH) - \frac{a\tau + b}{c\tau + d} (c dC_2 + dH) \stackrel{(1.2.37)}{=} \frac{1}{c\tau + d} G_3. \quad (1.2.42)$$

To see why this symmetry is remarkable, consider for example $s = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Starting from a configuration where $C_0 = 0$, we see that

$$e^\phi \rightarrow e^{-\phi}. \quad (1.2.43)$$

The string coupling $g_s \rightarrow 1/g_s$: strong coupling is mapped to weak coupling.

We hasten to add that this is a symmetry of the *supergravity* approximation; it does not fully survive in string theory. One reason is the general expectation that theories of quantum gravity should have no continuous symmetries, which we will review in Chapter 12. A perhaps more concrete argument is that the fundamental

string couples to B ; an arbitrary transformation in (1.2.38) transforms $B \rightarrow cC_2 + dB$, and if this were a symmetry for arbitrary c and d there would have to exist a continuum of string-like objects, coupling to any such linear combination. We will see in Section 1.4.3 that generalizations of the fundamental string do exist, but only a *discrete* infinity of them. This argument suggests that $SL(2, \mathbb{R})$ gets discretized; we will see this in Section 1.4.3.

1.2.4 Heterotic supergravity

Action

The bosonic part of the action for heterotic supergravity is

$$S_{\text{het}} = S_{\text{kNS}} + \frac{l_s^2}{8\kappa^2} \int_{M_{10}} d^{10}x \sqrt{-g_{10}} e^{-2\phi} \text{Tr}|F^2|. \quad (1.2.44)$$

(The positive sign is because $\text{Tr}(T^a T^b)$ is negative-definite; we do not include i 's in the gauge fields.) The term S_{kNS} is the usual (1.2.17), but now

$$H \equiv dB - \frac{l_s^2}{4} CS_A, \quad (1.2.45)$$

where CS_A was given in (1.2.20). As in type II supergravity, there are l_s corrections; some of these will play a role in anomaly cancellation, to which we now turn.

Anomaly cancellation

Since we only supersymmetrized the right-movers, (1.2.44) only has half the supersymmetry of the type II theories: 16 supercharges. Each line marked R in Table 1.3 is related by supersymmetry to the line marked NS above it. As in IIB, the spectrum is chiral: the gravitino $\psi_{M\alpha}$ and the gaugino χ_α have positive chirality, while the $\lambda_{\dot{\alpha}}$ have negative chirality. Again, this creates the danger of an anomaly, this time not only for diffeomorphisms but also for gauge transformations. Superficially, this does not appear to vanish, but exactly for the two gauge groups (1.1.44) it takes the factorized form

$$\delta\Gamma = \int \omega_2^1 \wedge Y_8. \quad (1.2.46)$$

Here the symbol δ denotes variation under both diffeomorphisms and gauge transformations; as usual in QFT, Γ is the quantum effective action (the action whose tree-level amplitudes equal the full quantum amplitudes of S). Y_8 is an eight-form, quartic in curvature and in the Yang–Mills field-strength F^a . The two-form ω_2^1 is such that

$$d\omega_2^1 = \delta(CS_g - CS_A), \quad (1.2.47)$$

where now CS_g is a three-form with the property $(dCS_g)_{MNPQ} = 6R^{AB}{}_{[MN}R_{PQ]BA}$, similar to (1.2.20).

Equation (1.2.46) is nonzero, but it is not the full story; there are contributions from the higher-derivative corrections. First of all, (1.2.45) should be modified to

$$H \equiv dB + \frac{l_s^2}{4} (CS_g - CS_A). \quad (1.2.48)$$

CS_g contains up to three derivatives, since its derivative contains by definition two Riemann tensors. Now (1.2.47) implies $\delta H \neq 0$, unless we also make B transform as $\delta B = -\frac{l_s^2}{4}\omega_2^1$. Another l_s correction is the term

$$\int B \wedge Y_8; \quad (1.2.49)$$

its transformation now cancels the anomaly (1.2.46). This only works for the two choices (1.1.44) of the gauge group, which allowed us to write (1.2.46) in the first place. For more general gauge groups, $\delta\Gamma$ does not factorize.

Exercise 1.2.1 Compute the equations of motion for the RR fluxes in IIA and IIB by varying the actions (1.2.15), (1.2.31), and (1.2.36) with respect to the C_p . Check that they are formally identical to the Bianchi identities (1.2.10), if we extend them to $k \geq 6$ by defining

$$F_6 = *F_4, \quad F_7 = -*F_3, \quad F_8 = -*F_2, \quad F_9 = *F_1, \quad F_{10} = *F_0, \quad (1.2.50)$$

in the notation of (1.2.34b).

Exercise 1.2.2 Use (1.2.38) and (1.2.39) to show

$$e^\phi \rightarrow |c\tau + d|^2 e^\phi. \quad (1.2.51)$$

1.3 D-branes

We saw that string theory has two expansion parameters: l_s (or rather the dimensionless l_s/r , with r a curvature radius), and e^ϕ , which when constant is also called g_s . In (1.1.21), we wrote the expansion as a power series, but we wondered whether this really captures the whole dependence. Recall that a function is called (real) analytic if it coincides with its Taylor series around every point. A famous nonanalytic function is

$$f(g) = e^{-1/g^2}. \quad (1.3.1)$$

Both f and all its derivatives vanish at $g = 0$, so its Taylor series vanishes identically, but $f(g) \neq 0$. In physics, we call an effect with such a dependence on the coupling *nonperturbative*. In attempts at describing nonperturbative aspects of field theory, two types of objects often come up: solitons, which are large and stable field configurations, localized in space; and instantons, which occur in the Euclidean path integral and are localized in time as well. In string theory, the most prominent objects that play these roles are *D-branes*; this section is devoted to them.

1.3.1 Solitons and instantons

We begin with a quick reminder of solitons and instantons in field theory.

Solitons

In perturbative field theory, we often deal with quanta of small field perturbations around the vacuum, which we call elementary particles. However, many field theories

also contain other objects. A *soliton* is a “localized” solution of the equations of motion that does not dissipate. This is in contrast with plane waves, which are delocalized. Already in a linear system we find both types of solutions. For example, for a free scalar ϕ in $d = 2$, the equation of motion $(-\partial_t^2 + \partial_x^2)\phi = 0$ is solved by $\phi = \phi_0(t - x)$ for any ϕ_0 ; this includes the plane wave solutions $\phi = e^{i(t-x)}$, but we can also take ϕ_0 to be any localized function, such as $\phi = (\cosh(x - t))^{-2}$, which is very small everywhere except for a narrow band of order one.

More notably, solitons exist in nonlinear systems also: the venerable Korteweg–De Vries (KdV) equation

$$\partial_t \phi + \partial_x^3 \phi + 6\phi \partial_x \phi = 0, \tag{1.3.2}$$

describing water waves in shallow channels, has many solitonic solutions, such as

$$\phi = \frac{c}{2} \left(\cosh \left(\frac{\sqrt{c}}{2} (x - ct) \right) \right)^{-2} \tag{1.3.3}$$

for any c . Once again, this is localized in a narrow band at any t , and velocity c . There also exist *multisoliton* solutions, where many waves similar to (1.3.3) coexist and interact, just like particles. The dissipative and nonlinear effects from the second and third term of (1.3.2) compete in exactly such a way that these waves do not dissipate. (At a deeper level, this phenomenon is really due to the presence of infinitely many conserved quantities for (1.3.2); see, for example, [51].)

Going back to relativistic field theories, many nonlinear theories also have such stable solutions. Any $d = 2$ theory of a single scalar ϕ with a potential $V(\phi)$ with two or more vacua ϕ_{\pm} will have solitons: in this case, they are field configurations where $\phi \rightarrow \phi_{\pm}$ at $x \rightarrow \pm\infty$. This is protected from dissipation by topological reasons: one cannot take such a configuration to a vacuum without an infinite energy expenditure. Unlike (1.3.3), these solitons can be at rest. Solitons also exist for $d > 2$, and again they are usually stable for topological reasons. They are usually heavy when the theory is weakly coupled. Nevertheless, just like the soliton waves of the KdV equation, they behave almost as particles, and we should be able to compute their dynamics, for example, their scattering amplitudes.

Monopoles

A *magnetic monopole* is a soliton around which the flux integral of the magnetic field is nonzero. In \mathbb{R}^3 , this means that

$$\int_{S_2} da_i B_i = q_m \neq 0 \tag{1.3.4}$$

on a sphere S_2 . da_i is an outward-directed vector whose norm da is the infinitesimal area. Usually in electromagnetism $q_m = 0$, because of the Maxwell equation $\partial_i B_i = 0$, the space part of $\partial_{[\mu} F_{\nu\rho]} = 0$. So to introduce such objects, we should change this equation to $\partial_i B_i = \Pi_{i=1}^3 \delta^i(x^i)$. A striking feature is their *charge quantization*. Suppose we have a monopole localized at the origin. Consider a potential A_{μ}^N , and integrate it on the equator E of S_2 . Since $B_i = \epsilon_{ijk} \partial_j A_k^N$, we can apply Stokes’s theorem to the semisphere U_N of S_2 bounded by E :

$$\oint_E dx_i A_i^N = \int_{U_N} da_i B_i. \tag{1.3.5}$$

However, we could do the same with the other portion of the sphere U_S ; because of orientation, there is a minus sign, so this time we would have $\oint_E dx_i A_i^N \stackrel{?}{=} -\int_{U_S} da_i B_i$. Subtracting this from (1.3.5), we would find $\int_{S_2} da_i B_i \stackrel{?}{=} 0$, a contradiction. The way out is to have a second potential A_μ^S for the region U_S ; then $\oint_E dx_i A_i^S = -\int_{U_S} da_i B_i$, and subtraction from (1.3.5) now just gives

$$\int_{S_2} da_i B_i = \oint_E dx_i (A^N - A^S)_i = q_m. \tag{1.3.6}$$

Two potentials are related by a gauge transformation, so $(A^N - A^S)_i = \partial_i \lambda_0$; now (1.3.6) tells us that λ_0 is not periodic, but undergoes a shift q_m after a turn around E . Now, a gauge transformation acts on the wave function of a particle with electric charge q_e by $\psi \rightarrow e^{-iq_e \lambda_0} \psi$. This should be periodic, so $e^{-iq_e q_m} = 1$, or in other words,

$$\frac{1}{2\pi} q_e q_m \in \mathbb{Z}. \tag{1.3.7}$$

This is called *Dirac quantization*, and will play many roles in this book.¹⁰

Monopole solutions can be found in the Yang–Mills–Higgs model

$$S_{\text{YMH}} = - \int d^4x \text{Tr} \left(\frac{1}{2g_{\text{YM}}^2} |F|^2 + D_\mu a D^\mu a + \lambda (a^2 - a_0^2)^2 \right), \tag{1.3.8}$$

where F is an $SU(2)$ gauge field, a an adjoint scalar, and $D_\mu a \equiv \partial_\mu a + [A_\mu, a]$ the gauge covariant derivative. For the *'t Hooft–Polyakov monopole* solutions, q_m is defined as the magnetic charge under $\text{Tr}(aF)$, and all fields are nonsingular. One can prove the *Bogomolnyi–Prasad–Sommerfield (BPS) bound* for the mass of any monopole solution:

$$m_{\text{mon}} \geq a_0 q_m = \frac{2\pi a_0}{g_{\text{YM}}}. \tag{1.3.9}$$

As anticipated, the mass is large at weak coupling. Conversely, at strong coupling they may become light.

These effects are under better control in supersymmetric theories. A famous example is the $\mathcal{N} = 2$ -supersymmetric version of Yang–Mills (*super-YM*), which is a bit similar to (1.3.8). The $\mathcal{N} = 1$ super-YM involves the gauge field A_μ and a gaugino λ_α . The $\mathcal{N} = 2$ version involves *two* gauginos, and a scalar a (now complex), all in the adjoint representation of the gauge group. The bosonic Lagrangian is¹¹

$$S_{\text{SYM}} = \int d^4x \text{Tr} \left(-\frac{1}{2g^2} |F|^2 + \frac{\theta}{64\pi^2} \epsilon_{(0)}^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} - D_\mu a^\dagger D^\mu a - \frac{1}{2} \text{Tr}([a, a^\dagger])^2 \right). \tag{1.3.10}$$

¹⁰ Monopoles in electromagnetism have never been detected; but in quantum chromodynamics (QCD), the theory of strong interactions, their condensation is believed to play an important role in confinement. The mechanism is similar to the Meissner effect in superconductors, where the electric field effectively acquires a mass and the magnetic field is zero almost everywhere, except in thin tubes. The same should happen in QCD, but now with the electric field confined in thin tubes; since its flux lines no longer disperse, this leads to a potential that grows with distance.

¹¹ The θ term is a total derivative, but it does have physical effects on instantons and on monopoles. In QCD, it is allowed but observed to be $< 10^{-9}$ for unknown reasons (the *strong CP* problem). In supersymmetric theories, it appears naturally.

The anticommutators of the $\mathcal{N} = 2$ algebra read

$$\{Q^I, \bar{Q}_J\} = P_\mu \gamma^\mu \delta^I_J, \quad \{Q^I, Q^J\} = \epsilon^{IJ} Z. \tag{1.3.11}$$

The generator Z commutes with all other generators in the algebra, and as such it is called *central charge*, just like c in (1.1.19). The BPS bound is now reinterpreted as $m_{\text{mon}} \geq |Z|$. The representation theory of (1.3.11) shows that BPS states form a special short representation; because of this, they are protected against time evolution and against deformations of the theory, so they cannot just disappear as the coupling is changed.

There is a low-energy effective description where we only keep the “abelian” part of the fields, in the Cartan subalgebra. For $G = \text{SU}(2)$, this is $\text{U}(1)$, which we can take along the σ^3 generator of $\text{su}(2)$; so we keep A_μ^3 and its supersymmetric partners. The bosonic part of the effective action is now an Abelian version of (1.3.10), but with both $g = g(a_3)$ and $\theta = \theta(a_3)$ depending on the vacuum expectation value for the scalar a_3 . $\mathcal{N} = 2$ supersymmetry determines this *Seiberg–Witten (SW) effective theory* [52] exactly as $\tau(a_3) = \partial_{a_3}^3 \mathcal{F}$, where the *prepotential* \mathcal{F} is a holomorphic function of a_3 , and we defined it:

$$\tau \equiv \frac{4\pi i}{g_{\text{YM}}^2} + \frac{\theta}{2\pi}. \tag{1.3.12}$$

In this solution, monopoles do become light at strong coupling, and there is a “dual” description of the theory, of which they are the elementary photons, described by a vector \tilde{A}_μ .

In conclusion, solitons behave a lot like particles; at weak coupling, they are collective excitations, but at strong coupling they may become the fundamental degrees of freedom. So in quantum field theory it is not always clear which objects are made of which others.

Instantons

In cases where exact results are available, such as in the $\mathcal{N} = 2$ SW theory, the effective action is not analytic: it does not coincide with its perturbative expansion. The prepotential \mathcal{F} depends on a_3 , but after reexpressing it in terms of the high-energy g_{YM} , we find a sum of contributions of the type

$$e^{-s/g_{\text{YM}}^2}, \tag{1.3.13}$$

with $s = 8\pi^2 k$, for k an integer.

Such effects are ubiquitous: they appear even in quantum mechanics, in the dependence on \hbar of the energy spectrum, or of the tunnel effect probability (see, for example, [53–55]). The easiest example is perhaps the energy of the vacuum in a double-well potential $V_{\text{dw}} = \lambda(x^2 - x_0^2)^2$. Near each vacuum $x = \pm x_0$, it is approximately harmonic, $V_{\text{dw}} \sim \frac{1}{2}\omega^2(x \mp x_0)^2$, $\omega^2 = 8x_0^2\lambda$; so we expect the lowest energies to be $\sim \frac{\hbar}{2}\omega$, with a small splitting due to tunneling between the two. To find this, one computes the probability $\langle x_0 | e^{-iHT/\hbar} | -x_0 \rangle$, which becomes an integral over histories:

$$\int Dx(t) e^{-iS/\hbar}. \tag{1.3.14}$$

After a Wick rotation $T \rightarrow -iT_E$, the integral is dominated by the Euclidean-time history $x(t_E)$ that solves the Euclidean equations of motion, which are obtained from the ordinary ones by an overall sign of the potential, $V_E = -V$. This classical solution x_{cl} needs to asymptote to $\pm x_0$ for $t_E \rightarrow \pm\infty$; it is called an *instanton* because most of its action comes from a particular time t_0 , when x switches from one vacuum to the other. Evaluating the integrand on this solution gives already a good approximation to the integral; a better one is found by performing the integral over fluctuations $(x - x_{\text{cl}})(t)$ around it. For the double-well V_{dw} , both these steps can be carried out exactly, and the energy of the lowest eigenstate is

$$E_0 \sim \frac{\hbar}{2}\omega - \hbar\omega\sqrt{\frac{6s}{\pi\hbar}}e^{-s/\hbar}, \quad s = \frac{2}{3}\omega x_0^2. \quad (1.3.15)$$

The splitting of the levels contains the nonanalytic $e^{-s/\hbar}$. The coefficient $s = S(x_{\text{cl}})$ is the action evaluated on the classical solution; the coefficient multiplying it comes from the integral over small fluctuations.

In field theory, one again Wick-rotates the path integral, looking for classical Euclidean solutions ϕ_{cl} with finite action $s = S(\phi_{\text{cl}})$, so that their contribution $e^{-s} \neq 0$. So a field theory instanton should be localized in space and time. A famous example occurs for YM theories in $d = 4$, where instantons are solutions of the *self-duality equation*

$$F_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu}{}^{\rho\sigma}F_{\rho\sigma}, \quad (1.3.16)$$

which in the notation of (1.2.34b) reads

$$F = *F. \quad (1.3.17)$$

Computing these effects exactly is more challenging, but sometimes it can be done; again, supersymmetry helps (see, for example, [56] for a review). While the SW $\mathcal{N} = 2$ solution was found with other methods, one can in fact reproduce it exactly by counting gauge instantons [57].

World-sheet instantons in string theory

In string theory, we have two expansion parameters, l_s and $g_s = e^\phi$. For l_s , we can apply the preceding discussion to the two-dimensional QFT on the world-sheet. An instanton for this model is a finite-action solution to the Euclidean equations of motion. Equation (1.1.4) is not directly the world-sheet area, but it is classically equivalent to it, so we can look for maps $x(\sigma): \Sigma \rightarrow M_{10}$, where the area $A(\Sigma)$ is minimized. In flat space, this would give the degenerate situation where Σ is shrunk to a point. But for spacetimes with nontrivial topology, such as compactifications, the internal space may contain a two-dimensional subspace S_2 that cannot be continuously shrunk to a point; so its minimal area $A(\Sigma) \neq 0$, and this embedding is a *world-sheet instanton*, which, recalling (1.1.4) and (1.1.5), contributes

$$e^{-A(\Sigma)/4\pi l_s^2} \quad (1.3.18)$$

to the path integral. Remarkably, these effects can sometimes be computed exactly, as we will see in Section 7.1.

Effects that are nonperturbative in g_s are a different matter altogether. The world-sheet approach is intrinsically perturbative in g_s ; so we need a new ingredient.

1.3.2 Open string definition

A *D-brane* is an additional extended object in string theory; so far we have not encountered it because we limited ourselves to a perturbative approach.

In general relativity, no solitons exist, if defined as fully regular solutions with localized energy and flat asymptotics ([58]; for a recent account, see, for example, [59, IV.8]). In a general gravitational theory, we might be tempted to say that black holes are the analogue of solitons: they have localized energy, and they are stable. They do have a singularity, unlike the solitons we have studied in the previous section. It is natural to think, however, that this singularity signals the breakdown of general relativity, and that in a full theory of quantum gravity it disappears.

Besides black holes, string theory has solutions whose energy is localized along a subspace, and not just a point in space; so they are called *p-branes*, a generalization of the word “membrane,” where *p* denotes the number of space dimensions they span. As solutions of the supergravity equations, they have a singularity at their core; but string theory does provide an alternative, smooth understanding, in terms of *open strings*.

An open string is one whose time slice is an interval rather than a circle. It requires choosing boundary conditions for the world-sheet fields; Dirichlet boundary conditions for some of the $x^M(\sigma)$ correspond to open strings that end on a subspace of spacetime, which is accordingly called a Dirichlet-brane, or more commonly a *D-brane*.

These D-branes are the fundamental definition of the *p-brane* gravity solutions [60]. Consider, for example, a D-brane in flat space. We have both closed and open strings, interacting with one another, both described by a world-sheet model with a flat background metric $g_{MN} = \eta_{MN}$. Suppose we now perform the path integral over the open string degrees of freedom – in QFT jargon, we “integrate them out.” As in QFT, the action for the remaining degrees of freedom, those of closed strings, is now modified: the background metric is distorted to a different metric g_{MN} . So we obtain an effective description, with only closed strings in a curved background metric, which is nothing but the *p-brane* metric, or in other words the “back-reaction” of the D-brane on spacetime. Because of the difficulties with the world-sheet description of RR fields that we mentioned at the end of Section 1.1.2, this identification is a bit difficult to demonstrate explicitly, but several indirect arguments point toward it.

We will now review the definition of D-branes as loci where open strings end, and then their gravitational description.

To describe an open string, we take $\sigma^1 \in [0, \pi]$, rather than a periodic coordinate as in Section 1.1. When we vary the action, we now have to pay attention to boundary terms. For example, let us consider flat space $g_{MN} = \eta_{MN}$ and $h_{\alpha\beta} = \eta_{\alpha\beta}$, and focus on a single coordinate x . Then the variation gives

$$\begin{aligned} \frac{1}{2} \delta \int_{\Sigma} d^2\sigma \partial^\alpha x \partial_\alpha x &= \int d^2\sigma \partial^\alpha x \partial_\alpha \delta x = \int d^2\sigma \left(\partial_\alpha (\delta x \partial^\alpha x) - \delta x \partial^2 x \right) \\ &= \int_{\partial\Sigma} d\sigma^0 \delta x \partial_{\sigma^1} x - \int_{\Sigma} d^2\sigma \delta x \partial^2 x. \end{aligned} \quad (1.3.19)$$

The second term gives us the free equation of motion $\partial^2 x = 0$, familiar from closed strings; the first is new. To set it to zero, we can either set $\partial_{\sigma^1} x = 0$ or $\delta x = 0$. The first

is called *Neumann* (N) boundary condition; the second sets to zero the variation of this world-sheet scalar at the boundary, and is called *Dirichlet* (D) boundary condition.

All this was for a single x ; still remaining in flat space, we can select N or D boundary conditions for each of the x^M independently. Each x^M for which we are choosing, a D boundary condition is fixed, $x^M = x_0^M$, at the boundary $\partial\Sigma$, or in other words at the endpoints of the open string. So if we choose N boundary conditions for $p + 1$ fields x^M (including time), and D for the remaining $9 - p$, we have a theory of open strings that end on a $p + 1$ -dimensional object; by definition, this is called a Dp -brane.

Clearly, the presence of a Dp breaks some of the symmetries of the background; in flat space, a flat Dp breaks the ten-dimensional Poincaré group $ISO(1, 9)$ to a $ISO(1, p + 1) \times SO(9 - p)$. For applications to compactifications, we will want Dp -branes that are completely extended along \mathbb{R}^4 and localized along some of the internal dimensions. This preserves the four-dimensional Poincaré group.

The analysis in (1.3.19) is enough for D -branes in bosonic string theory; for type II superstrings, we also need to give boundary conditions for world-sheet spinors. The procedure (1.3.19) applied to the spinorial action (1.1.30) shows that at the boundary $\partial\Sigma$ the combination $\psi_+ \delta\psi_+ - \psi_- \delta\psi_-$ should vanish; this can be arranged by having

$$(\psi_+ = \pm\psi_-)|_{\partial\Sigma}. \quad (1.3.20)$$

Both for the x^M and ψ^M , the boundary conditions now relate left- and right-movers, and the open string spectrum has only one set of oscillators. Just like in the closed string sector, massless states come from the fermionic oscillators; the sign in (1.3.20) gives two different sectors, which again we call NS and R . After GSO projection, we have the massless spectrum

$$b_{-1/2}^M |0\rangle_{NS}, \quad \mathbf{8}_V; \quad |0, \alpha\rangle_R, \quad \mathbf{8}_S. \quad (1.3.21)$$

If we are considering N boundary conditions, the endpoints are not fixed to lie anywhere, and we have a $D9$ -brane; in this case (1.3.21) is interpreted as a vector field a_M and a gaugino λ_α . If we have D directions $i = p + 1, \dots, 9$, then the components $b_{-1/2}^i |0\rangle_{NS}$ are transverse to the Dp -brane, and behave as scalars under $ISO(1, p + 1)$; the parallel components $b_{-1/2}^a |0\rangle_{NS}$, $a = 0, \dots, p$ still represent a vector field. As for the $|0, \alpha\rangle_R$, they represent a spinor under both the parallel and transverse rotations. So in total we have

$$a^a, \quad x^i, \quad \lambda^\alpha. \quad (1.3.22)$$

Calling the scalars x^i might seem confusing, given that we also called x^M the world-sheet scalars. But we will see shortly that the scalars x^i on a D -brane parameterize its transverse fluctuations, and thus give an embedding of Dp into spacetime.

Just as closed (bosonic) strings can couple to condensates of their massless fields (1.1.9), we can couple open strings to condensates of (1.3.22). For example, the coupling to a^a reads

$$\int_{\partial\Sigma} d\sigma^0 a_a \partial_0 x^a; \quad (1.3.23)$$

recalling (1.1.3), this means that the endpoints behave as charged particles under a . Rewriting this as $-\int_\Sigma d^2\sigma \partial_1 (a_a \partial_0 x^a)$ and adding the total derivative

$0 = \int d^2\sigma \partial_0(a_a \partial_1 x^a)$, this becomes $\int_{\Sigma} d^2\sigma f_{ab} \partial_0 x^a \partial_1 x^b$, where the two-form $f = da$ is the field-strength of the vector a . This has the same form as in (1.1.10); in the parallel directions, the two can be collected together, and the combination $(B + 2\pi l_s^2 f)_{ab}$ appears.

1.3.3 Effective action

So far, we considered a Dp -brane extended along a flat subspace $\mathbb{R}^{p+1} \subset \mathbb{R}^{10}$. We can try to generalize this to arbitrary subspaces in arbitrary background metrics, but we still need to impose conformal invariance. For closed strings, this led to the beta functions (1.1.14), which we reinterpreted as spacetime equations of motion for the effective action (1.1.17); for open strings, it leads to equations of motion for the brane, which again come from an effective action S_{Dp} . The fields appearing in this action are those of the closed string spectrum in Tables 1.1 and 1.2, plus the open string fields (1.3.22). Once again, we only give here the bosonic part:

$$S_{Dp} = \tau_{Dp} \left[- \int d^{p+1}\sigma e^{-\phi} \sqrt{-\det(g|_{Dp} + \mathcal{F})} \pm \sum_k (-1)^k \frac{1}{k!} \int_{Dp} C_{p+1-2k} \wedge \mathcal{F}^k \right] \tag{1.3.24}$$

where

$$\tau_{Dp} = \frac{1}{(2\pi)^p l_s^{p+1}}. \tag{1.3.25}$$

The coordinates σ^a , $a = 0, \dots, p$ parameterize the world-volume¹² Dp . The restriction or pull-back operation $|_{Dp}$ consists in contracting each index M with $\partial_a x^M$, as in (1.1.3) and (1.1.10); the $x^M(\sigma^0, \dots, \sigma^p)$ gives the embedding of Dp into spacetime. In the second term of (1.3.24), the spacetime forms are integrated on the Dp world-volume by first pulling them back; so, for example, we should really write $\int_{Dp} C_p \equiv \int_{Dp} C_p|_{Dp}$. The sign \pm is related to the ‘‘orientation’’ on Dp , namely to the order of the coordinates in the world-volume measure; we will see this in more detail later, and fix this convention more precisely in Chapter 10.

Recall that $f^k = f \wedge \dots \wedge f$; we also defined

$$\mathcal{F}_{ab} \equiv (2\pi l_s^2 f + B|_{Dp})_{ab} = 2\pi l_s^2 f_{ab} + B_{MN} \partial_a x^M \partial_b x^N, \tag{1.3.26}$$

in line with the remark that follows (1.3.23). The gauge transformation for B no longer leaves the action invariant, unless it also acts on a :

$$B \rightarrow B + d\hat{\lambda}_1, \quad a \rightarrow a - 2\pi l_s^2 \hat{\lambda}_1. \tag{1.3.27}$$

For open strings, (1.1.12) has to be modified as $g_s^{2g-2+\#b}$, where $\#b$ is the number of boundaries added to Σ ; the prefactor $e^{-\phi}$ in (1.3.29) indicates then that this action originates from a sphere to which we add a single boundary, which is topologically a disk. This is the simplest open string diagram: an open string that is created and later disappears. The term containing C_p does not contain this prefactor because of the

¹² We will call ‘‘world-volume’’ the subspace swept in spacetime by a D-brane, and keep using ‘‘world-sheet’’ for a fundamental string.

customary RR rescaling noted following (1.2.17). For constant $e^\phi = g_s$ effectively the tension of a D-brane is not really τ_{Dp} , but rather

$$T_{Dp} = \frac{\tau_{Dp}}{g_s}. \quad (1.3.28)$$

So at small g_s , D-branes are heavy, similar to the bound on monopole mass in (1.3.9) in terms of g_{YM} .

The DBI term

If we set to zero all fields except g_{MN} , (1.3.24) reduces to

$$- \tau_{Dp} \int d^{p+1} \sigma e^{-\phi} \sqrt{-\det g_{MN} \partial_a x^M \partial_b x^N}. \quad (1.3.29)$$

This is the natural generalization of the particle action (1.1.2) to an extended object; it measures the volume of the $(p+1)$ -dimensional object Dp in spacetime, relative to the background metric g_{MN} . In the flat background metric $g_{MN} = \eta_{MN}$, a subspace that is itself flat extremizes (1.3.29); a curved Dp will tend to relax to such a flat subspace, or shrink to a point. Without loss of generality, we can locally take the first $p+1$ coordinates of spacetime x^M to coincide with the σ^a ; then the remaining $9-p$ can be identified with the transverse scalars x^i in (1.3.22). If we choose the latter so that $g_{aj} = 0$, then

$$(g|_{Dp})_{ab} = g_{ab} + g_{ij} \partial_a x^i \partial_b x^j. \quad (1.3.30)$$

For small x^i , (1.3.29) contains then the usual kinetic term, and terms with more than two derivatives.

The full first term $\sqrt{-\det(g|_{Dp} + \mathcal{F})}$ is called (*Dirac–Born–Infeld (DBI)* because it is reminiscent of early proposals to improve the short-distance behavior of the electromagnetic field, inspired by the resemblance to the relativistic contraction factor $\gamma^{-1} = \sqrt{1-v^2}$. While this combination might seem odd, we will see that it has many natural properties (including a duality with Pythagoras' theorem in Section 1.4.2). We can expand it for small open string fields. Recall that $\log \det M = \text{Tr} \log(M)$; then for a matrix $M \sim 1 + m$ near the identity, $m \ll 1$:

$$\begin{aligned} \det(1 + m) &= \exp[\text{Tr} \log(1 + m)] = \exp\left[\text{Tr} m - \frac{1}{2} \text{Tr}(m^2) + \dots\right] \\ &= 1 + \text{Tr} m + \frac{1}{2} ((\text{Tr} m)^2 - \text{Tr}(m^2)) + \dots \end{aligned} \quad (1.3.31)$$

Using this and (1.3.30), for $B = 0$:

$$\sqrt{-\det(g|_{Dp} + \mathcal{F})} \sim \sqrt{-\det g_{ab}} \left(1 + \frac{1}{2} g_{ij} \partial_a x^i \partial^a x^j + \frac{1}{2} |2\pi l_s^2 f|^2 \right) + \dots, \quad (1.3.32)$$

where we kept only terms with two derivatives, and indices are raised and contracted with g_{ab} . Taking into account the prefactors in (1.3.24), the YM coupling is given by

$$g_{YM}^2 = (2\pi)^{p-2} l_s^{p-3} g_s. \quad (1.3.33)$$

In particular, it is dimensionless for $p = 3$, as expected.

The Wess–Zumino term

We now turn to a discussion of the term involving C_{p+1-2k} in (1.3.24). This is variously called *Chern–Simons* or *Wess–Zumino* (WZ) term, inspired by the names of two famous actions that don’t involve the metric. When $\mathcal{F} = 0$, the integral on Dp is defined as in (1.2.8), after the pull-back operation described following (1.3.24). Explicitly,

$$\int_{Dp} C_{p+1} = \int C_{M_0\dots M_p} \partial_0 x^{M_0} \dots \partial_p x^{M_p} . \tag{1.3.34}$$

For $k = 1$ this has the form (1.1.3); for $k = 2$, it looks like the coupling (1.1.10) of the fundamental string to B . So the coefficient of (1.3.34) is a charge density; we see from (1.3.24) that it equals the brane tension T_{Dp} .

Recall from Section 1.1.2 that the C_{p+1} exist with $p = \text{even}$ in IIA, and $p = \text{odd}$ in IIB; so there are

$$Dp\text{-branes: } p = \begin{cases} \text{even} & \text{IIA,} \\ \text{odd} & \text{IIB.} \end{cases} \tag{1.3.35}$$

In IIA, the smallest object is a point-like soliton in IIA, the D0-brane. In IIB we also have the possibility of a $D(-1)$ -brane or *D-instanton*, with a D boundary condition for the time coordinate as well, so that it is localized also in time. In the context of compactifications, there are additional brane instanton that wrap the internal directions and are completely localized in the noncompact directions.

A subtlety mentioned following (1.3.24) is that the overall sign of (1.3.34) can change if we embed the world-volume differently: if we flip the sign of one of the σ^a ,

$$x^M(\sigma^0, \sigma^1, \dots, \sigma^p) \rightarrow x^M(\sigma^0, -\sigma^1, \dots, \sigma^p) , \tag{1.3.36}$$

the image of the embedding remains the same, but (1.3.34) changes sign, because $\partial_{\sigma^1} x^M$ appears only a single time in it. A similar sign change happens upon exchanging two coordinates. Since the charge density changes sign, we call this an *anti-Dp-brane*. The difference between a brane and antibrane is conventional.

Effect of world-volume flux

When $\mathcal{F} \neq 0$, a Dp -brane also couples to RR fields C_k with $k < p$. For example, the WZ term for a D2-brane becomes

$$\int_{D2} (C_3 - \mathcal{F} \wedge C_1) . \tag{1.3.37}$$

In (1.3.34), C_1 would couple to a D0; so we interpret the second term in (1.3.37) as the presence of a distribution of D0-branes on the D2. In other words, we have a D2 that also has D0 charge, or a *D2/D0 bound state*.

A final subtlety about the WZ term regards the Romans mass F_0 [61, 62]. Since this flux has no potential, it might seem that no brane couples to it. In fact, for $F_0 \neq 0$ the WZ term in (1.3.24) needs to be modified to

$$\int_{Dp} w_{Zp+1} , \quad dw_{Zp+1} = \sum \frac{1}{k!} (-1)^k F_{p+2-2k} \wedge \mathcal{F}^k . \tag{1.3.38}$$

For $F_0 = 0$, $w_{Zp+1} = \sum_k \frac{1}{k!} (-1)^k C_{p+1-2k} \wedge \mathcal{F}^k$ as in (1.3.24); for $F_0 \neq 0$, we can use $\sum_k F_{p+2-2k} \text{cs}_k$, where cs_k generalizes the Chern–Simons form (1.2.20). For example,

for $p = 0$, taking (1.3.24) literally would give a term $\int C_1 - C_{-1} \wedge \mathcal{F}$; the correct coupling can be obtained by formally integrating the second term by part:

$$\int_{D0} (C_1 - F_0 a). \tag{1.3.39}$$

Magnetic dual potentials

In (1.3.34), RR potentials C_p appear for any p , while in Section 1.2 we only saw $p \leq 4$. We need to extend the definition to higher p by introducing the *magnetic dual* potentials, defined through

$$dC_p - H \wedge C_{p-2} \equiv F_{p+1} \equiv (-1)^{p(p-1)/2} * F_{9-p} \quad (p > 4) \tag{1.3.40}$$

in the notation of (1.2.34b). These new C_p , $p \geq 5$ are the higher-dimensional analogue of the magnetic dual gauge fields we mentioned for the SW solution that follows (1.3.12). For example, a D6-brane couples to C_7 , which when $H = 0$ is defined by $dC_7 = F_8$, or more explicitly

$$F_{M_1 \dots M_8} = -\sqrt{-g} \epsilon_{M_1 \dots M_8}^{(0) M_9 M_{10}} \partial_{M_9} C_{M_{10}}. \tag{1.3.41}$$

In Exercise 1.2.1, it was noted that the RR equations of motion look like an extension of the Bianchi identities in terms of the magnetic duals. This allows to rewrite the action by swapping one or more of the original potentials with their magnetic duals. In presence of a Dp with $p \geq 4$, it is convenient to use this reformulation to vary with respect to C_{p+1} . The combined action, (1.3.24) plus the supergravity action in (1.2.31) or (1.2.36), can be rewritten as an integral over ten dimensions by using delta functions. In the language of forms, recalling our notation x^i for the directions transverse to the Dp , we can introduce a form

$$\delta_{Dp} \equiv \delta(x^1) \dots \delta(x^{9-p}) dx^1 \wedge \dots \wedge dx^{9-p}. \tag{1.3.42}$$

After the variation, we then obtain (for $\mathcal{F} = 0$)

$$dF_p - H \wedge F_{p-2} = \mp 2\kappa^2 \tau_{D(8-p)} \wedge \delta_{D(8-p)}; \tag{1.3.43}$$

the source term on the right-hand side is the effect of D-branes on the RR field strengths. A similar source term appears in the equations of motion for the other closed string fields.

1.3.4 Flux quantization

The coupling (1.3.34) is the natural generalization of (1.1.3) to extended objects. In this sense, a Dp -brane is the elementary charge for the RR field C_{p+1} , much as the electron for the electric field. The generalization of the large gauge transformation (1.2.13) is

$$C_{p+1} \rightarrow C_{p+1} + \Lambda_{p+1}, \quad \frac{\tau_{Dp}}{2\pi} \int_{S_{p+1}} \Lambda_{p+1} \in \mathbb{Z} \tag{1.3.44}$$

for any subspace S_{p+1} . This includes our old “small” RR gauge transformations (1.2.11): $\Lambda_{p+1} = d\lambda_p$ is a total derivative, and the integral (1.3.44) is zero.

The non-single-valuedness of a gauge transformation λ_0 was used in the Dirac quantization argument (1.3.7). This suggests a generalization of that result to RR

fields. Surround a Dp -brane with a sphere S^{8-p} , consider an “equator” E on it (itself a sphere S^{7-p}), and the two semispheres U_N, U_S into which it divides S^{8-p} . The integral $\tau_{D(6-p)} \int_{U_N} (dC)_{8-p}$ can be rewritten using an analogue of Stokes’s theorem (which we will formalize in Section 4.1.10) as

$$\tau_{D(6-p)} \int_E C_{7-p}. \tag{1.3.45}$$

Repeating this over U_S yields a competing result for (1.3.45); the two results in general disagree, their difference being exactly the integral of F_{8-p} over all of S^{8-p} . But there is no inconsistency if we use two different C_{p+1} on U_N and U_S , differing by a large gauge transformation (1.3.44). This leads to the condition

$$\tau_{D(6-p)} \int_{S^k} (dC)_{8-p} = 2\pi n, \tag{1.3.46}$$

where $n \in \mathbb{Z}$. Recalling (1.3.25),

$$\frac{1}{(2\pi l_s)^{k-1}} \int_{S^k} (dC)_k \in \mathbb{Z}. \tag{1.3.47}$$

Although for concreteness we presented our argument with spheres, (1.3.47) holds for any k -dimensional subspace S_k . Often such a subspace can be continuously deformed to a point, and in that case the integral in (1.3.47) just vanishes. The integral can be nonzero in two types of situations:

- If S_k surrounds a brane, we cannot continuously deform it to zero without crossing it. The integral $\int F_{8-p}$ can be taken as the definition of the charge of a Dp ; one could use this to rederive (1.3.25). In this case, flux quantization can also be derived by suitably integrating (1.3.43).
- For compactifications, some S_k in the internal space cannot be continuously deformed to a point for *topological* reasons (Section 4.1.10). In this case, (1.3.47) can be nonzero even if there are no D-branes.

The logic behind (1.3.47) can also be applied to the magnetic dual potentials (1.3.40); one concludes that (1.3.47) is valid for all $k \leq 10$, and not just for the F_k with $k \leq 5$ that appear in the supergravity actions. Notice that the $*$ in (1.3.40) contains the metric, as we see explicitly in (1.3.41). This seems to create a paradox: if we impose flux quantization both for F_3 and for its dual $*F_3$ (say), it seems we are imposing some quantization conditions on the metric itself. In practice, however, there are always some noncompact directions in spacetime, and only one of the two flux integrals makes sense. For example, for $M_{10} = \mathbb{R}^4 \times M_6$, if we impose flux quantization for F_3 , the one for $*F_3$ would involve an integral over \mathbb{R}^4 , which would diverge.

The case $k = 0$ in (1.3.47) deserves a separate discussion. There is no potential C_{-1} , and $S^0 = \{x_1^2 = 1\}$ is just the union of two points. But integrating (1.3.43) for $p = 0$ on a segment crossing a D8, we still obtain

$$2\pi l_s \Delta F_0 = N_{D8} \in \mathbb{Z}. \tag{1.3.48}$$

Strictly speaking, in this case we only quantized the jumps of F_0 rather than F_0 itself. It would now be possible to entertain the notion that $2\pi l_s F_0 = n_0 + \theta$, with $n_0 \in \mathbb{Z}$ and $\theta \in [0, 1)$, fixed for any background. This looks unlikely; we will see that T-duality relates the RR fluxes to one another, and if $\theta \neq 0$ were allowed it would eventually violate one of the (1.3.47) for $k > 0$. So we will take

$$2\pi l_s F_0 \in \mathbb{Z}. \tag{1.3.49}$$

Our discussion can also be easily applied to B , since its coupling to the fundamental string is identical to the one of C_1 to the D1. Looking at (1.3.44) and (1.3.47), we conclude

$$B \rightarrow B + \tilde{\Lambda}_2, \quad \frac{1}{(2\pi l_s)^2} \int_{S^2} \tilde{\Lambda}_2 \in \mathbb{Z}; \tag{1.3.50a}$$

$$\frac{1}{(2\pi l_s)^2} \int H \in \mathbb{Z}. \tag{1.3.50b}$$

1.3.5 Supersymmetry

D-branes break some of the bosonic symmetries of a background; perhaps more importantly, D-branes also partially break some of its supersymmetry. Recall that both type II string theories are invariant under 32 supercharges, whose infinitesimal parameters are two spacetime fermions ϵ^a . A Dp extended along a flat $\mathbb{R}^{p+1} \subset \mathbb{R}^{10}$ is invariant only if the two ϵ^a are related by

$$\epsilon^1 = \Gamma_{\parallel} \epsilon^2, \tag{1.3.51}$$

where

$$\Gamma_{\parallel} = \frac{1}{p! \sqrt{-\det(g_{Dp})}} \epsilon_{(0)}^{a_0 \dots a_p} \Gamma_{a_0 \dots a_p}, \quad \Gamma_a \equiv \Gamma_M \partial_a x^M \quad (\mathcal{F} = 0). \tag{1.3.52}$$

The flat space background is invariant under all 32 supercharges. If we have a Dp -brane extended along a $\mathbb{R}^{p+1} \subset \mathbb{R}^{10}$ subspace, without loss of generality we can take it to be $\{x^{p+1} = \dots = x^9 = 0\}$; (1.3.52) then becomes

$$\Gamma_{\parallel} = \Gamma_0 \dots \Gamma_p. \tag{1.3.53}$$

Equation (1.3.51) gives a relation among the two ϵ^a : only ϵ^1 is now independent, and thus we are left with 16 supercharges. This preserved supersymmetry makes these flat D-branes the analogue of the field-theory BPS states in Section 1.3.1. In particular, they are stable against time evolution and deformations of the theory.¹³ This is the reason they are useful: if we make the string coupling g_s large, we lose perturbative control, but we know that D-branes have to remain in the spectrum. Since their tension (1.3.28) is the inverse of the string coupling, they become light at strong coupling and might become new fundamental objects, similar to the SW solution of $\mathcal{N} = 2$ -supersymmetric Yang–Mills (Section 1.3.1).

We saw in Section 1.1.2 that the ϵ^a have opposite chiralities in IIA, and equal in IIB, and that multiplication by a single Γ^M changes chirality. It follows that

¹³ A D-brane with a more general shape is not guaranteed to preserve supersymmetry, and hence to be stable under time evolution. In Section 9.2, we will consider this further, and also generalize (1.3.52) to $\mathcal{F} \neq 0$.

(1.3.51) can only be solved for $p = \text{even}$ in IIA, and for $p = \text{odd}$ in IIB; this confirms (1.3.35).¹⁴

The origin of the constraint (1.3.51) is roughly the following. The fermionic completion of the effective action (1.3.24) includes two world-sheet fermions θ^a . Closed string supersymmetry acts on these as a spinorial translation:

$$\delta\theta^a = \epsilon^a. \quad (1.3.54)$$

Imposing supersymmetry would then set both $\epsilon^a = 0$. But there is also a gauge equivalence among the θ^a , called κ -symmetry, acting as

$$\delta\theta^1 = \Gamma_{\parallel}\kappa, \quad \delta\theta^2 = \kappa. \quad (1.3.55)$$

Imposing invariance under a combined supersymmetry and κ -symmetry gives (1.3.51).

1.3.6 Multiple D-branes

D-brane stacks and nonabelian gauge groups

To each endpoint of an open string, it is possible to add an extra discrete quantum number $I = 1, \dots, N$, called the *Chan–Paton (CP) label*. It can be interpreted as the presence of $N > 1$ superimposed D-branes, or in other words a stack of N D-branes.

Let us see why. Since open strings have two endpoints, the states (1.3.21) and the fields (1.3.22) acquire two extra labels IJ , and so they are now promoted to Hermitian matrices. Now a becomes a nonabelian $U(N)$ gauge field, and x a scalar in the adjoint representation. In the action (1.3.24), we should turn all fields into matrices with an overall trace, but there are potential ordering ambiguities.

The situation is much clearer for the two-derivative approximation (1.3.32), whose nonabelian extension is dictated by supersymmetry. We know from (1.3.51) that a brane extended along a flat $\mathbb{R}^{p+1} \subset \mathbb{R}^{10}$ subspace preserves 16 supercharges; we expect its effective action to have this invariance. This is the maximum number of supercharges for a QFT model (not including gravity), and models with this property are quite constrained. For example, for $p = 9$, when we have a D9 extended along all of spacetime, there are no scalars x^i , and in (1.3.32) only the $|f|^2$ term remains. Its nonabelian version is the YM Lagrangian density $\text{Tr}|f|^2$; supersymmetry involves also the gaugino λ^α of (1.3.22), and the requirement of 16 supercharges fixes the action uniquely.

For lower p , some components of the gauge field now become x^i , because of their common origin in (1.3.57). The resulting supersymmetric YM theory with 16 supercharges is again uniquely fixed: it is the dimensional reduction of the action for $p = 9$ along the directions to the Dp . The components of the gauge field along the transverse directions become the transverse scalars x_i ; the world-volume field strengths then become

$$F_{ai} = D_a x_i = \partial_a x_i + [A_a, x_i], \quad F_{ij} = [x_i, x_j]. \quad (1.3.56)$$

¹⁴ There also exist non-BPS branes that violate (1.3.35) and are of course not described by the effective action (1.3.24). These branes are unstable, but they become stable after some quotients, including the orientifolds we will introduce in Section 1.4.4 [63].

In particular, the YM term generates a potential for the scalars

$$V \propto \text{Tr} \left([x_i, x_j][x_i, x_j] \right), \quad (1.3.57)$$

which vanishes in the Abelian case. The x_i are diagonalizable because Hermitian; the vacua of (1.3.57) are given by configurations where $[x_i, x_j] = 0$, which means that the x_i are simultaneously diagonalizable. In the generic vacuum where they are all different, the BEH mechanism gives a mass to all fields except the diagonal ones; the action of these massless modes is again the Abelian (1.3.32). The D-brane positions are the eigenvalues λ_i^I . This picture motivates the proposed interpretation of the CP label as N superimposed D-branes.

An additional check of this conclusion is that the off-diagonal modes receive a mass that is proportional to the $\sum_i (\lambda^I - \lambda^J)_i^2$; these can be interpreted as the lengths of the strings going from the I th to the J th D-brane. Another is supersymmetry: the condition for (1.3.51) is the same for parallel Dp , since $\partial_a x^M$ is insensitive to translations $x^M \rightarrow x^M + x_0^M$. So having parallel Dp still preserves 16 supercharges, which is what we assumed for the nonabelian super-YM theory in presence of the CP label.

Brane–antibrane system

Recall from (1.3.36) that parity in one of the embedding coordinates changes the WZ coupling (1.3.34) by a sign; we called this an anti- Dp . Now in the condition for supersymmetry (1.3.51) the matrix Γ_{\parallel} changes sign, too. If we have a Dp and an anti- Dp together, we have to solve

$$\epsilon^1 = \Gamma_{\parallel} \epsilon^2, \quad \epsilon^1 = -\Gamma_{\parallel} \epsilon^2, \quad (1.3.58)$$

which is impossible. So a brane–antibrane system breaks supersymmetry.

Moreover, an analysis of the open string modes reveals the presence of a tachyon. It has been shown [64, 65] that in fact this tachyon has a nontrivial potential, with a stable vacuum, which represents the closed-string vacuum without any branes; in other words, the tachyon can condense (get a nonzero expectation value) and become stable. The spacetime interpretation is that the Dp –anti- Dp pair annihilates, much as a particle–antiparticle pair. The story becomes even more interesting when one of the two has a nontrivial flux of the world-volume field-strength f ; in this case, after the annihilation we are left with a lower-dimensional brane instead of the vacuum. This motivates the *K-theory* interpretation of D-branes [66], also suggested by the higher-derivative corrections to (1.3.24) [67]. (This will reappear in Section 9.2.3.)

1.3.7 Gravity solutions

We anticipated at the beginning of this section that a D-brane back-reacts on closed string fields, distorting the flat space metric to a so-called p -brane solution. These can be found by solving the closed string equations of motion with the symmetries we expect the brane to preserve, namely $\text{ISO}(p+1) \times \text{SO}(d-p)$ (parallel Poincaré and transverse rotations), and 16 supercharges. Here we give the result; we will check in later chapters that they are supersymmetric solutions, in various ways.

The solutions for N superimposed Dp -branes, extended along a flat $\{x^{p+1} = \dots = x^9 = 0\} = \mathbb{R}^{p+1} \subset \mathbb{R}^{10}$, is

$$ds_{Dp}^2 = h^{-1/2} ds_{\parallel}^2 + h^{1/2} ds_{\perp}^2, \quad e^{\phi} = g_s h^{\frac{3-p}{4}}, \quad (1.3.59a)$$

$$F_{i_1 \dots i_{8-p}} = \frac{f_{8-p}}{r^{9-p}} x^i \epsilon_{i i_1 \dots i_{8-p}}^{\perp}, \quad f_{8-p} \equiv \frac{(2\pi l_s)^{7-p} N}{v_{8-p}}. \quad (1.3.59b)$$

$ds_{\parallel}^2 = -(dx^0)^2 + \sum_{a=1}^p (dx^a)^2$, and $ds_{\perp}^2 = \sum_{i=p+1}^9 (dx^i)^2$ are the metrics respectively of the $1 + p$ parallel and $9 - p$ transverse dimensions prior to introducing the D-branes; recall that a and i are our names in this section for the parallel and transverse indices respectively. ϵ^{\perp} is the completely antisymmetric tensor in the transverse directions (such that $\epsilon_{p+1 \dots 9}^{\perp} = 1$); $v_d \equiv \text{Vol}(S^d) = 2 \frac{\pi^{(d+1)/2}}{\Gamma((d+1)/2)}$ is the volume of the unit-radius sphere S^d ; in form notation, as we will see in [Exercise 4.1.9](#), $F = f_p \text{vol}_{S^d}$, the volume form of S^d .

h is a function of the radial direction $r = (\sum_{i=p+1}^9 (x^i)^2)^{1/2}$, which satisfies

$$\sum_{i=p+1}^9 \partial_i^2 h = 0 \quad (1.3.60)$$

away from $r = 0$. For $p < 7$, we can take

$$h = 1 + N \frac{r_0^{7-p}}{r^{7-p}}, \quad r_0^{7-p} \equiv \frac{g_s (2\pi l_s)^{7-p}}{(7-p)v_{8-p}}. \quad (1.3.61)$$

The fields not mentioned in (1.3.59) are zero, and in particular so is $H = dB$.

For the sake of a unified description, in (1.3.59) we have given the transverse RR field-strength F_{8-p} . For $p \geq 4$, this is one of the RR forms that appear in the type II supergravity actions of [Section 1.2](#). For $p \leq 2$, this is one of the magnetic duals defined in (1.3.40); using that definition backward on (1.3.59) gives the original RR form F_{p+2} , which has components $F_{a_0 \dots a_p r}$. For $p = 3$, the flux F_5 is self-dual, and both the transverse and parallel directions appear.

The $x^i \epsilon_{i i_1 \dots i_{8-p}}^{\perp}$ in (1.3.59b) can be interpreted as the angular part of the transverse directions, as we will see better in [Section 4.1.3](#); its integral is simply v_{8-p} , and the prefactor is fixed so that (1.3.46) and (1.3.47) holds with $n = 1$. The equations of motion then determine the coefficient of r^{7-p} in (1.3.61), but only up to a sign; the correct choice can be decided by examining the long-distance behavior of the gravitational field, which should be attractive given that we want positive tension as in (1.3.28).

Generalizations

The solutions (1.3.59) are part of a more general family, where there are two horizons; (1.3.59) is the extremal limit where the two horizons have coalesced, and have been set to $r = 0$ by a coordinate change.¹⁵ This extremality is related to the equality of the tension and the charge density of a D-brane, or in other words the coefficients of the DBI and WZ terms in (1.3.24). It is quite common for the extremal limit to saturate the BPS bound, and vice versa.

¹⁵ See, for example, [68, sec. 1] or [69, chap. 19] for more details; the change of variables is similar to the one we will see in [Section 11.1](#) for charged black holes.

Another generalization, which will be more important for us, consists in considering parallel branes. This can be achieved by solving (1.3.60) with a harmonic function, which has several point-like sources in the transverse space rather than just one. For $p < 7$, (1.3.61) is replaced by

$$h = 1 + r_0^{7-p} \sum_{\alpha=1} \frac{1}{|x - x_I|^{7-p}}, \quad (1.3.62)$$

where x_I^i is the position of the I th D-brane in transverse space. The fact that this solution is static signals that *parallel Dp-branes don't exert any force on each other*. We could also have expected this from the analysis in Section 1.3.6, where we concluded that parallel Dp preserve the same 16 supercharges as a single one.

Asymptotics and singularities

Returning to a single stack (1.3.61), at large r we would expect the solution to be asymptotic to flat space. The radius r_0 is interpreted as the region where the gravitation field is strong. For $g_s \rightarrow 0$, it shrinks to zero; this might seem strange, since the D-brane's tension (1.3.28) gets large in this limit. The reason is the prefactor $e^{-2\phi}$ in the NSNS sector action (1.2.17), so that the effective Newton constant is $2\kappa^2 g_s^2$; this overcomes the g_s^{-1} in the D-brane tension, so that their product goes like g_s , which is the power observed in r_0^{7-p} . (Indeed, $(2\pi l_s)^{7-p} = 2\kappa^2 \tau_{Dp}$.)

In fact, as we will now see, $r = 0$ is in most cases a singularity; either curvature, or dilaton, diverge there. As we mentioned, this singularity is expected to be resolved in fully fledged string theory. For example, in Section 7.2.3 we are going to see how this happens for the D6-brane solution.

The behavior of this solution as $r \rightarrow 0$ depends on p :

- If $p < 3$, the curvature invariants remain finite as $r \rightarrow 0$; in particular, the Ricci scalar $R \rightarrow 0$. But the string coupling diverges. $r = 0$ is at infinite distance.
- If $p = 3$, again the curvature invariants are finite, and moreover ϕ is constant. In fact, an analytic continuation beyond $r = 0$ exists [70], but once again $r = 0$ is at infinite distance.
- If $3 < p < 7$, the curvature diverges as $r \rightarrow 0$, but the string coupling $e^\phi \rightarrow 0$.
- For $p = 7$, there are two transverse directions, and a function satisfying (1.3.60) is a logarithm:

$$h = -\frac{g_s N}{2\pi} \log(r/r_0). \quad (1.3.63)$$

$r = 0$ is again a singularity. The metric is no longer asymptotic to flat space at large distance: for $r > r_0$ the function h becomes negative, and the solution loses meaning. In practice, this is seldom a problem, because in the context of compactifications the transverse directions to a D7 are compact anyway.

- For $p = 8$, there is only one transverse direction x_9 , and a function satisfying (1.3.60) is piecewise-linear:

$$h = h_0 + \frac{g_s N}{2\pi l_s} |x_9|. \quad (1.3.64)$$

Again, the solution is no longer asymptotic to flat space, and has a critical distance, dependent on the integration constant h_0 , which is related to the value of the dilaton at $x_9 = 0$.

NS5-branes

Similar solutions to (1.3.59) exist that are supersymmetric and charged under the Kalb–Ramond field B .

One represents the back-reaction of a stack of superimposed fundamental strings $F1$; we will see it later in an exercise. Another describes a new object, extended along five space dimensions and hence called *NS5-brane*. The solution is rather similar to (1.3.59) for $p = 5$:

$$ds_{\text{NS5}}^2 = ds_{\parallel}^2 + h ds_{\perp}^2, \quad e^{\phi} = g_s h^{1/2}, \quad (1.3.65a)$$

$$H_{ijk} = 2N x^l \epsilon_{lijk}^{\perp}. \quad (1.3.65b)$$

$ds_{\parallel}^2, ds_{\perp}^2$ describe the six parallel and four transverse directions; $h = 1 + Nl_s^2/r^2$. (In form notation, (1.3.65b) reads $H = 2N \text{vol}_{S^3}$.) A definition of this solitonic object from open strings is not available, but we will see later that it plays an important role in string dualities, which give indirect information. One can infer from this an effective action in the style of (1.3.24). In particular, one obtains that it couples to the dual potential B_6 , defined similar to (1.3.40) as

$$*H = -dB_6. \quad (1.3.66)$$

The tension is

$$T_{\text{NS5}} \equiv \frac{\tau_{\text{NS5}}}{g_s^2} \equiv \frac{1}{(2\pi)^5 l_s^6 g_s^2}, \quad (1.3.67)$$

which differs from $T_{D5} = 1/((2\pi)^5 l_s^6 g_s)$ from (1.3.28) in the power of g_s , signaling that the NS5 is not a D-brane. Its effect on the Bianchi identity is

$$dH = -2\kappa^2 \tau_{\text{NS5}} \delta_{\text{NS5}}, \quad (1.3.68)$$

in the spirit of (1.3.43).

Exercise 1.3.1 In flat ten-dimensional space \mathbb{R}^{10} , consider a D-brane on $S^p \times \mathbb{R}$, where $S^p = \{\sum_{m=1}^p (x^m)^2 = R^2\}$. Does it satisfy (1.3.51)?

Exercise 1.3.2 Obtain (1.3.47) from (1.3.43).

Exercise 1.3.3 Show that the metric of the $D8$ solution can be put in the *conformally flat* form

$$ds_{10}^2 = f ds_{\text{Mink}_{10}}^2, \quad (1.3.69)$$

where f is a function and Mink_{10} denotes as usual *flat* Minkowski space.

Exercise 1.3.4 Check the sign of the coefficient of r^{7-p} in (1.3.61) by considering the potential of a p -brane with DBI action (1.3.29) in the solution (1.3.59): the force should be attractive. Next, consider a Dp -brane probe, with its full action (1.3.24); check that the total force in this case is zero.

1.4 Dualities

We have learned that in flat space D-branes extended along flat subspaces are BPS solitons, the analogue of monopoles in supersymmetric YM theories; in this section, we will use them to find information on the strong-coupling behavior of string theory.

1.4.1 From IIA to eleven dimensions

D0-branes as KK-modes

D0-branes are particle states. When $g_s = e^\phi$ is constant, from (1.3.28) we see that they have a mass

$$m_{\text{D0}} = \frac{1}{l_s g_s}. \quad (1.4.1)$$

States made of k coincident D0-branes have masses $k/(l_s g_s)$. They are all multiple of the same fundamental value (1.4.1), just like a tower of KK states (0.2.1). Taking this analogy seriously suggests that a new dimension has been generated dynamically, of size

$$L_{10} = 2\pi l_s g_s. \quad (1.4.2)$$

At weak coupling, (1.2.18) implies $L_{10} \ll l_{\text{Pl}}$: the size of this extra circle is sub-Planckian, and has unclear physical meaning. But at strong coupling $L_{10} \gg l_{\text{Pl}}$: the new dimension is macroscopic, and cannot be ignored. So if the KK interpretation of D0-branes is correct, we should expect the strong-coupling limit of IIA string theory to have an eleven-dimensional interpretation.

Since these states are BPS, supersymmetry should not be broken in the process, and we would expect the eleven-dimensional theory to still have 32 supercharges. Such a theory exists and is unique.

Eleven-dimensional supergravity

The fields are the metric, a three-form potential, and a single gravitino:

$$g_{MN}, \quad A_{MNP}; \quad \psi_{\alpha M}. \quad (1.4.3)$$

The field-strength of the three-form is $G_4 = dA_3$, in the form notation of Section 1.2.1. The bosonic action is [71]

$$S_{11} = \frac{1}{2\kappa_{11}^2} \left[\int d^{11}x \sqrt{-g_{11}} \left(R_{11} - \frac{1}{2} |G_4|^2 \right) - \frac{1}{6} \int A_3 \wedge G_4 \wedge G_4 \right]. \quad (1.4.4)$$

The supersymmetry transformations contain a single Majorana ϵ , which has 32 independent components.

On $\text{Mink}_{10} \times S^1$, if the circle is small we can ignore dependence on it, and describe the theory by ten-dimensional fields. The metric g_{MN}^{11} generates in $d = 10$ a metric, a vector field g_{M10} and a scalar g_{1010} . The latter two can be identified with C_M and the dilaton in IIA. The three-form potential A_{MNP} generates a three-form and two-form A_{MN10} ; these can be identified with C_{MNP} and B_{MN} respectively. To reduce the gravitino, we notice that a spinor in $d = 11$ becomes a pair of spinors of both chiralities in $d = 10$; so $\psi_{M\alpha}$ generates $\psi_{M\alpha}$, $\psi_{M\dot{\alpha}}$, $\psi_{10\alpha}$, $\psi_{10\dot{\alpha}}$, which can be identified with the gravitinos and dilatinos in Table 1.1.

More precisely, with the field identification

$$ds_{11}^2 = e^{-\frac{2}{3}\phi} (ds_{10}^2 + e^{2\phi} (dx^{10} - C_1)^2), \quad (1.4.5a)$$

$$A_3 = C_3 - B \wedge dx^{10}, \quad (1.4.5b)$$

the action (1.4.4) turns into (1.2.15), the IIA action for $F_0 = 0$. The four-form field-strength is

$$G_4 = dC_3 - H \wedge dx^{10} \stackrel{(1.2.9)}{=} F_4 - e^{-2/3\phi} H \wedge e^{10}. \quad (1.4.5c)$$

To identify κ_{11} , notice that in (1.4.4) the integral over dx^{10} gives an overall L_{10} ; this would give $\kappa_{11}^2 = \kappa^2 L_{10}$. For the physical Planck length, however, we also want to include the $e^{-2\phi} = g_s^{-2}$ factor in (1.2.17), as we did in (1.2.18). Defining $2\kappa_{11}^2 = (2\pi)^8 l_{\text{P}11}^9$ by analogy with (1.2.16):

$$l_{\text{P}11} = g_s^{1/3} l_s. \quad (1.4.6)$$

L_{10} is macroscopic with respect to this length again when $g_s \gg 1$.

The dimensional reduction we have just described might look dangerous, since in this regime IIA is strongly coupled, and strictly speaking we don't know its action. However, as we mentioned after (1.2.23), the two-derivative action is protected by supersymmetry, and so for slowly varying fields (1.2.15) is still appropriate.

Still (1.4.4) cannot be the end of the story: it is nonrenormalizable, so it cannot be itself the strong-coupling definition of IIA string theory. Rather, we should think of it as another effective action, useful in the regime where $g_s \gg 1$, just like (1.2.15) is useful in the regime $g_s \ll 1$. Of what theory is it an effective action? It cannot be one of the perturbative string theories we know, since it is defined on an eleven-dimensional spacetime. Moreover, in (1.4.4) there is no coupling parameter that can be made small, like g_s was for perturbative strings. So the chances that we can find a weakly coupled description at this point appear slim. We will proceed anyway, and see what we can learn about it. This mysterious theory in $d = 11$ is called *M-theory*, and we conclude [72]

$$\text{M-theory on } S^1 \cong \text{IIA}. \quad (1.4.7)$$

Membranes

The relation to IIA strings suggests that M-theory should also have extended objects.

All BPS objects are charged under some potential. Dp -branes are charged under C_{p+1} ; the M-theory potential A_3 suggests the existence of a 2-brane, also called a membrane. One can indeed find a gravitational solution charged under A_3 , along the lines of (1.3.59) (Exercise 1.4.6). Perhaps more impressively, one can write a supersymmetric theory [73] for a membrane embedded in spacetime \mathbb{R}^d , *only for* $d = 11$. This is strongly reminiscent of string theory, where the one-dimensional object F1 is embedded in $d = 10$. Unfortunately this theory is difficult to quantize; we will find a better version in Chapter 11. Still we can postulate this object's existence

and see what follows from it. This membrane is called *M2-brane*, and is one of the inspirations for the name “M-theory.”

It is natural to think that the M2 becomes the D2 when the eleventh dimension becomes small, $L_{10} \rightarrow 0$. In particular, the two tensions should be the same when viewed by a ten-dimensional observer:

$$T_{\text{M2}} = T_{\text{D2}} \stackrel{(1.3.28)}{=} \frac{1}{4\pi^2 l_s^3 g_s} \stackrel{(1.4.6)}{=} \frac{1}{4\pi^2 l_{\text{P11}}^3}. \quad (1.4.8)$$

We can also consider a D2 extended along x^{10} . In the limit $L_{10} \rightarrow 0$, in $d = 10$ this will look like a $p = 1$ -brane. The only such object is the fundamental string F1. So we are led to also postulate

$$L_{10} T_{\text{M2}} = T_{\text{F1}} \stackrel{(1.1.5)}{=} \frac{1}{2\pi l_s^2}. \quad (1.4.9)$$

Indeed, this follows from (1.4.2) and (1.4.8); we can regard this as an alternative derivation for (1.4.2).

When several M2-branes are brought together, one expects a nonabelian theory to arise, since this happens for D2-branes. As we noted earlier, for M-theory we don't have direct control over the membrane dynamics, so we cannot just derive such a theory directly. Moreover, in IIA the YM parameter on the D-brane world-volume is (1.3.33), and we could make it small by changing g_s ; in M-theory we don't have a similar parameter. In spite of these difficulties, the non-abelian M2 theory was found [74–77]. The trick was essentially to consider M2-branes on a geometry obtained as an *orbifold*, namely by identifying some space directions by the action of a discrete group \mathbb{Z}_k ; the geometry is still flat, but the M2 dynamics now contains factors of $1/k$ that provide perturbative control. We will comment on this further in Section 11.3.

M5-branes

D-branes can couple to the magnetic dual potentials defined in (1.3.40). In the same spirit we can define a potential A_6 by

$$*G_4 = dA_6. \quad (1.4.10)$$

A brane coupling to it would be extended along $p = 5$ space dimensions; hence we call it *M5-brane*.

Its properties can again be inferred from its dimensional reductions. It is expected to have a two-form potential b_2 on its world-volume, whose IIA reduction becomes the gauge field a . To avoid it from also generating an unwanted two-form, one imposes a duality condition $h_3 = *h_3$, $h_3 = db_2$. As for IIB supergravity, this creates a problem in writing an action, although a proposal for overcoming it is known [78]. This world-volume theory has $\mathcal{N} = (2, 0)$ in $d = 6$. For coincident M5-branes, the world-volume theory is expected to have $\sim N^3$ degrees of freedom, rather than the familiar gauge theory $\sim N^2$. This can be inferred from anomalies [79], holography [80], field theory [81–83], and conformal bootstrap [84]. This theory remains mysterious to this day; for recent efforts, see, for example, [85, 86].

We can reduce an M5 to IIA in two ways, by wrapping it along the S^1 or not; this produces branes with $p = 4$ and $p = 5$. Given the list of available BPS objects in IIA, these must be the D4 and the NS5 respectively. This leads to

$$\begin{aligned}
 T_{M5} &= T_{NS5} \stackrel{(1.3.67)}{=} \frac{1}{(2\pi)^5 l_s^6 g_s^2} \stackrel{(1.4.6)}{=} \frac{1}{(2\pi)^5 l_{P11}^6}, \\
 L_{10} T_{M5} &= T_{D4} \stackrel{(1.3.28)}{=} \frac{1}{(2\pi)^4 l_s^5 g_s}.
 \end{aligned}
 \tag{1.4.11}$$

Again the two are in agreement once we recall (1.4.2). We could also regard the first relation as a derivation of T_{NS5} .

There are still some objects in IIA that we have not recovered from M-theory: D6- and D8-branes. D6-branes are a source for the flux F_2 ; we see from (1.4.5a) that its potential C_1 becomes part of the metric in eleven dimensions. So we expect a D6-brane to become a purely geometrical object in $d = 11$; we will see how this comes about in Section 9.2.

Massive IIA

D8-branes are more troubling. They are a source for the Romans mass F_0 ; but the dimensional reduction (1.4.5) of eleven-dimensional supergravity only reproduced the $F_0 = 0$ version (1.2.15) of IIA. Even our original argument for the emergence of an eleventh dimension no longer works with $F_0 \neq 0$, because D0-branes have a term $\int F_0 a$ from (1.3.39), and the equation of motion on the $0 + 1$ -dimensional world-sheet gives $F_0 = 0$, a contradiction. This is known as a ‘‘tadpole’’ problem because the field a has a nonzero one-point function.

The nonperturbative definition of massive IIA cannot be conventional M-theory; it is an interesting open question. We will see in Section 10.1 that there are no classical supergravity solutions with $F_0 \neq 0$ and $g_s \gg 1$ [87]; but from a quantum-mechanical point of view, the path integral should be over all configurations, so the nonperturbative formulation is still needed. Applying the formulas (1.4.5) to the D8-brane solution, one obtains a metric in eleven dimensions [88], but this does not clarify what the action is. There are in fact arguments against the existence of deformations of eleven-dimensional supergravity that preserve 32 supercharges [89–91]. More recently, it has been suggested that these can be overcome by a ‘‘dual’’ formulation for the graviton [92].

This might make us doubt that massive IIA is really part of string theory. There are several reasons to believe that it is. Anticipating a bit some later discussions:

- We will see in Section 1.4.2 that *T-duality* relates F_0 to the other RR fluxes, and the D8 to the other Dp -branes.
- Related to this, one can get to M-theory from massive IIA with a *sequence* of dualities. At the end of Section 1.4.5, a particular quotient of massive IIA will be related to heterotic string theories, and to M-theory on spaces with boundaries. There are other related duality sequences [93, 94].
- Several AdS solutions exist with $F_0 \neq 0$, some of which we will see in Chapter 11. These can be tested quantitatively using the AdS/CFT correspondence [87, 95–97].

1.4.2 T-duality

T-duality can be obtained from the world-sheet; we could have discussed it already in Section 1.1. But it can also be interpreted in terms of KK states, similar to our motivation for eleven-dimensional supergravity.

T-duality can be used when at least one of the directions is an S^1 . For definiteness and simplicity, in this section we consider a spacetime

$$\mathbb{R}^{10-k} \times (S^1)^k, \quad (1.4.12)$$

with a line element $ds_{10}^2 = ds_{\mathbb{R}^{10-k}}^2 + \sum_{M=k+1}^9 (dx^M)^2$. The only data of the torus are the periodicities L_k of the S^1 s. In Section 4.2.6 we will deal with more complicated situations, which we will cast in the language of fiber bundles.

We first consider $k = 1$, so that there is a single periodic coordinate $x^9 \sim x^9 + L$. In this case, T-duality is the equivalence $L \cong \tilde{L}$,

$$\tilde{L} = \frac{4\pi^2 l_s^2}{L}. \quad (1.4.13)$$

For superstrings, the duality also exchanges IIA and IIB.

KK argument

We first show (1.4.13) in the bosonic string. A first piece of evidence comes from the formula for the spectrum: (1.1.23) is invariant under

$$R \rightarrow \tilde{R} = \frac{l_s^2}{R}, \quad w \leftrightarrow q, \quad (1.4.14)$$

matching (1.4.13) for $L = 2\pi R$, $\tilde{L} = 2\pi \tilde{R}$. We did not derive (1.1.23), so here is a perhaps more intuitive version of the same statement, modeled on the D0 argument leading to (1.4.2). Consider a state with a string winding $w \in \mathbb{Z}$ times around the S^1 . From the nine-dimensional point of view, it looks like a particle, with mass

$$L T_{F1} = \frac{Rw}{l_s^2}. \quad (1.4.15)$$

Comparison with (0.2.1) suggests that such states form a KK tower for an extra S^1 dimension of size $\tilde{L} = 2\pi \tilde{R} = 2\pi l_s^2/R$, reproducing (1.4.14). This version of the argument also applies directly to superstrings.

Dualization of world-sheet scalars

We now show that the world-sheet action is invariant under (1.4.13) (see, for example, [98, 99]). We focus on the periodic scalar $x \equiv x^9$, ignoring the other directions; for simplicity, we also set $B = 0$ and the dilaton to a constant. The relevant part of (1.1.4) is then

$$S_x = -\frac{1}{4\pi l_s^2} \int d^2\sigma \partial_\alpha x \partial^\alpha x. \quad (1.4.16)$$

Introduce a vector field b_α and consider the alternative action

$$S_{b,x} = -\frac{1}{4\pi l_s^2} \int d^2\sigma (b_\alpha b^\alpha - 2\epsilon^{\alpha\beta} \partial_\alpha x \partial_\beta x). \quad (1.4.17)$$

Since b_α has no kinetic term, we can substitute it with the solution of its equation of motion, $b_\alpha = \epsilon_{\alpha\beta} \partial^\beta x$; then $S_{b,x}$ reduces to (1.4.16). On the other hand, the x equations of motion are

$$\epsilon^{\alpha\beta} \partial_\alpha b_\beta = 0. \quad (1.4.18)$$

Naively, the solution is simply $b_\alpha = \partial_\alpha \tilde{x}$ for a new scalar \tilde{x} . Substituting this in (1.4.17) gives an action identical to (1.4.16), but with $x \rightarrow \tilde{x}$.

This does not seem very interesting: the action for a free scalar is equivalent to that of another free scalar. Our real interest is how the periodicities are mapped. Let us Wick-rotate σ^0 and consider the Euclidean model for (1.4.17). For definiteness, we take $\sigma^i \sim \sigma^i + 2\pi$. Since $x \sim x + 2\pi R$, we can parameterize

$$\partial_\alpha x = \partial_\alpha x_{\text{per}} + 2\pi n_\alpha R, \tag{1.4.19}$$

where x_{per} is periodic. $\partial_\alpha x$ is only locally the gradient of a function: its integral along the σ^i can be nonzero. Likewise, the solution to (1.4.18) is not just $b_\alpha = \partial_\alpha \tilde{x}$, but more generally

$$b_\alpha = \partial_\alpha \tilde{x}_{\text{per}} + 2\pi \lambda_\alpha, \tag{1.4.20}$$

where λ_α are constant. If we replace this and (1.4.19) in (1.4.17), we get a term $-\frac{2\pi}{l_s^2} \epsilon^{\alpha\beta} n_\alpha R \lambda_\beta$. The n_α are part of the degrees of freedom of x ; since they are discrete we cannot vary them, but in the path integral we should still sum over them. But now $\sum_m e^{ixm} = 2\pi \delta(x - 2\pi m)$, the Fourier series analogue of the Fourier transform of the delta function; this implies $\lambda_\alpha = (l_s^2/R) m_\alpha$ with $m_\alpha \in \mathbb{Z}$. Comparing (1.4.20) with (1.4.19), the periodicity of \tilde{x} is determined as $2\pi \tilde{R}$, with $\tilde{R} = l_s^2/R$. This reproduces (1.4.14).

Returning to Lorentzian signature, the equation of motion (1.4.18) is $\partial_\alpha \tilde{x} = \epsilon_\alpha^\beta \partial_\beta x$; so

$$\partial_\pm \tilde{x}^9 = \pm \partial_\pm x^9, \tag{1.4.21}$$

while of course the x^M with $M \neq 9$ are unchanged. In other words, T-duality is equivalent to a parity transformation acting on the right-movers only. (In the bosonic string, this can also be seen from the spectrum (1.1.23).) As familiar from QFT (and reviewed in Section 2.1.1), a parity acts on spinors by a gamma matrix. For type II superstrings, this changes the chirality of the GSO projection in such a way that IIA is mapped to IIB. To summarize:

$$\text{IIA on } S^1 \cong \text{IIB on } \tilde{S}^1. \tag{1.4.22}$$

Action on D-branes

Another consequence of (1.4.21) is that N and D boundary conditions are exchanged. Indeed, the N condition is $\partial_{\sigma^1} x^9 = 0$, which becomes $\partial_{\sigma^0} \tilde{x}^9 = 0$; this fixes \tilde{x}^9 to be a constant, so it is the D condition. More precisely, since the a_a and x^i have a common origin from the 8_V in (1.3.21), the component a_9 is exchanged by T-duality with the scalar x^9 . By (1.2.14) and (1.3.23), $a_9 \sim a_9 + 2\pi/l_9$. We conclude

$$Dp \text{ along } x^9 \text{ with } a_9 = a \xleftrightarrow{T_9} D(p-1) \text{ at } \tilde{x}^9 = 2\pi l_s^2 a. \tag{1.4.23}$$

We know from (1.3.59) that a Dp sources an RR field along the angular directions surrounding it. Since in (1.4.23) a Dp loses a parallel direction, the RR flux gains an index:

$$F_{m_1 \dots m_k} \xleftrightarrow{T_9} F_{m_1 \dots m_k 9}. \tag{1.4.24}$$

We are going to give a more complete description of the T-duality’s action on fields in Section 4.2.6.

Let us compare D-brane tensions on both sides of (1.4.23), in the spirit of (1.4.15). From a nine-dimensional point of view, both objects in (1.4.23) look $(p - 1)$ -dimensional. T-duality states that physics should be the same on both sides, so the tensions should be equal:

$$L_9 T_{Dp} = T_{D(p-1)} \Rightarrow \frac{L_9}{(2\pi)^p g_s l_s^{p+1}} = \frac{1}{(2\pi)^{p-1} \tilde{g}_s l_s^p}; \tag{1.4.25}$$

so the value of g_s after T-duality is

$$\tilde{g}_s = g_s \frac{2\pi l_s}{L_9}. \tag{1.4.26}$$

Thus weak coupling is mapped to weak coupling, unlike the $SL(2, \mathbb{R})$ of Section 1.2.3. This is consistent with our use on both sides of the world-sheet approach, which is perturbative in g_s .

Tori and bound states

In presence of a torus (1.4.12), we can choose to T-dualize along any subset of the k compact directions. If $k = 2$, T-duality along a single direction exchanges IIA and IIB, so T-duality T_{89} along both maps IIA to IIA, and IIB to IIB. The action on D-branes is also a straightforward iteration of the rule (1.4.23). For example, a D2 extended along both x^8 and x^9 , or in other words “wrapping” the torus $T^2 = S^1 \times S^1$, is mapped by a single T-duality T_9 to a D1, and by T_{89} to a D0. In the top part of Figure 1.1, the T^2 is represented as a rectangle; the opposite sides are meant to be identified. The D1-branes along x^9 are mapped by T_9 to D0-branes; and a D1 along x^8 is mapped to a D2. But what is the T-dual of an *oblique* D1-brane, which winds w_8 times along S^1_8 and w_9 times along S^1_9 , such as the one on the bottom-left of Figure 1.1?




Figure 1.1 T-duality acting on D1-branes on a T^2 . It maps horizontal and vertical D1-branes to D0- and D2-branes respectively; it maps oblique D1s to D2/D0 bound states.

Such a D1 is still straight: so it should be BPS. From the point of view of an observer in the noncompact eight dimensions, it is a particle with mass

$$m_{\text{obD1}} = T_{\text{D1}} \text{Length}(\text{obD1}) = \frac{1}{2\pi l_s^2 g_s} \sqrt{w_8^2 L_8^2 + w_9^2 L_9^2}. \tag{1.4.27}$$

We may imagine a process where w_8 D1-branes wrapping S_8^1 and w_9 D1s wrapping S_9^1 snap together to create a single oblique D1. In Figure 1.1, this process would connect the top-left and bottom-left situations. In this process, the total mass decreases: because of the triangular inequality, (1.4.27) is smaller than $T_{\text{D1}} \times (w_8 L_8 + w_9 L_9)$. In a sense, the oblique D1 is a bound state of the horizontal and vertical ones. Now, the latter are mapped under T_9 to w_8 D2s and w_9 D0s respectively; so we expect the oblique D1 to be mapped to a bound state of these objects on the IIA side.

This is confirmed by the duality between world-volume gauge fields and transverse scalars. To avoid having to deal with nonabelian branes, we take $w_8 = 1$. We describe an oblique D1 by $x^9 = \frac{w_9 L_9}{L_8} x^8$, so (1.4.23) gives

$$a_9 = \frac{1}{2\pi l_s^2} \frac{w_9 L_9}{L_8} x^8. \tag{1.4.28}$$

As a cross-check, $\frac{1}{2\pi} \int_{\text{D2}} f_{89} dx^8 dx^9 = w_9 \in \mathbb{Z}$, in agreement with Dirac quantization. Now recall from the discussion in (1.3.37) that the presence of f turns the D2 into a bound state of a D2 and w_9 D0s, confirming our expectations.

This D2 is viewed as a particle in eight dimensions, but to compute its mass we cannot just use the formula $T_{\text{D2}} \times L_8 \tilde{L}_9$; because of the presence of $f \neq 0$, we have to use the DBI term. The potential energy is obtained by integrating the Lagrangian density over the space dimensions; we need to evaluate the determinant of the 2×2 matrix $\begin{pmatrix} 1 & \mathcal{F}_{89} \\ -\mathcal{F}_{89} & 1 \end{pmatrix}$. Recalling also (1.3.26) with $B = 0$:

$$\begin{aligned} m_{\text{D2/D0}} &= T_{\text{D2}} L_8 \tilde{L}_9 \sqrt{1 + (2\pi l_s^2 f_{89})^2} \stackrel{(1.4.28)}{=} \frac{L_8 \tilde{L}_9}{(2\pi)^2 l_s^3 \tilde{g}_s} \sqrt{1 + w_9^2 \frac{L_9^2}{L_8^2}} \\ &\stackrel{(1.4.13), (1.4.26)}{=} \frac{1}{2\pi l_s^2 g_s} \sqrt{L_8^2 + w_9^2 L_9^2} \stackrel{(1.4.27)}{=} m_{\text{obD1}}. \end{aligned} \tag{1.4.29}$$

So under T-duality the square root appearing in (1.4.27) because of Pythagoras' theorem is exchanged with the square root in the DBI term of (1.3.24).

1.4.3 S-duality in IIB

In Section 1.4.1, we found a nonperturbative completion to IIA string theory by focusing on the behavior of D0-branes at strong coupling. In IIB, we don't have (BPS) D0-branes; the next "smallest" object is a D1-brane.

The D1 tension $T_{\text{D1}} = \frac{1}{2\pi l_s^2 g_s}$ becomes small at strong coupling, suggesting that D1s might become fundamental objects. We have described all known perturbative string theories in Section 1.1; could this be a new one? The fields are (1.3.22); the world-volume a field can be integrated out, because gauge fields in two dimensions are nondynamical. One finds that the D1 action becomes at strong coupling simply the IIB perturbative string again, in the Green–Schwarz formulation. So the strong-coupling limit of IIB is IIB, but with $F1 \leftrightarrow D1$.

The $SL(2, \mathbb{R})$ symmetry (1.2.38) can relate the weak and strong coupling regimes of IIB supergravity; but in Section 1.2.3 we gave arguments against it being a symmetry of IIB strings. One argument was that it would have required a continuum of string states, coupling to any linear combination $cC_2 + dB$. Now we know at least one such a string state: the D1, which couples to C_2 . So it is possible that at least the element

$$s = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad (1.4.30)$$

is a symmetry of the full IIB string theory: it exchanges B and C_2 , and hence F1s and D1s. This is called *S-duality*.

D-brane duals

As usual, let us check how the various extended objects transform, setting for simplicity $C_0 = 0$. From (1.2.38) and (1.4.30):

$$\phi \rightarrow -\phi, \quad g_{MN} \rightarrow e^{-\phi} g_{MN}, \quad F_3 \leftrightarrow -H, \quad F_5 \rightarrow F_5. \quad (1.4.31)$$

Because of how the metric transforms, a p -brane's tension picks up a factor

$$g_s^{-(p+1)/2}. \quad (1.4.32)$$

So, for example,

$$T_{F1} = \frac{1}{2\pi l_s^2} \rightarrow \frac{1}{2\pi l_s^2 g_s} = T_{D1}. \quad (1.4.33)$$

On the other hand, T_{D1} picks up a g_s^{-1} , but at the same time the g_s in the denominator gets inverted; so the two factors cancel out and we obtain T_{F1} . This confirms that (1.4.30) indeed exchanges F1s and D1s.

We next transform the D3:

$$T_{D3} = \frac{1}{(2\pi)^3 l_s^4 g_s} \xrightarrow{(1.4.32)} \frac{g_s^{-2}}{(2\pi)^3 l_s^4 g_s^{-1}} = T_{D3}. \quad (1.4.34)$$

So the D3-brane should be invariant under (1.4.30), consistent with the invariance of F_5 in (1.4.31), of which it is a source. Next we have

$$T_{D5} = \frac{1}{(2\pi)^5 l_s^6 g_s} \xrightarrow{(1.4.32)} \frac{g_s^{-3}}{(2\pi)^3 l_s^4 g_s^{-1}} \stackrel{(1.3.67)}{=} T_{NS5}. \quad (1.4.35)$$

Again this is consistent with the transformation in (1.4.31) of the fields they create, B and C_2 .

The next object would be the D7-brane. The field it creates is C_0 , which we have chosen to set to zero. More importantly, we have seen after (1.3.63) that its back-reaction on the metric is not localized, but grows logarithmically; so it is not really a soliton. We will discuss its S-dual in Section 9.4. Finally, we would have the D9-brane, which fills all of spacetime. In fact, a D9 by itself is inconsistent. We discussed its world-volume theory preceding (1.3.57), where we argued it to be a supersymmetric YM with 16 supercharges. This theory is chiral, and suffers from a gauge anomaly. This can be cured by introducing more ingredients, as we will do in Section 1.4.4. It is still possible, but subtler, to define an S-dual to the D9, dubbed NS9 [100, sec. 6].

Other S-dualities

We now would like to know whether there are other elements of $SL(2, \mathbb{R})$ besides (1.4.30) that survive in full string theory. A natural candidate is

$$t = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}. \quad (1.4.36)$$

Since $|c\tau + d| = 1$, the metric does not transform under (1.2.38). Moreover $\tau \rightarrow t \cdot \tau = \tau + 1$; taking real and imaginary parts, $\phi \rightarrow \phi$, and

$$C_0 \rightarrow C_0 + 1. \quad (1.4.37)$$

By (1.3.44), this is a large gauge transformation, suggesting that t is indeed a symmetry of string theory.

Equations (1.2.38) and (1.4.36) also give $B \rightarrow B$, $C_2 \rightarrow C_2 + B$. The latter implies that the D1 turns into an object that couples to both C_2 and B : a *D1/F1 bound state*. As a cross-check, consider placing a D1 and an F1 next to each other, both wrapping an S^1 [101]. The F1 can break in two open strings that end on the D1. Recall from (1.3.23) that the endpoints of an open string behave on a D-brane as electric charges for the world-sheet potential a . Now the endpoints of the F1 can recombine, making the open F1 disappear altogether. This leaves behind a flux $f_{01} \neq 0$ on the D1, constant because of the Maxwell equation. Equation (1.3.32) contains a term $|\mathcal{F}|^2$, which in turn contains $-2\pi l_s^2 f_{01} B_{01}|_{D1}$. This is a coupling to B ; so we have indeed obtained a D1/F1 bound state.

Since s and t , (1.4.30), (1.4.36) are both symmetries, so are all the elements of the group they generate:

$$SL(2, \mathbb{Z}). \quad (1.4.38)$$

This is called *S-duality group* to distinguish it from the particular element (1.4.30).

Acting on D1s and F1s, (1.4.38) generates more general bound states of p F1s and q D1s called (p, q) -strings. More precisely, it acts on $\begin{pmatrix} p \\ q \end{pmatrix}$ in the vector representation, with an F1 represented by the vector $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, and a D1 by $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Since F_5 is invariant under (1.2.38), D3-branes are invariant under the whole $SL(2, \mathbb{Z})$. As we discussed around (1.3.57), the field theory living on a D3 stack has 16 supercharges; for $p = 3$, this is $\mathcal{N} = 4$ -supersymmetric YM in $d = 4$. Equation (1.4.38) implies a QFT duality that maps weak to strong coupling, because $g_{\text{YM}}^2 = 2\pi g_s$, by (1.3.33). This was indeed discovered long before it resurfaced in string theory [102]:

$$\tau_{\text{YM}} \rightarrow \frac{a\tau_{\text{YM}} + b}{c\tau_{\text{YM}} + d}, \quad (1.4.39)$$

with τ_{YM} as in (1.3.12); this is a strong check of S-duality. (The θ angle is created by the C_0 contribution to the WZ term, similar to the discussion leading to (1.3.37).)

Next we consider D5- and NS5-branes. They couple to the magnetic potentials C_6 and B_6 respectively, which were defined in (1.3.40) and (1.3.66) from F_3 and H_3 . This implies the existence of bound states of D5s and NS5s, called (p, q) -five-branes. We expect (p, q) -seven-branes for similar reasons, but we postpone their discussion to Section 9.4.

From M-theory to IIB

Consider now M-theory on $\mathbb{R}^9 \times T^2$ [103, 104]; let the two sides of the T^2 have lengths L_A and L_{10} . From Section 1.4.1, this is equivalent to IIA on $\mathbb{R}^9 \times S^1$, with $g_s \equiv g_A = L_{10}/2\pi l_s$. Further T-dualizing along the remaining S^1 gives IIB on the same space; from (1.4.13) and (1.4.26), the length of the S^1 and the string coupling are

$$L_B = \frac{4\pi^2 l_s^2}{L_A}, \quad g_B = g_A \frac{2\pi l_s}{L_A} = \frac{L_{10}}{L_A}. \tag{1.4.40}$$

There is also an equivalent but more direct logic taking M-theory on $\mathbb{R}^9 \times T^2$ to IIB. Naively, shrinking the T^2 results in a nine-dimensional spacetime. However, M2s wrapping the T^2 w times give particle states of mass

$$T_{M2} \text{Area}(M2) = \frac{wL_A L_{10}}{(2\pi)^2 l_{P11}^3}. \tag{1.4.41}$$

Interpreting this once again as a KK tower (0.2.1) gives

$$\frac{L_A L_{10}}{(2\pi)^2 l_{P11}^3} = \frac{2\pi}{L_B}, \tag{1.4.42}$$

suggesting the emergence of a new S^1 of length L_B , that becomes large as the T^2 gets small; so spacetime is ten-dimensional after all. Wrapping the M2s along the two circles of the torus also gives rise to two types of string states; so the ten-dimensional theory is IIB, which has the F1 and D1. Comparing the tensions:

$$L_A T_{M2} = T_{D1} \Rightarrow \frac{L_A}{(2\pi)^2 l_{P11}^3} = \frac{1}{2\pi l_s^2 g_B}, \tag{1.4.43a}$$

$$L_{10} T_{M2} = T_{F1} \Rightarrow \frac{L_{10}}{(2\pi)^2 l_{P11}^3} = \frac{1}{2\pi l_s^2}. \tag{1.4.43b}$$

Combining (1.4.42) with (1.4.43), we recover (1.4.40). We can generalize (1.4.43) by wrapping one direction of the M2 along an oblique direction (k, l) direction, winding p times along the first circle and q along the second (as in the bottom-left panel of Figure 1.1). This gives the (p, q) -strings, the bound states of p D1s and q F1s.

It was a bit arbitrary to decide in (1.4.43) that wrapping the M2s on the first circle gives a D1, and on the second an F1. Had we done the opposite, we would have obtained (1.4.40) with $L_{10} \leftrightarrow L_A$, and hence with $g_B \rightarrow 1/g_B$. So exchanging the role of the two directions of the torus in M-theory gives rise to the S-duality (1.4.30) in IIB. To see what generates the other elements, we consider a more general geometry for the torus. We define the two-dimensional torus as a quotient

$$T^2 \equiv \mathbb{C}/\mathbb{Z}^2, \tag{1.4.44}$$

much as we did in (1.1.40) when we compactified 16 of the bosonic coordinates to define the heterotic string. A fundamental region for this equivalence relation is now a parallelogram, with the two opposite sides identified. The metric on $\mathbb{C} \cong \mathbb{R}^2$ is taken to be the Euclidean one. Up to rescalings and rotations, we can always make one of the generators of the \mathbb{Z}^2 lattice to be 1; but the second is then an arbitrary complex number τ , called *modular parameter*. So the two identifications in (1.4.44) are $z \sim z+1 \sim z+\tau$. All this is illustrated in Figure 1.2. The tori we have discussed so far, in particular (1.4.12) for $k = 2$, were “rectangular”: their τ was purely imaginary.




Figure 1.2 Fundamental region for the identification (5.3.64) defining a torus. The sides marked by the same symbol are to be considered as identified.




Figure 1.3 M-theory on $\mathbb{R}^9 \times T^2$.

Sometimes, different τ correspond to the same lattice: for example, τ and $\tau + 1$. More generally, any Möbius transformation $\tau' = \frac{a\tau+b}{c\tau+d}$, with $a, b, c, d \in \mathbb{Z}$, also defines the same torus as τ . This is how the $SL(2, \mathbb{Z})$ symmetry of IIB manifests itself in M-theory. An M2 wrapping one of the two sides of the fundamental region in Figure 1.2 is now mapped under a Möbius transformation to one wrapping an oblique line, which we argued corresponds to a (p, q) -string.

We summarize the relation among M-theory, IIA, and IIB with the parameter space in Figure 1.3. When both radii are large, spacetime is eleven-dimensional, and the most appropriate description is the mysterious M-theory. When one of the radii is small and the other is large, there are ten macroscopic dimensions; the theory becomes IIA on $\mathbb{R}^9 \times S^1$. When both radii are small, spacetime naively becomes nine-dimensional, but in fact there is a dual IIB description where it is ten-dimensional.

1.4.4 Orientifolds

It is common to define a space by means of an equivalence relation, as for example in the definition (1.4.44) of the torus. In string theory, one can also quotient by symmetries that act on the world-sheet.

A time slice of the world-sheet is one-dimensional; the only possible nontrivial action on it is world-sheet parity $\Omega: \sigma^1 \rightarrow -\sigma^1$. This squares to the identity, so the action R on spacetime should be an *involution*, a map $x^M \rightarrow R^M(x)$ such that $R^2 = 1$. We define an *orientifold* to be the quotient of a string theory by a simultaneous action of Ω and such a spacetime involution [105–107]. (For more thorough discussions, see [108, 109].)

R acts on fermions as a product of the gamma matrices in the transverse directions, as we mentioned after (1.4.21) and will review in Section 2.1.1. This sometimes squares to -1 , and one then has to include a factor $(-1)^{F_L}$, defined as -1 on fields that are fermionic in the left-mover Hilbert space, or in other words on the RNS and RR sectors. If R_p is locally the reflection of $9-p$ coordinates, the orientifold action is

$$\begin{aligned} \Omega R_p & \quad \text{if } p = 0, 1, 4, 5, 8, 9; \\ \Omega(-1)^{F_L} R_p & \quad \text{if } p = 2, 3, 6, 7. \end{aligned} \quad (1.4.45)$$

The action of Ω in the NSNS sector states (1.1.8) is obtained by noting that exchanging left- and right-movers is equivalent to exchanging the two indices MN . So g_{MN} and ϕ remain invariant, while B_{MN} picks up a sign. The Ω action on the RR sector can be inferred likewise. The overall conclusion is that the bosonic fields should satisfy

$$R_p^* \phi = \phi, \quad R_p^* g = g, \quad R_p^* B = -B, \quad R_p^* F_k = -(-1)^{\lfloor p/2 \rfloor + \lfloor k/2 \rfloor} F_k, \quad (1.4.46)$$

where $R_p^* \alpha_{M_1 \dots M_k}(x) = \partial_{M_1} R_p^{N_1} \dots \partial_{M_k} R_p^{N_k} \alpha_{N_1 \dots N_k}(x)$ (a definition that will find its proper place in Section 4.1.4). So an orientifold acts a bit like a mirror, dictating that the physics in one half of spacetime should be a specular copy of that in the other half.

R_p has a $(p+1)$ -dimensional fixed locus, which is called an *orientifold plane* and denoted by Op , in analogy with the notation for D-branes. Op -planes have a tension and RR charge, as Dp -branes do; but as we will see, they can have *negative tension*, so they are in a sense a source of antigravity. Usually objects with negative mass can generate instabilities,¹⁶ but this is avoided here because Op -planes are stuck at the fixed locus and are not dynamical. Since open strings do not end on an O -plane (unless D-branes also happen to be located on top of it), the open string modes (1.3.22) are absent, and in particular there is no transverse fluctuation x^i .

Op -planes also preserve half-supersymmetry in flat space, as D-branes do; this makes them stable under time evolution. The preserved supercharges are again those that satisfy (1.3.51).

Op_{\pm} -planes

The action on open strings needs to be specified separately. World-sheet parity still reverses the orientation of the string, and now exchanges the two endpoints: $\Omega: \sigma^1 \rightarrow \pi - \sigma^1$. Since Ω exchanges the endpoints, it should also reverse the order of CP labels (Section 1.3.6). Omitting other quantum numbers, this would lead to $\Omega|IJ\rangle = |JI\rangle$, or more generally to

$$\Omega|IJ\rangle = \sum_{JK} M_{IK} M_{LJ}^{-1} |LK\rangle. \quad (1.4.47)$$

Imposing $\Omega^2 = 1$ gives

$$M^{-1} M^t = \mp 1. \quad (1.4.48)$$

¹⁶ Already in Newtonian mechanics, a naive argument for instability is the following: if we have two objects with mass m , $-m$, the first will feel a repulsive force, the second an attractive one, so they will both start accelerating in the same direction.

The choice of sign has important consequences on the O-plane, which is then called Op_{\pm} .¹⁷ Most notably, the tension is

$$\tau_{Op_{\pm}} = \mp 2^{p-5} \tau_{Dp} = \mp \frac{1}{32\pi^p l_s^{p+1}}. \tag{1.4.49}$$

Both Op_{\pm} are charged under C_{p+1} with a charge density equaling the tension. In other words, their effective action is

$$S_{Op} = \tau_{Op} \left[- \int d^{p+1} \sigma e^{-\phi} \sqrt{-\det g|_{Op}} + \int_{Op} C_{p+1} \right], \tag{1.4.50}$$

to be compared with (1.3.24). Notice that the terms involving \mathcal{F} are absent: $B = 0$ vanishes on the Op because of (1.4.46), while f vanishes because there are no dynamical fields, as already observed. Other orientifold projections are possible, but we will not need them in this book. See, for example, [111].

Dp-branes on Op-planes

Consider now Dp-branes parallel to an Op-plane, for simplicity in flat space, along $\{x^{p+1} = 0, \dots, x^9 = 0\}$. This configuration is BPS, just like parallel Dps. If the branes are not atop the O-plane, then the orientifold action dictates that they should come in mirror pairs, related by the involution R_p . So for a stack of N Dps at $\{x^{p+1} = x_0^{p+1}, \dots, x^9 = x_0^9\}$, there should be a mirror stack at $\{x^{p+1} = -x_0^{p+1}, \dots, x^9 = -x_0^9\}$. In this case, the gauge group on each stack is still $U(N)$ (see Section 1.3.6). But for a Dp stack on an Op-plane, the orientifold action relates the open string states to themselves, projecting some of them out. Consider, for example, the states that give rise to the vector field, which are the parallel components of (1.3.21) with the additional CP labels; a linear combination of such states is of the form

$$\sum_{IJ} \lambda_{IJ} b_{-1/2}^a |0, IJ\rangle_{NS}. \tag{1.4.51}$$

The action of Ω on $b_{-1/2}^a |0, IJ\rangle_{NS}$ gives a minus sign. Recalling (1.4.47), the states that survive are those such that

$$\lambda = -M\lambda^t M^{-1}. \tag{1.4.52}$$

Now the gauge group depends on whether we have an Op_+ or Op_- :

- For an Op_- -plane, the sign in (1.4.48) is -1 . Then M is symmetric; it can be brought to the identity by a change of basis ($\lambda \rightarrow C^{-1}\lambda C, M \rightarrow CMC^t$). So (1.4.52) says $\lambda \in \mathfrak{so}(N)$, and the gauge group is $SO(N)$.
- For an Op_+ -plane, the sign in (1.4.48) is $+1$, and M is antisymmetric; by a change of basis, it can be taken to be $\begin{pmatrix} 0 & 1_N \\ -1_N & 0 \end{pmatrix}$. By definition, (1.4.52) then says that $\lambda \in \mathfrak{sp}(N)$. The gauge group is $Sp(N)$, also known as $USp(2N)$.

If we start from the earlier situation with N Dp-branes parallel to the Op, and move them on top of it, the N mirror images will also do so, resulting in $2N$ Dp-branes and a gauge group $SO(2N)$ or $Sp(N)$. Sometimes one calls a pair of two mirror images a “full” brane. A single Dp cannot be moved away from an Op and is called a “half” brane.

¹⁷ In older literature, such as [110], this convention is sometimes reversed.

If a Dp -brane is instead orthogonal to an Op , it is mapped to itself by the orientifold action; the gauge group is $U(N)$ away from the orientifold, with some boundary conditions on the gauge fields that depend on how many directions the Op -plane and the Dp -brane have in common.

T-duality

T-duality acts on orientifolds as in (1.4.23) for D-branes. Under T_9 , an orientifold whose spacetime involution is a reflection R_p becomes a reflection $R_p R_9$; a factor $(-1)^{F_L}$ is added or erased according to (1.4.45). For example, on $\mathbb{R}^9 \times S^1$,

$$T_9: \Omega \rightarrow \Omega R_9. \quad (1.4.53)$$

The left-hand side Ω has no spacetime involution. The O-plane extends over all of spacetime, so we call it an O9. On the right-hand side, on the periodic direction, $x^9 \sim x^9 + L_9$, the involution $R_9: x^9 \rightarrow x^9$ has two fixed loci, at $\{x^9 = 0\}$ and $\{x^9 = L_9/2\}$. So there are *two* O8-planes. If we compactify p directions and T-dualize along all of them, we get 2^p O $(9-p)$ -planes. The \pm type is preserved by this operation, so an O9₋ will generate 2^p O $(9-p)$ ₋-planes. Notice that the total tension and RR charge is unchanged, because of (1.4.49).

It is possible to include Op -planes of different types in the same model [110]. If we have the ΩR_9 orientifold above, and choose one of the planes to be O8₊ and the other O8₋, and T-dualize back along x^9 , the result is no longer an O9, but a *shift-orientifold*, defined by Ω sh, where

$$\text{sh}: x_9 \rightarrow x_9 + L_9/2 \quad (1.4.54)$$

is the translation by half a period. This has no fixed loci, and so there are no O-planes. This is consistent with the O8_± having opposite tensions and RR charges.

Gravitational description

Even though Op -planes have a fixed position, they have a tension and charge, and thus have an effect on the gravitational and RR fields. Given the similarity between (1.4.50) and (1.3.24), this Op gravitational solution is formally identical to that for Dp -branes, (1.3.59); but the constant in the harmonic function h changes, because of the tension (1.4.49). For the Op_- , (1.3.61) changes to

$$h = 1 - 2^{p-5} \frac{r_0^{7-p}}{r^{7-p}}, \quad (1.4.55)$$

again valid for $p < 7$. (For the Op_+ , the relative sign is +.) The negative sign changes the behavior of the solution: now in the region $r < 2^{p-5} r_0$ the function h is negative, and the solution (1.3.59) is meaningless.¹⁸ This is in fact not worrisome because the supergravity approximation breaks down already well outside the hole $r < r_0$, due to large curvature and sometimes string coupling. This unphysical hole should be resolved in full-fledged string theory, much like what we expect for the singularity at $r = 0$ for Dp -branes; we will see examples in Sections 7.2.3 and 9.4.3.

¹⁸ This should not be confused with the region inside a Schwarzschild black hole horizon, where $g_{tt} < 0$ but the metric is real.

Equation (1.4.55) can also be combined with Dp -branes, similar to (1.3.62). This confirms that parallel Op -planes and Dp -branes are also simultaneously BPS, just like parallel Dp .

Here are some more details:

- For $p < 3$, at the hole boundary $r = 2^{p-5}r_0$ the scalar curvature diverges, while the string coupling goes to zero.
- For $p = 3$, $R = 0$, but other curvature invariant diverge at $r = 2^{p-5}r_0$. The string coupling is constant.
- For $3 < p < 7$, both curvature and string coupling diverge at $r = 2^{p-5}r_0$.
- For $p = 7$, the harmonic function is

$$h = 4 \frac{g_s}{2\pi} \log(r/r_0), \quad (1.4.56)$$

the -4 of difference with (1.3.63) for $D7$ -branes again due to (1.4.49). The unphysical hole is $r < r_0$.

- For $p = 8$,

$$h = h_0 - 8 \frac{g_s}{2\pi l_s} |x_9|. \quad (1.4.57)$$

For $h_0 > 0$, there is no hole: the dilaton approaches $e^\phi \sim g_s h_0^{-5/4}$. The curvature is finite except at $x_9 = 0$, where the second derivatives of h generate a delta function that can be interpreted as the effect of the $O8$ source. This is a much milder singularity than all the others we have encountered, especially for large h_0 . However, for small h_0 the string coupling is large; for $h_0 = 0$, both dilaton and curvature diverge at $x_9 = 0$.

O9-plane and type I superstrings

We now focus on the $O9$, where the spacetime involution is the identity, which we already encountered after (1.4.53). The fields that survive the projection (1.4.46) for $p = 9$ are

$$\phi, \quad g_{MN}, \quad C_2. \quad (1.4.58)$$

(The magnetic dual C_6 (1.3.40) is not independent.) The fermionic fields are also projected: only one gravitino and one dilatino remain. The branes that are still BPS should be

$$D1, \quad D5, \quad D9. \quad (1.4.59)$$

The $F1$ and the $NS5$ should not be BPS, because they would be charged under B , which has been projected out in (1.4.58). Indeed, a closed $F1$ is unstable toward breaking in open strings. Ordinarily this can happen only near D -branes, but in type I we have $D9$ -branes, which are everywhere.

The $O9$ by itself has a problem: the chiral fermions have an anomaly. In IIB, this was canceled by the contribution of C_4 (Section 1.2.3), but now that field has been projected out. A curious alternative way of noticing the problem is to focus on the WZ term in (1.4.50), which for $p = 9$ gives

$$T_{O9} \int C_{10}. \quad (1.4.60)$$

Since C_{10} only appears here, its equation of motion reads $1 = 0$. (This “tadpole” is reminiscent of the one on D0s in massive IIA (Section 1.4.1).) This point of view suggests a way out: D9-branes also have a WZ coupling to C_{10} , and adding N of them results in a total $(N T_{D9} + T_{O9}) \int C_{10}$. In view of (1.4.49), we need $N = 16$ full D9-branes, leading to a gauge group

$$\mathrm{SO}(2N) = \mathrm{SO}(32). \quad (1.4.61)$$

This is the gauge group for one of the heterotic theories, (1.1.44); the anomaly is now canceled in a similar fashion as in that theory. This O9 plus 16 (full) D9 is called a *type I superstring*, because the O9 breaks supersymmetry by half. The spacetime action S is obtained from the IIB (1.2.36) by setting to zero the fields not included in (1.4.58), $B_{MN} = C_0 = C_4 = 0$, and adding an action for the D9–O9 system. By Section 1.3.6, this is supersymmetric Yang–Mills with gauge group $\mathrm{SO}(32)$.

This finally completes the list of perturbative string theories at the beginning of Section 1.1.

1.4.5 Heterotic dualities

In Sections 1.4.1–1.4.3, we have seen that IIA, IIB and eleven-dimensional supergravity are all connected by a web of dualities, which can be explained by postulating a theory in $d = 11$ of extended objects called M-theory. We will now see how to obtain from M-theory the remaining perturbative string theories, namely type I and the two heterotic theories [112, 113].

M-theory on an interval and heterotic strings

We begin by considering M-theory on a spacetime

$$\mathbb{R}^{10} \times I, \quad (1.4.62)$$

with I an interval. In general, even in QFT, introducing space boundaries requires care with boundary conditions; in a theory of gravity, where geometry is dynamical, one might even worry that a boundary can be unstable and expand. In string theory, so far we have only seen one possible spacetime boundary, the O8-plane. We can try to introduce a boundary in a similar way in M-theory, by quotienting it.

We then view $I = S^1/\mathbb{Z}_2$: if S^1 is defined by a periodic coordinate $x^{10} \sim x^{10} + 2\pi R_{10}$, then \mathbb{Z}_2 identifies opposite points according to $x^{10} \rightarrow -x^{10}$. The two fixed loci of this action, $\{x^{10} = 0\}$ and $\{x^{10} = \pi R_{10}\}$, are the endpoints of the interval. Once we take the product with \mathbb{R}^{10} , these loci become ten-dimensional boundaries. Unfortunately, this $x^{10} \rightarrow -x^{10}$ is not a symmetry: in the $d = 11$ action (1.4.4), the CS term $A_3 \wedge G_4 \wedge G_4$ is not invariant, because its definition contains an ϵ tensor (similar to (1.2.19)), which picks up a minus sign under a parity. Said differently, $A_3 \wedge G_4 \wedge G_4$ only contains the index $M = 10$ a single time. We can cure this problem by also reversing the sign of A_3 :

$$x^{10} \rightarrow -x^{10}, \quad A_3 \rightarrow -A_3. \quad (1.4.63)$$

This is now a symmetry of (1.4.4), and we can quotient M-theory by it.

In the limit where the interval is small, $R_{10} \ll l_{P11}$, the resulting quotient theory will look ten-dimensional. In a KK reduction, in first approximation we can keep only

the zero modes in x^{10} . (1.4.63) implies that the components A_{MNP} , $M, N, P \neq 10$, are odd functions of x^{10} ; so the zero KK mode, which would be the constant, is projected out. The components A_{MN10} are even functions, and the zero KK mode remains. So the ten-dimensional theory we obtain has a two-form B_{MN} , but no three-form. There are also zero modes for the metric and dilaton; so the $d = 10$ bosonic fields are those of the common NSNS sector or of the bosonic string, (1.1.9).

Equation (1.4.63) should be supplemented by the spinorial parity action, multiplication by Γ_{10} . The result for the zero modes is that we keep a positive-chirality gravitino $\psi_{M\alpha}$, and a negative-chirality dilatino $\lambda_{\dot{\alpha}}$. The supersymmetry parameter ϵ becomes a single Majorana spinor ϵ of positive chirality; so there are 16 supercharges.

The massless fields we have found in ten dimensions so far are the same as the first two rows of Table 1.3 for the heterotic theories. This is problematic, because as we saw there the fermions $\psi_{M\alpha}$ and $\lambda_{\dot{\alpha}}$ give rise to an anomaly. Before taking the limit $R_{10} \ll l_{P11}$, we might think we can have no anomaly because we are in $d = 11$ dimension, which is odd; but there can be anomalies localized on the boundaries. So on a spacetime boundary in M-theory, there should be degrees of freedom that cancel such anomalies. There are two identical boundaries, and it is natural to expect that they give the same contribution. Given that in the heterotic theory the anomaly is canceled by the χ_{α}^a , it must then be that each of the two boundaries gives half that contribution.

One of the heterotic gauge groups (1.1.44) is indeed a product of two copies of E_8 ; so the anomaly is canceled if on each $d = 10$ spacetime boundary there is a χ_{α}^a in the adjoint of E_8 . Supersymmetry then demands that a bosonic vector field should also be present. In other words, the remaining two lines of Table 1.3 are also present in the spectrum, but are localized on the two spacetime boundaries, each with an E_8 gauge group.

We conclude then that the $R_{10} \ll l_{P11}$ limit of M-theory on $\mathbb{R}^{10} \times I$ is the $E_8 \times E_8$ heterotic string:

$$\text{M-theory on } I \cong E_8 \times E_8 \text{ heterotic} . \quad (1.4.64)$$

M-theory on (1.4.62) can be regarded as a nonperturbative completion of heterotic $E_8 \times E_8$, much as M-theory on $\mathbb{R}^{10} \times S^1$ can be regarded as a nonperturbative completion of IIA.

Heterotic T-duality

The two heterotic theories are *T-dual*. To see this, we put both theories on $\mathbb{R}^9 \times S^1$. This is more complicated than IIA or IIB on such space: the length of the S^1 is now no longer the only parameter. There is also A_9 , which is physical because of (1.2.14). By a gauge transformation, we can make it lie in the Cartan subalgebra, which for both heterotic gauge groups (1.1.44) is 16-dimensional. So we have 17 physical parameters:

$$(A_9^I, L_9) . \quad (1.4.65)$$

The gauge group is broken to the commutant of A_9^I . On generic points of the parameter space, this is $U(1)^{16}$; moreover, there is an extra $U(1)^2$ coming from g_{M9} and B_{M9} , much as in (1.1.24) for the bosonic string. At the special points where $A_9^I = 0$ in either heterotic theory, this is enhanced back to

$$E_8 \times E_8 \times U(1)^2, \quad SO(32) \times U(1)^2. \quad (1.4.66)$$

There is also a world-sheet description of such S^1 compactifications [114, 115]. In Section 1.1.3, the requirements of level matching constrained the T^{16} for the left-moving bosons so much that only two possibilities remained, leading to the two heterotic theories. Compactifying one spacetime direction, we have 17 compact left-movers and one compact right-mover; the spectrum analysis is a mix of those for the heterotic theory and for the compactified bosonic string, (1.1.23). We now need an $(1 + 17)$ -dimensional lattice. These are all mathematically equivalent under rotations that mix all 18 bosons, but physically we should regard as equivalent only those that are related by rotations of the 17 left-movers. Another possible perspective is that we should choose a single linear combination of the 17 compact left-movers to pair with the single compact right-mover. This again gives 17 physical parameters, as in (1.4.65). As for the gauge group, a generic $(1 + 17)$ -lattice won't have elements of length two, so the generic gauge group is Abelian. For special values of the lattice, one can obtain (1.4.66).

From the world-sheet point of view, we didn't have to specify which of the two heterotic theories we were discussing. Before compactifications, there are two possibilities; after compactification, there is a 17-dimensional space of possibilities, and each point in this space can be thought of as arising from either of the two heterotic theories. In other words, a configuration obtained from the $E_8 \times E_8$ theory for one choice of (1.4.65) also arises from the $SO(32)$ theory for a different choice of parameters. In this identification, the length of the S^1 gets inverted, as in type II T-duality. In this sense, the two heterotic theories are T-dual.

Heterotic-type I duality

Earlier in this subsection, we have argued that the strong-coupling limit of the $E_8 \times E_8$ heterotic theory is M-theory on (1.4.62). What is the strong-coupling limit of the $SO(32)$ heterotic theory?

The $SO(32)$ gauge group suggests that it might be type I. Indeed, this works at the level of the effective action. The map is (1.4.31), applied to the fields (1.4.58) that survive the O9 orientifold projection, and leaving the $SO(32)$ gauge field unchanged. In particular, the C_2 potential of type I becomes the heterotic B ; the D1-brane of type I becomes the heterotic F1. We also argued after (1.4.59) that the F1 in type I is unstable, and this is dual to the lack of D-branes in the heterotic theory.

M-theory on a cylinder

The dualities of this subsection can be summarized conveniently by considering M-theory on

$$\mathbb{R}^9 \times S^1 \times I, \quad (1.4.67)$$

sketched in Figure 1.4. This is similar to how M-theory on $\mathbb{R}^9 \times T^2$ summarizes the dualities among M-theory, IIA, and IIB (see Figure 1.3).




Figure 1.4 M-theory on $\mathbb{R}^9 \times S^1 \times I$.

We call R_9 and R_{10} the radii of S^1 and I respectively. When both are large, spacetime is eleven-dimensional, and the best description is M-theory. When R_9 is large and R_{10} is small, we recover the $E_8 \times E_8$ heterotic string by (1.4.63). In the figure, we have gone from the top-right corner of Figure 1.4 to the bottom-right (from “M” to “HE”). If we now also make R_{10} small, naively spacetime becomes nine-dimensional, but we argued that a T-duality can make it ten-dimensional again, turning the theory into SO(32) heterotic (bottom-left corner “HO” in the figure).

For M-theory on $\mathbb{R}^9 \times T^2$, the exchange of the two radii was interpreted as the S-duality (1.4.31). This exchanges the two axes in Figure 1.3 and leaves the bottom-left IIB corner invariant, consistent with S-duality being a symmetry of IIB. For M-theory on (1.4.67), the same operation

$$R_9 \leftrightarrow R_{10} \quad (1.4.68)$$

is again interpreted as an S-duality, this time relating the SO(32) heterotic theory to type I. This is why the bottom-left corner is labeled with both theories.

The two bottom corners in the figure are related by T-duality; (1.4.68) then predicts that the duality taking type I to the top-left corner should also be a T-duality. By (1.4.53), this is a configuration with two O8₋-planes in IIA. It is called *type IA*, although it is simply an orientifold of IIA (just like type I is just an orientifold of IIB). The 16 full D9-branes in type I are T-dualized to 16 full D8-branes. Reducing directly from M-theory on the S^1 , we would have expected to find IIA on $\mathbb{R}^9 \times I$; the only boundaries we know in both type II theories are indeed O8-planes.

In type I, there are also 16 full D9-branes, which are T-dualized to 16 full D8-branes parallel to the boundaries. By (1.4.23), their positions in the interval are dual to the values of the gauge field a_9 on the D9s (in the Cartan subalgebra). Going further back in the duality chain, this originates from the values of a_9 in the two copies of E_8 that live on the M-theory boundaries.

D8-branes couple to F_{10} , the dual of the Romans mass parameter F_0 . When the D8 positions are generic, there will thus be a nonzero value for F_0 in the spacetime on the IA side, thus achieving a lift of sorts of massive IIA to eleven dimensions. This map from IA to M-theory looks quite nonlocal: the position of the D8s is encoded in the values of two gauge fields that live on the boundaries of the M-theory spacetime. It is not obvious how to generalize this to cases without O8-planes.

Exercise 1.4.1 Show with an argument similar to (1.3.47) that in eleven-dimensional supergravity

$$\frac{1}{(2\pi l_{\text{P11}})^3} \int G_4 \in \mathbb{Z}, \quad \frac{1}{(2\pi l_{\text{P11}})^6} \int *G_4 \in \mathbb{Z}. \quad (1.4.69)$$

Exercise 1.4.2 The M5 solution is

$$ds_{\text{M5}}^2 = h^{-1/3} ds_{\parallel}^2 + h^{2/3} ds_{\perp}^2, \quad G_{ijkl} = 3\pi l_{\text{P11}}^3 x^m \epsilon_{mijkl} r^{-5}, \quad (1.4.70)$$

in the notation of (1.3.59): the parallel and transverse directions are six and five, and $h = 1 + r_0^3/r^3$, $r_0^3 = \pi N l_{\text{P11}}^3$.

Identify periodically one of the transverse directions, $\mathbb{R}^5 \rightarrow \mathbb{R}^4 \times S^1$ and reduce along it: namely, calling the compact direction x^{11} , write (1.4.70) in the form (1.4.5a). Check that you obtain the NS5 solution from (1.3.65).

Exercise 1.4.3 Now reduce (1.4.70) along one of the parallel directions, and check that you obtain the D4 solution from (1.3.59).

Exercise 1.4.4 Use S-duality (1.2.38) and (1.4.30) to derive the NS5 solution (1.3.65) from the D5 solution (1.3.59).

Exercise 1.4.5 Now use the same S-duality to derive the back-reaction of an F1 from the D1 solution.

Exercise 1.4.6 Finally, use the F1 solution to obtain that of an M2 whose metric reads

$$ds_{\text{M2}}^2 = h^{-2/3} ds_{\parallel}^2 + h^{1/3} ds_{\perp}^2, \quad h = 1 + \frac{r_0^6}{r^6}, \quad (1.4.71)$$

with $r_0^6 = 32\pi^2 N l_{\text{P11}}^6$. What happens if we try to obtain it by lifting a D2?

Exercise 1.4.7 Check that (1.4.46) is a symmetry of the type II actions (1.2.15) and (1.2.36).