

Dietary intake in pregnant women in a Spanish Mediterranean area: as good as it is supposed to be?

Clara L Rodríguez-Bernal^{1,*}, Rosa Ramón², Joan Quiles², Mario Murcia^{1,3},
Eva M Navarrete-Muñoz⁴, Jesús Vioque^{3,4}, Ferran Ballester^{1,3,5} and
Marisa Rebagliato^{2,3}

¹Centro Superior de Investigación en Salud Pública, CSISP, Avenida Cataluña 21, E-46020 Valencia, Spain:

²Dirección General de Salud Pública, Generalitat Valenciana, Valencia, Spain: ³CIBER de Epidemiología y Salud Pública, CIBERESP, Barcelona, Spain: ⁴Departamento de Salud Pública, H^a Ciencia y Ginecología, Universidad Miguel Hernández, Elche-Alicante, San Juan de Alicante, Spain: ⁵Escuela de Enfermería, Universitat de Valencia, Valencia, Spain

Submitted 28 July 2011: Final revision received 1 April 2012: Accepted 17 June 2012: First published online 9 August 2012

Abstract

Objective: To assess food and nutrient intakes and compliance with nutritional recommendations in pregnant women according to selected sociodemographic characteristics.

Design: Cross-sectional study based on data from the INMA-Valencia cohort (Spain), which recruited pregnant women between 2004 and 2005. Information on maternal sociodemographics and anthropometry was collected. Dietary intake was assessed through an FFQ. Intakes of foods were compared with Spanish food-based dietary guidelines. Intake inadequacy for nutrients was assessed using the Dietary Reference Intakes of the US Institute of Medicine.

Setting: Valencia, Spain.

Subjects: We studied 822 pregnant women who had information on dietary intake during their first trimester of pregnancy.

Results: More than 50% of pregnant women did not meet the guidelines for cereals and legumes; reported intakes of carbohydrates, *n*-3 and *n*-6 fatty acids were below recommendations and exceeded the total fat intake according to dietary references. Dietary inadequacy for folate, Fe and vitamin E ranged from 99% to 68%. Vegetable intake was related to age only. Younger and less educated women showed lower intakes of protein and *n*-3 fatty acids and higher intakes of *trans*-fatty acids as well as greater inadequacy for micronutrients. Spanish women reported lower intakes of fruit and carbohydrates and higher intakes of protein, total fat, SFA, MUFA and *n*-3 fatty acids compared with their foreign-born counterparts.

Conclusions: Women in the studied area have inadequate intakes of several nutrients relevant during pregnancy. Age, education and country of origin are factors significantly related to dietary intake and adequacy.

Keywords

Diet

Pregnancy

Adequacy

Nutritional guidelines

Sociodemographic characteristics

Maternal diet during the periconceptional and pregnancy period has been shown to be an important determinant of birth outcomes, such as birth defects, preterm delivery and fetal growth^(1–4), and of maternal health problems such as pre-eclampsia⁽⁵⁾ and gestational diabetes⁽⁶⁾. Moreover, nutrient intake during pregnancy has been related to cognitive development and allergic disorders in childhood^(7,8).

Nutritional requirements increase during pregnancy in order to support fetal growth and the development of maternal tissue specific to reproduction⁽⁹⁾. This may lead to deep deficiencies of micronutrients such as Fe and folate, the recommended daily intakes of which are not usually

achieved from food sources even in populations assumed to be well nourished^(10–12). As a result, recommendations on specific supplementation during pregnancy for these micronutrients have been formulated^(13,14). On the other hand, it has been shown that recommended levels of other micronutrients are met by dietary intake from food in developed countries^(12,15,16), making supplementation unnecessary in some cases.

Neither the dietary intake of pregnant women in Mediterranean areas^(17–19) (where diet is considered to be healthier than in other Western societies) nor their compliance with food-based dietary guidelines for pregnancy has been examined extensively⁽²⁰⁾. Furthermore,

*Corresponding author: Email rodriguez_claber@gva.es

from a global perspective, few studies have estimated total nutrient intakes in pregnant women taking into account food and supplements jointly^(15,19,21). Evaluating nutrient intakes from both food and supplements would reflect in a more realistic way the adequacy with respect to recommendations⁽²²⁾.

We assessed intakes from foods and supplements during the first trimester of pregnancy in a cohort of women in a Spanish Mediterranean area, and estimated the compliance with pregnancy-specific recommendations for food groups and nutrients.

Materials and methods

Population and study design

INMA (Spanish acronym for Childhood and Environment) is a network of research groups in Spain that set up a collaborative mother and child cohort study in 2003⁽²³⁾.

The present study was based on data from the INMA-Valencia birth cohort. The mother's recruitment and follow-up procedures have been described in detail elsewhere⁽²³⁾. In brief, pregnant women from a well-defined geographic area in the Valencia province, who attended their first prenatal visit at La Fe Hospital between February 2004 and June 2005, were eligible if they fulfilled the inclusion criteria⁽²³⁾. Pregnant women were identified at the hospital admission desk and were invited to participate by the team staff if they were at least 16 years old, had a singleton pregnancy, were between 10 and 13 weeks of gestation at enrolment, did not follow any programme of assisted reproduction for the current pregnancy, had no previous history of hypertension, had no communication handicap and planned to deliver at the reference hospital.

Eligible women who agreed to participate signed an informed consent form; 855 out of 1563 eligible women were included in the study (55% participation rate). Participants were similar in educational level to non-participants but the former were slightly older and more likely to work at the time of the participation request.

Of the 855 included women, twenty-three had an abortion before their first appointment, four were lost to follow-up, one abandoned the study and five were excluded because their dietary information from food sources was missing. Finally, 822 women were included in the present analysis. The study protocol was approved by the Ethics Committee of La Fe Hospital.

Data collection and processing

Information on sociodemographic and lifestyle factors and maternal anthropometric variables was obtained from a questionnaire, administered by trained interviewers, in the first trimester of pregnancy (week 12 approximately) and from maternal medical records. The variables were: age at recruitment; country of origin; educational level (primary, 6–9 years of schooling; secondary, 10–14 years of schooling;

university, ≥ 15 years of schooling); social class following a widespread Spanish classification⁽²⁴⁾; working status during the first trimester; smoking during the first trimester; pre-pregnancy BMI; parity; planned pregnancy; drug use; and self-reported vomiting during the first trimester.

Information on usual daily food and nutrient intakes during the first trimester of pregnancy was collected using an FFQ with 101 food items. The FFQ was an adapted version of Willett's questionnaire⁽²⁵⁾ developed and validated for use among adults living in Spain⁽²⁶⁾, with satisfactory coefficients for validity and reproducibility^(26,27).

In our population of pregnant women we asked how often, on average, participants had consumed a particular amount of a specific type of food from the last menstrual period until the time of the interview (10–13 weeks). Standard units and serving sizes were specified for each food item. The questionnaire included nine possible answers to determine frequency of intake, ranging from 'never or less than once per month' to 'six or more times per day'. The response to each food item was converted to average daily intake for each participant. Nutrient values and total energy intake were obtained from the US Department of Agriculture food composition tables and other published sources^(28,29). We calculated nutrient intakes by multiplying the frequency of consumption for each food item by the nutrient composition of the portion size specified on the FFQ and by addition across all foods to obtain total intake of each nutrient for each individual. Nutrient intakes were adjusted for total energy intake by calculating the residuals from a linear regression with the natural logarithm of the nutrient modelled as the dependent variable and the natural logarithm of total energy intake as the independent variable⁽²⁵⁾.

Information on supplement use was collected by asking women at their first visit (weeks 10–13) the question: 'In three months before becoming pregnant and until now, have you ever taken specific supplements or multivitamins?' Monthly intake of nutrients from supplements was estimated based on supplement brand name, composition, daily dose and timing of consumption.

From food sources, intakes of protein, carbohydrates, total fat, SFA, MUFA, PUFA, *n*-3 fatty acids, *n*-6 fatty acids, dietary fibre, vitamins A, D, E, C, B₁₂, folic acid, Ca, Fe, iodine, Zn, Mg and energy were calculated. Dietary iodine intake was assessed taking into account iodized salt consumption.

From supplements, intakes of vitamins A, D, E, C, B₁₂, folic acid, Ca, Fe, iodine and Zn were calculated using the mean intake of the first, second and third months to obtain the intake of the first trimester of pregnancy. In the case of folic acid, its intake was calculated in folate equivalents (1 μg dietary folate = 0.6 μg of folic acid from supplements). Supplementary iodine intake in our cohort was mainly from multivitamins, so we decided to consider 'iodine users' those women who took at least 100 μg of iodine from multivitamins. Then, we added the nutrient intake from

supplements to that previously obtained from diet alone, to obtain the intake 'from food and supplements'.

Lack of compliance with food-based dietary guidelines was assessed by calculating the proportion of women not reaching the minimum recommended daily number of servings of each food group according to the Spanish recommendations – those specific for pregnancy⁽³⁰⁾. The cut-off values for the food groups assessed were the lowest values of the following ranges: farinaceous foods (cereals and legumes), 4–5 servings/d; vegetables, 2–4 servings/d; fruits, 2–3 servings/d; dairy, 3–4 servings/d; protein-rich foods (meat, poultry, fish, eggs, nuts), 2 servings/d; drinks (water, herbal teas, low-sugar, non-alcoholic drinks), 4–8 glasses/d. Inadequate intake of macronutrients was calculated by taking as cut-off points the Acceptable Macronutrient Distribution Range⁽³¹⁾ and that of micronutrients using the Estimated Average Requirement^(32–36). We then estimated the percentage of the population whose intake was below the recommended values for micronutrients or outside the range established for macronutrients. These values correspond to the Dietary Reference Intakes established by the Institute of Medicine for the US population and assessment methods used in the present study were made following recommendations for evaluation of dietary adequacy in population groups^(22,37).

Statistical analysis

Descriptive statistics of maternal characteristics and dietary variables are shown by means of percentages, means and standard deviations and percentiles. Differences in the intake of food groups according to sociodemographic characteristics were contrasted using the non-parametric Kruskal–Wallis test. Since the distribution of some macronutrients was right skewed, they were transformed by the natural logarithm in order to approximate normality. Subsequently, their intake according to sociodemographic characteristics was compared using ANOVA. When significant differences were found, Tukey's *post hoc* test was performed in order to detect paired differences between groups. Homogeneity of variances was assessed with Levene's test. Heteroscedasticity ($P < 0.05$ in Levene's test) was treated using the robust Welch statistic and Dunnett's D3 *post hoc* test. We verified the robustness of ANOVA to small deviations from normality: additional analysis with the Kruskal–Wallis test led to equivalent conclusions. Differences in intake inadequacy of food groups, macro- and micronutrients according to maternal characteristics were assessed using the χ^2 test. Statistical differences were determined by a level of significance $P < 0.05$ in all tests. Statistical analysis was carried out using the statistical software package SPSS version 15.0.

Results

The mean age of our study population was 29.8 years, with 53% of the sample being ≥ 30 years of age (Table 1).

Table 1 Sociodemographic and lifestyle characteristics among pregnant women of the INMA-Valencia cohort (n 822), 2004–2005

Characteristic	<i>n</i>	%
Age (years)		
<25	99	12.0
25–29	290	35.3
≥ 30	433	52.7
Educational level		
Primary school	286	34.8
Secondary school	348	42.3
University studies	188	22.9
Country of origin		
Spain	723	88.0
Latin America	64	7.8
Others	35	4.3
Socio-economic status		
I+II (highest)	127	15.5
III	194	23.6
IV+V (lowest)	501	60.9
Number of previous pregnancies		
0	374	45.5
≥ 1	448	54.5
Smoking*	194	24.8
Alcohol consumption*	251	30.5
Drugs use*	10	1.3
Pre-pregnancy BMI (kg/m ²)		
<18.5	41	5.0
18.5–25.0	549	67.0
>25.0	230	28.0
Self-reported vomiting*		
No	474	42.3
Yes	348	57.7
Planned pregnancy		
No	198	24.4
Yes	615	75.6
Multivitamin supplements use	459	55.8
Folic acid supplements use*	785	95.5
Fe supplements use*	88	10.7
Iodine from supplements*,†	407	49.6
Iodized salt consumption*	502	61.1

*During first trimester.

†From multivitamins (at least 100 μ g).

Concerning educational level, 65% of women completed at least secondary studies. Eighty-eight per cent of the women were born in Spain. About 95% and 11% of the mothers took folic acid and Fe supplements, respectively. At least 50% of women consumed iodine through multivitamin supplements or iodized salt in early pregnancy (Table 1).

Food groups

For the whole population, median intakes (in servings/d) were within the range of recommendations for all food groups except for cereals and legumes, which was below the minimum recommended (Table 2). Older women consumed a greater amount of vegetables ($P < 0.001$); more educated women had a lower intake from the meat group ($P = 0.016$); and women of foreign origin had a higher intake of fruits compared with Spanish women ($P = 0.001$).

When frequency of compliance with recommendations was assessed (Table 3), overall, lack of compliance was very high for cereals (77%), followed by dairy (52%) and fruit and vegetables (about 47%). Equal proportions of

Table 2 Intake of food groups (servings/d) during the first trimester of pregnancy according to age, educational level and country of origin among women of the INMA-Valencia cohort (*n* 822), 2004–2005

Food group	Recommended (servings/d)	Age (years)									Education						Country origin							
		All women		<25		25–29		≥30		<i>P</i>	Primary		Secondary		University		<i>P</i>	Spain		Latin America		Others		<i>P</i>
		Median	IQR	Median	IQR	Median	IQR	Median	IQR		Median	IQR	Median	IQR	Median	IQR		Median	IQR	Median	IQR	Median	IQR	
Cereals and legumes	4–5	3·1	1·9	3·2	2·2	3·1	1·8	3·0	1·9	0·600	3·2	1·9	3·1	1·8	2·9	2·1	0·255	3·1	1·8	2·4	2·1	3·5	3·1	0·095
Vegetables	2–4	2·1	1·5	1·6	1·6	2·1	1·6	2·3	1·5	<0·001	2·0	1·6	2·2	1·6	2·2	1·3	0·437	2·1	1·6	2·2	1·3	2·4	1·8	0·378
Fruits	2–3	2·0	1·8	2·0	1·9	2·0	1·8	2·1	1·9	0·306	2·1	2·1	2·1	1·8	2·0	1·5	0·857	2·0	1·8	2·6	2·2	2·7	2·3	0·001
Milk and dairy	3–4	2·9	2·0	2·8	2·1	2·9	2·0	3·0	2·0	0·897	3·1	2·1	2·9	1·9	2·7	2·2	0·187	2·9	2·0	3·2	2·3	3·1	2·0	0·618
Meat, poultry, fish and eggs	2	2·3	0·9	2·4	0·8	2·3	0·9	2·3	0·9	0·789	2·4	0·9	2·3	0·8	2·2	0·9	0·016	2·3	0·9	2·2	0·8	2·4	0·9	0·119
Non-alcoholic low-sugar drinks	4–8	6·5	2·3	6·5	2·1	6·6	2·6	6·5	2·3	0·712	6·5	2·1	6·5	2·4	6·5	2·6	0·813	6·5	2·4	6·5	2·4	6·6	2·5	0·179

IQR, interquartile range (25th–75th percentile).
P values obtained using the Kruskal–Wallis test.

Table 3 Percentage not meeting recommendations* for food intake during the first trimester of pregnancy according to age, educational level and country of origin among women of the INMA-Valencia cohort (*n* 822), 2004–2006

Food group	Recommended (servings/d)	Age (years)						Education				Country of origin			
		All women	<25	25–29	≥30	<i>P</i>	Primary	Secondary	University	<i>P</i>	Spain	Latin America	Others	<i>P</i>	
		%	%	%	%		%	%	%		%	%	%		
Cereals and legumes	4–5	76·6	69·7	77·9	77·4	0·216	75·9	77·9	75·5	0·771	77·6	76·6	57·1	0·020	
Vegetables	2–4	46·2	62·6	48·3	41·1	0·000	49·0	44·5	45·2	0·514	47·6	35·9	37·1	0·110	
Fruits	2–3	47·9	50·5	50·3	45·7	0·410	47·9	46·6	50·5	0·679	49·8	34·4	34·3	0·016	
Milk and dairy	3–4	52·3	56·6	53·8	50·3	0·440	49·0	53·4	55·3	0·340	53·1	45·3	48·6	0·441	
Meat, poultry, fish and eggs	2	30·2	25·3	29·0	32·1	0·349	29·0	26·1	39·4	0·006	29·7	39·1	22·9	0·187	
Non-alcoholic low-sugar drinks	4–8	11·6	7·1	13·1	11·5	0·269	13·3	8·9	13·8	0·124	11·2	17·2	8·6	0·304	

P values obtained using the χ^2 test.

*Food-based dietary guidelines of the Spanish Society of Community Nutrition.

Spanish and Latin-American women did not meet guidelines for cereals intake (about 77%); regarding vegetable consumption, significant differences were found by age only, with the highest percentage of women not meeting dietary guidelines being observed among those under 25 years ($P < 0.001$); approximately 50% of Spanish women had a fruit intake below the minimum recommended and inadequacy was lower (about 34%) for foreign-born women ($P = 0.016$; Table 3).

Macronutrients

The distribution of macronutrient intakes for the whole sample is presented in Appendix 1.

Regarding differences in mean intakes by selected maternal characteristics (Table 4), it was found that the intakes of protein and *n-3* fatty acids increased, whereas intake of *trans*-fatty acids decreased, with older age ($P < 0.001$). Women with the lowest educational level had the lowest intakes of protein ($P = 0.002$) and *n-3* fatty acids ($P = 0.012$) and the highest intake of *trans*-fatty acids ($P = 0.003$) but also the highest consumption of fibre ($P = 0.048$). Women of Spanish origin had the highest intakes of protein ($P < 0.001$), total fat ($P < 0.001$), SFA ($P = 0.008$), MUFA ($P < 0.001$) and *n-3* fatty acids ($P < 0.001$); Latin-American women had higher intake of carbohydrates ($P < 0.001$) and women from other origins, higher intake of *trans*-fatty acids ($P = 0.050$; Table 4).

When the adequacy with respect to the Institute of Medicine's recommendations was assessed (Table 5), it was found that about 57% of women had an intake of carbohydrates under the recommended range. Virtually all women had protein intake which fell within recommendations. Total fat intake was too high (almost 71% of women exceeded the recommendation). More than 50% of women had deficient intakes of *n-3* and *n-6* fatty acids (Table 5). Spanish women had the highest percentage of inadequacy for carbohydrates (61% had low intakes, $P < 0.001$) and total fat (74% exceeded the recommended consumption, $P < 0.001$). Latin-American women showed the highest percentages of inadequacy for *n-3* fatty acids (almost 77% of them were below the recommendation, $P < 0.001$). Additionally, younger ($P < 0.001$) and less educated women ($P = 0.048$) showed the highest percentages of inadequacy for *n-3* fatty acids because of low intake (Table 5).

Micronutrients

The distribution of micronutrient intakes (from food only and total intake – food plus supplements) for the whole sample is presented in Appendix 2. It is worthy to highlight a very high – sevenfold – increase in folate intake after including intake from supplements: 298 µg/d from food only *v.* 2112 µg/d from food plus supplements.

Regarding inadequacy, data obtained from food intake only, showed that the greatest percentages of inadequacy for the whole study population were observed – in

Table 4 Intake of macronutrients (g/d) during the first trimester of pregnancy according to age, educational level and country of origin among women of the INMA-Valencia cohort (n 822), 2004–2005

Macronutrient	Age (years)						Education						Country of origin										
	<25		25–29		≥30		Primary		Secondary		University		Spain		Latin America		Others						
	Mean	sd	Mean	sd	Mean	sd	Mean	sd	Mean	sd	Mean	sd	Mean	sd	Mean	sd	Mean	sd					
Protein	100.5	14.5	95.4 ^a	12.3	98.9 ^a	14.4	102.6 ^b	14.6	<0.001	98.0 ^a	14.4	101.7 ^b	14.8	101.8 ^b	13.6	0.002	101.2 ^a	14.4	93.5 ^b	14.4	98.1 ^{a,b}	13.9	<0.001
Carbohydrates*	249.5	1.2	256.0	1.2	250.6	1.2	247.3	1.2	0.079	250.4	1.2	249.3	1.2	248.5	1.2	0.846	246.4 ^a	1.1	274.5 ^b	1.2	270.7 ^b	1.2	<0.001
Total fat	96.1	14	95.3	15.2	96.1	13.6	96.2	14.1	0.843	96.6	14.6	95.6	13.7	96.3	13.8	0.634	97.1 ^a	13.4	88.5 ^b	16	88.2 ^b	16.3	<0.001
SFA	29.9	5.9	29.9	6.1	30.3	5.8	29.8	5.9	0.53	30.1	5.4	29.8	6.3	30.1	5.8	0.789	30.3 ^{a†}	5.7	27.8 ^{b†}	6.6	28.2 ^{a,b†}	7.4	0.008†
Trans-fatty acids*	1.49	1.5	1.80 ^a	1.5	1.60 ^b	1.5	1.36 ^c	1.5	<0.001	1.59 ^{a†}	1.5	1.46 ^{b†}	1.6	1.40 ^{b†}	1.5	0.003†	1.47 ^{a†}	1.5	1.52 ^{a,b}	1.5	1.76 ^b	1.5	0.050
MUFA	44.8	8.8	43.8	8.9	44.7	8.8	45.2	8.8	0.338	45	9.2	44.6	8.6	44.9	8.8	0.852	45.6 ^a	8.5	39.5 ^b	8.9	39.1 ^b	9.3	<0.001
PUFA*	13.9	1.3	14.1	1.3	13.9	1.2	13.9	1.3	0.903†	14.0	1.3	13.8	1.2	13.9	1.3	0.731	13.9	1.3	13.8	1.4	13.8	1.3	<0.001
n-3 fatty acids*	1.51	1.3	1.36 ^a	1.2	1.49 ^b	1.3	1.56 ^c	1.2	<0.001	1.47 ^a	1.3	1.55 ^b	1.3	1.52 ^{a,b}	1.2	0.012	1.54 ^a	1.2	1.31 ^b	1.2	1.39 ^b	1.2	<0.001
n-6 fatty acids*	12.2	1.3	12.5	1.3	12.2	1.3	12.1	1.3	0.601†	12.3	1.3	12.0	1.3	12.2	1.3	0.508	12.2	1.3	12.3	1.4	12.2	1.3	0.972†
Dietary fibre*	22.3	1.4	23.1	1.4	21.8	1.5	22.5	1.4	0.293	23.2 ^a	1.4	22.1 ^{a,b}	1.4	21.4 ^b	1.4	0.048	22.2	1.4	21.9	1.4	24.8	1.6	0.181

P values obtained using ANOVA, unless otherwise stated.
^{a,b,c}Mean values within a row with unlike superscript letters were significantly different using Tukey's post hoc test or †Dunnett's T3 post hoc test ($P < 0.05$).
*Variables were transformed by the natural logarithm before analysis. Geometric means and geometric standard deviations are presented.
†P value obtained using Welch's test.

Table 5 Intake inadequacy* of selected macronutrients during the first trimester of pregnancy according to age, educational level and country of origin among women of the INMA-Valencia cohort (*n* 822), 2004–2005

Macronutrient	Type of inadequacy	Age (years)					Education				Country of origin			
		All women	<25	25–29	≥30	<i>P</i>	Primary	Secondary	University	<i>P</i>	Spain	Latin America	Others	<i>P</i>
		%	%	%	%		%	%	%		%	%	%	
Protein	Deficit	0.2	1.0	0.3	0.0	0.167	0.3	0.3	0.0	0.734	0.0	3.1	0.0	<0.001
	Excess	0.0	0.0	0.0	0.0		0.0	0.0	0.0		0.0	0.0	0.0	
Carbohydrates	Deficit	57.4	51.5	58.3	58.2	0.291	57.7	56.0	59.6	0.350	61.1	29.7	31.4	<0.001
	Excess	0.2	1.0	0.3	0.0		0.7	0.0	0.0		0.0	1.6	2.9	
Total fat	Deficit	0.2	0.0	0.3	0.2	0.799	0.3	0.0	0.5	0.732	0.1	0.0	2.9	<0.001
	Excess	70.8	68.7	73.1	69.7		71.7	69.8	71.3		74.0	42.2	57.1	
<i>n</i> -3 fatty acids	Deficit	54.1	72.7	56.2	48.5	<0.001	59.1	49.4	55.3	0.048	51.5	76.6	68.6	<0.001
	Excess	0.4	0.0	0.7	0.2		0.0	0.9	0.0		0.4	0.0	0.0	
<i>n</i> -6 fatty acids	Deficit	62.9	55.6	62.1	65.1	0.288	58.4	66.4	63.3	0.284	63.2	60.9	60.0	0.621
	Excess	2.1	2.0	1.4	2.5		2.4	1.4	2.7		1.8	4.7	2.9	

P values obtained using the χ^2 test.

*Values of intake inadequacy are expressed as the percentage of women whose intake is below (deficit) or above (excess) the Acceptable Macronutrient Distribution Range established for each macronutrient by the US Institute of Medicine.

Table 6 Intake inadequacy* for selected micronutrients from food and food plus supplements during the first trimester of pregnancy according to age, educational level and country of origin among women of the INMA-Valencia cohort (*n* 822), 2004–2005

Nutrient	Age (years)					Education				Country of origin			
	All women	<25	25–29	≥30	<i>P</i>	Primary	Secondary	University	<i>P</i>	Spain	Latin America	Others	<i>P</i>
	%	%	%	%		%	%	%		%	%	%	
From food													
Vitamin A ($\mu\text{g}/\text{d}$)	4.6	12.1	5.5	2.3	<0.001	8.0	3.2	2.1	0.003	4.8	1.6	5.7	0.465
Vitamin C (mg/d)	14.4	22.2	15.5	11.8	0.022	16.1	14.9	10.6	0.234	14.8	9.4	14.3	0.495
Vitamin D ($\mu\text{g}/\text{d}$)	99.8	100.0	99.8	99.7	0.832	100.0	99.7	99.5	0.504	99.7	100.0	100.0	0.872
Vitamin E (mg/d)	67.8	77.8	69.0	64.7	0.036	71.0	67.5	63.3	0.215	67.8	65.6	71.4	0.840
Folate ($\mu\text{g}/\text{d}$)	99.6	99.0	99.3	100.0	0.169	99.7	99.4	100.0	0.574	99.6	100.0	100.0	0.814
Fe (mg/d)	67.9	71.7	72.4	64.0	0.040	77.6	65.2	58.0	<0.001	68.2	71.9	54.3	0.177
Iodine ($\mu\text{g}/\text{d}$)	24.3	31.3	24.8	22.4	0.171	28.3	22.1	22.3	0.150	24.1	21.9	34.3	0.346
Ca (mg/d)	5.5	11.1	4.8	4.6	0.031	6.3	6.0	3.2	0.290	5.1	6.2	11.4	0.266
From food & supplements													
Vitamin A ($\mu\text{g}/\text{d}$)	3.0	8.1	3.4	1.6	0.003	5.2	1.7	2.1	0.026	3.0	1.6	5.7	0.516
Vitamin C (mg/d)	8.9	15.2	9.7	6.9	0.029	10.5	8.6	6.9	0.398	9.0	6.3	11.4	0.657
Vitamin D ($\mu\text{g}/\text{d}$)	88.1	91.9	87.9	87.3	0.439	88.1	89.1	86.2	0.611	87.7	85.9	100.0	0.077
Folate ($\mu\text{g}/\text{d}$)	40.8	51.5	42.8	37.0	0.020	42.7	42.0	35.6	0.263	39.7	43.8	57.1	0.107
Fe (mg/d)	13.4	29.3	15.5	8.3	<0.001	16.4	14.9	5.9	0.002	11.6	26.6	25.7	<0.001
Iodine ($\mu\text{g}/\text{d}$)	50.9	57.6	54.3	47.1	0.061	59.8	49.3	40.4	<0.001	51.3	52.4	40.0	0.413
Ca (mg/d)	14.2	21.2	14.1	12.7	0.091	15.7	13.8	12.8	0.633	13.6	18.8	20.0	0.317
Vitamin A ($\mu\text{g}/\text{d}$)	4.1	7.1	3.8	3.7	0.294	4.2	4.9	2.7	0.466	3.7	4.7	11.4	0.081

P values obtained using the χ^2 test.

*Values of intake inadequacy are expressed as the percentage of women whose intake is below (deficit) or above (excess) the Dietary Reference Intake established for each micronutrient by the US Institute of Medicine.

descending order – for vitamin D, folate, Fe and vitamin E (Table 6). Virtually none of the women of our study reached the recommended intake for folate or vitamin D (under deficient sun exposure) and nearly 68% of them did not comply with recommendations for Fe or vitamin E either. Intake inadequacy was significantly higher in younger women for vitamins A, C, E and Fe and Ca ($P < 0.001$, $P = 0.022$, $P = 0.036$, $P = 0.040$ and $P = 0.031$, respectively). With regard to educational level, the overall pattern was a decrease in inadequacy as level of studies increased. Nevertheless, this relationship was significant for vitamin A ($P = 0.003$) and Fe ($P < 0.001$) only (Table 6). Although Latin-American women had a lowest proportion of inadequacy for all nutrients except for Fe and folate, the

differences found according to women's country of origin were not statistically significant.

When the addition of supplements to dietary consumption was assessed (Table 6), an important reduction in inappropriate intake was observed for all the nutrients examined, especially for folate. None the less, inadequacy remained high for vitamin E, Fe and vitamin D (40.8%, 50.9% and 88.1%, respectively). Differences regarding maternal characteristics followed a similar pattern to that found for intake from food only, except that according to women's country of origin, foreign-born women showed higher inadequacy for folic acid than Spain-born women (about 26% *v.* 12%; $P < 0.001$). Regarding vitamin D and Ca intakes, there was a trend showing that inadequacy

did not change in non-Latin-American foreign women after intake from supplements was considered.

Discussion

Pregnant women in our study did not reach the recommendations established for this life stage for some food groups, macro- and micronutrients. A very high percentage of women were found to have an intake of cereals and legumes, and to a lesser extent, of fruit and vegetables, below the recommendations. Composition of the diet regarding macronutrients was not adequate, according to recommendations. More than 50% of women had insufficient intakes of carbohydrates, *n*-3 and *n*-6 fatty acids and about 70% exceeded the recommendations for total fat. The greatest deficiencies in micronutrient intakes were found for vitamin D, folate and Fe. Supplementation reduced the frequency of intake inadequacy, but not in a substantial way for vitamin D, Fe and vitamin E. It was also found that dietary intake varied significantly according to socio-demographic characteristics such as age, education and country of origin.

Intakes by food group

Ferrer *et al.* performed a cross-sectional study in pregnant women in a Spanish Mediterranean island⁽²⁰⁾, finding intakes of cereals, legumes and vegetables below the recommendations, which coincides with our results. Nevertheless, the consumption of fruit and vegetables found in our study is more adequate than that seen in non-Mediterranean settings⁽¹⁵⁾.

Taking into account sociodemographic characteristics, we found that older women consumed a greater amount of vegetables. Studies in other industrialized settings^(15,38,39) have found the same association. Nevertheless, no differences according to educational level were found for fruit and vegetable intakes in our study. This evidence differs from studies in non-Mediterranean settings in which higher education was related to higher vegetable intake^(15,38,40). In agreement with our findings, Ferrer *et al.* did not find significant differences in fruit or vegetable intake by educational level⁽²⁰⁾. Furthermore, there is evidence that in Southern European countries, fruit and vegetable consumption is common due to cultural and economic reasons and not determined by occupation or education^(40–42).

Country of origin was found to be related to fruit intake. In Spain only one study, to our knowledge, has included foreign-born pregnant women when examining dietary intake and found no differences in food consumption⁽²⁰⁾. One study in the USA comparing the diet of Mexican immigrants with that of Mexican-Americans found that Mexico-born women consumed significantly more fruits, grains and dairy products than US-born women⁽⁴³⁾, partially coinciding with our results.

Macronutrient intakes

Other studies set in Spain and in Portugal also revealed a diet with an excessive content of total fat and a poor intake of carbohydrates^(17,44,45). None of these studies assessed macronutrient intake by sociodemographic characteristics. Seventy-four per cent of Spanish women in the present study had an excessive total fat intake; it is important to note, however, that it came mainly from MUFA. Nevertheless, SFA intake was also high. Latin-American, younger and less educated women were found to have the highest percentages of inadequacy for *n*-3 fatty acids because of low intake. Regarding carbohydrates, Spanish women showed the highest percentage of inadequacy because of low intake. Evidence suggests that the quality of lipids and carbohydrates during pregnancy is important. Excessive consumption of saturated fat and low *n*-3 fatty acid intake, as well as high intakes of sugar and lower intakes of starchy food, have been related to adverse maternal and infant health outcomes^(46–49). Thus, pregnant women in the studied area should be advised to reduce their fat intake, especially from saturated fats (reducing the consumption of foods such as red and processed meats), and increase their consumption of *n*-3 fatty acids (raise their consumption of fish other than red tuna or swordfish, and also that of walnuts and leafy vegetables) as well as their intake of starchy carbohydrates from sources such as cereals and legumes.

Micronutrient intakes

The mean vitamin D intake from food found in the present study was lower than that reported in other studies including Nordic or Mediterranean pregnant women^(15,17,19,21). In our population, virtually all women had inadequate intakes of vitamin D according to the Institute of Medicine reference values. When taking into account total intake (food plus supplements), the percentage of inadequacy observed was still elevated, and higher than that reported in studies among Nordic pregnant women^(15,21). None the less, it is important to consider that the Estimated Average Requirement established for vitamin D is based on an assumption of minimal or no sun exposure. In the Valencia region, sun exposure might be relatively high all year round, so it is likely that the inadequacy is lower than that estimated in the present study. Yet, women should be encouraged to consume foods rich in vitamin D such as dairy products, enriched breakfast cereals and types of fish recommended for pregnancy. Dietary deficiencies of other important nutrients such as folate, Fe and vitamin E, as found in the present study, are not uncommon in industrialized countries^(12,16,17,36,41,50). The fact that a high percentage of women consumed low amounts of folate sources such as cereals, vegetables and fruit partly explains our results. Supplementation greatly reduced deficient intakes for folic acid; however, it increased mean intake in a disproportionate way, which might lead to an important proportion of women taking daily doses exceeding the safety limits as shown in a previous study

within this cohort⁽⁵¹⁾. Considering sociodemographic characteristics, younger, less educated and foreign-born women showed the greatest percentage of inadequacy for folate after intake from supplements was considered. As fruit and vegetables are widely available in Mediterranean settings and not related to socio-economic status, strategies at the population level should be established to promote a higher consumption. Besides, the appropriateness in the use of folic acid supplements in the periconceptional period and during pregnancy as well as the causes of differences found by maternal characteristics should be assessed as part of public health policies.

The small reduction in inadequacy for Fe found after considering supplements consumption is likely to be due to the current recommendations in Spain of starting supplementation in the second half of pregnancy⁽⁵⁰⁾. Nevertheless, it is important to consider that, even in developed countries, some women might start pregnancy with deficient Fe status⁽⁵²⁾, which is a strong predictor of maternal and infant adverse outcomes^(53,54).

Other studies in Mediterranean areas have found mean intakes and/or percentages of inadequacy for vitamin E from food similar to those shown here^(17–19), which makes evident the need to promote a higher consumption of vitamin-E-rich foods other than olive oil, of which intake is already high. Such foods include nuts and seeds. Vitamin E might protect against maternal and childhood illnesses such as pre-eclampsia or asthma^(8,55).

Iodine intake from different sources (foods, iodized salt and supplements) has already been described in our pregnant population⁽⁵⁶⁾, as has its effect on maternal thyroid function⁽⁵⁷⁾ and infant neurodevelopment⁽⁵⁸⁾. In summary, inadequate intake of iodine was very low in women who consumed either iodized salt or iodine supplements; however, a high iodine intake from supplements was associated with both maternal thyroid dysfunction and decreased psychomotor scores in infants at the age of 1 year. These findings suggest the need to further evaluate the safety and effectiveness of iodine supplementation during pregnancy in iodine-sufficient or mildly deficient populations.

The differences in dietary intake and adequacy found according to the maternal characteristics studied reveal the influence of socio-economic and cultural factors. There is evidence that populations are less likely to meet dietary guidelines if food plans are not socially or culturally accepted⁽⁵⁹⁾. Consumption at population level depends also on food supply and availability⁽⁶⁰⁾. Therefore, public policies aiming to reduce food-related diseases should take into account these issues.

The present study has some limitations such as the moderate participation rate, which could make the generalization of our results difficult; however, when sociodemographic characteristics of participants and non-participants were assessed, no significant differences were found, except for the fact that participants were

slightly older and more likely to work at the time of enrolment. As older age was related to better adequacy with respect to recommendations, one could expect that in the general population the inadequacy will, if anything, be slightly higher than that found here. Regarding the validity of dietary intake estimates, the validity and reproducibility of a similar version of the FFQ were satisfactory when comparing the FFQ with four 1-week dietary records in an adult population of the same region.

Conclusions and implications for public health

An important percentage of women in the studied area did not meet the guidelines for certain food groups and were also found to have inadequate intakes of several nutrients which are relevant during pregnancy. Age, education and country of origin seem to be variables strongly related to dietary intake and adequacy.

A nutrient-rich and balanced diet is necessary and must be promoted in women of childbearing age before pregnancy, to avoid the risk of deficient intake until women know they are pregnant or their first antenatal visit takes place. In the light of the situation observed in the present study, policies to improve supplementation schemes based on the nutritional status of pregnant women are needed in order to reduce the likelihood of both deficient and excessive intakes and the resulting potential risks for health. Monitoring the nutritional status of pregnant women is important and a good start point. However, based on these findings, and acknowledging that people's food consumption is conditioned to a certain extent by factors different from individual choices, it would be also necessary to implement intervention programmes that take into account social and cultural contexts and guarantee a healthier food environment.

Acknowledgements

Source of funding: This work was supported by grants from Instituto de Salud Carlos III (Rd INMA G03/176 and CB06/02/0041), the Spanish Ministry of Health (FIS 03/1615, FIS 04/1509, FIS 04/1436, FIS 05/1079, FIS 06/1213, FIS 06/0867, FIS 09/02647), Conselleria de Sanitat – Generalitat Valenciana and Fundació Roger Torné. *Conflict of interest:* None of the authors declared a conflict of interest. *Authors' contributions:* C.L.R.-B. and R.R. conceived the study, conducted the data analysis and prepared the manuscript; J.Q. contributed to the conceptual approach and manuscript preparation; M.M. carried out the statistical analysis and contributed to the data interpretation and manuscript preparation; E.M.N.-M assisted with the nutritional data and provided feedback; J.V. designed the FFQ and contributed to manuscript preparation; F.B. and M.R. supervised the data collection, helped with data interpretation and manuscript preparation, and provided critical

revision of the manuscript. *Acknowledgements*: The authors would particularly like to thank all participants for their generous collaboration; Amparo Cases and Amparo Quiles for their support with the fieldwork and the collection of nutritional data; and Daniel Giménez and Manoli Garcia for their assistance with the nutritional data. A full list of the INMA Project researches can be found at <http://www.proyectoINMA.org>.

References

- Scholl T & Johnson W (2000) Folic acid: influence on the outcome of pregnancy. *Am J Clin Nutr* **71**, 5 Suppl., 1295S–1303S.
- Olsen S & Secher N (2002) Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: prospective cohort study. *BMJ* **324**, 447.
- Ramón R, Ballester F, Iñiguez C *et al.* (2009) Vegetable but not fruit intake during pregnancy is associated with newborn anthropometric measures. *J Nutr* **139**, 561–567.
- Rodríguez-Bernal C, Rebagliato M, Iñiguez C *et al.* (2010) Diet quality in early pregnancy and its effects on fetal growth outcomes: the Infancia y Medio Ambiente (Childhood and Environment) Mother and Child Cohort Study in Spain. *Am J Clin Nutr* **91**, 1659–1666.
- Olafsdottir A, Skuladottir G, Thorsdottir I *et al.* (2006) Relationship between high consumption of marine fatty acids in early pregnancy and hypertensive disorders in pregnancy. *BJOG* **113**, 301–309.
- Thomas B, Ghebreskel K, Lowy C *et al.* (2006) Nutrient intake of women with and without gestational diabetes with a specific focus on fatty acids. *Nutrition* **22**, 230–236.
- Oken E, Wright R, Kleinman K *et al.* (2005) Maternal fish consumption, hair mercury, and infant cognition in a US cohort. *Environ Health Perspect* **113**, 1376–1380.
- Nurmatov U, Devereux G & Sheikh A (2011) Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol* **127**, 724–733.
- Picciano M (2003) Pregnancy and lactation: physiological adjustments, nutritional requirements and the role of dietary supplements. *J Nutr* **133**, issue 6, 1997S–2002S.
- Erkkola M, Karppinen M, Järvinen A *et al.* (1998) Folate, vitamin D, and iron intakes are low among pregnant Finnish women. *Eur J Clin Nutr* **52**, 742–748.
- Giddens J, Krug S, Tsang R *et al.* (2000) Pregnant adolescent and adult women have similarly low intakes of selected nutrients. *J Am Diet Assoc* **100**, 1334–1340.
- Turner R, Langkamp-Henken B, Littell R *et al.* (2003) Comparing nutrient intake from food to the estimated average requirements shows middle- to upper-income pregnant women lack iron and possibly magnesium. *J Am Diet Assoc* **103**, 461–466.
- Becker D, Braverman L, Delange F *et al.* (2006) Iodine supplementation for pregnancy and lactation – United States and Canada: recommendations of the American Thyroid Association. *Thyroid* **16**, 949–951.
- World Health Organization (2006) Iron and folate supplementation. In: *Integrated Management of Pregnancy and Childbirth. Standards for Maternal and Neonatal Care*, section 1.8. Geneva: Department of Making Pregnancy Safer, WHO.
- Arkkola T, Uusitalo U, Pietikäinen M *et al.* (2006) Dietary intake and use of dietary supplements in relation to demographic variables among pregnant Finnish women. *Br J Nutr* **96**, 913–920.
- Mouratidou T, Ford F, Prountzou F *et al.* (2006) Dietary assessment of a population of pregnant women in Sheffield, UK. *Br J Nutr* **96**, 929–935.
- Pinto E, Barros H & dos Santos Silva I (2009) Dietary intake and nutritional adequacy prior to conception and during pregnancy: a follow-up study in the north of Portugal. *Public Health Nutr* **12**, 922–931.
- Arija V, Cucó G, Vila J *et al.* (2004) Consumo, hábitos alimentarios y estado nutricional de la población de Reus en la etapa preconcepcional, el embarazo y el posparto. *Med Clin (Barc)* **123**, 5–11.
- Petrakos G, Panagopoulos P, Koutras I *et al.* (2006) A comparison of the dietary and total intake of micronutrients in a group of pregnant Greek women with the Dietary Reference Intakes. *Eur J Obstet Gynecol Reprod Biol* **127**, 166–171.
- Ferrer C, García-Esteban R, Mendez M *et al.* (2009) Determinantes sociales de los patrones dietéticos durante el embarazo. *Gac Sanit* **23**, 38–43.
- Haugen M, Brantsaeter AL, Alexander J *et al.* (2008) Dietary supplements contribute substantially to the total nutrient intake in pregnant Norwegian women. *Ann Nutr Metab* **52**, 272–280.
- Murphy S, Barr S & Poos M (2002) Using the new dietary reference intakes to assess diets: a map to the maze. *Nutr Rev* **60**, 267–275.
- Guxens M, Ballester F, Espada M *et al.* (2011) Cohort profile: the INMA – Infancia y Medio Ambiente – (Environment and Childhood) Project. *Int J Epidemiol* (Epublication ahead of print version).
- Domingo-Salvany A, Regidor E, Alonso J *et al.* (2000) Una propuesta de medida de la clase social. *Aten Primaria* **25**, 350–363.
- Willett W, Sampson L, Stampfer M *et al.* (1985) Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol* **122**, 51–65.
- Vioque J, Weinbrenner T, Asensio L *et al.* (2007) Plasma concentrations of carotenoids and vitamin C are better correlated with dietary intake in normal weight than overweight and obese elderly subjects. *Br J Nutr* **97**, 977–986.
- Vioque J (2006) Validez de la evaluación de la ingesta dietética. In *Nutrición y Salud Pública. Métodos, bases científicas y aplicaciones*, 2nd ed., pp. 199–210 [L Serra Majem and J Aranceta Bartrina, editors]. Mason-Elsevier: Barcelona.
- US Department of Agriculture, Agricultural Research Service (2007) USDA National Nutrient Database for Standard Reference Release 21. <http://www.nal.usda.gov/fnic/foodcomp/Data/SR21/nutrlist/sr21a306.pdf> (accessed November 2008).
- Palma I, Farran A & Cantós (2008) *Tablas de Composición de Alimentos por Medidas Caseras de Consumo Habitual en España*. CESNID. Madrid: McGraw-Hill Interamerican.
- Sociedad Española de Nutrición Comunitaria (2007) Guía de la Alimentación Saludable. <http://www.nutricioncomunitaria.org/index.jsp> (accessed March 2011).
- Food and Nutrition Board, Institute of Medicine (2005) *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids (Macronutrients)*. Washington, DC: The National Academies Press.
- Food and Nutrition Board, Institute of Medicine (2001) *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington, DC: The National Academies Press.
- Food and Nutrition Board, Institute of Medicine (2000) *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids*. Washington, DC: The National Academies Press.

34. Food and Nutrition Board, Institute of Medicine (1998) *Dietary Reference Intakes for Thiamine, Riboflavin, Niacin, Vitamin B₆, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline*. Washington, DC: The National Academies Press.
35. Food and Nutrition Board, Institute of Medicine (2001) *Dietary Reference Intakes*. Washington, DC: National Academy Press.
36. Food and Nutrition Board, Institute of Medicine (2011) *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press.
37. Barr S, Murphy S & Poos M (2002) Interpreting and using the dietary references intakes in dietary assessment of individuals and groups. *J Am Diet Assoc* **102**, 780–788.
38. Bodnar L & Siega-Riz A (2002) A diet quality index for pregnancy detects variation in diet and differences by sociodemographic factors. *Public Health Nutr* **5**, 801–809.
39. Uusitalo L, Uusitalo U, Ovaskainen ML *et al.* (2008) Sociodemographic and lifestyle characteristics are associated with antioxidant intake and the consumption of their dietary sources during pregnancy. *Public Health Nutr* **11**, 1379–1388.
40. Darmon N & Drewnowski A (2008) Does social class predict diet quality? *Am J Clin Nutr* **87**, 1107–1117.
41. Roos G, Johansson L, Kasmel A *et al.* (2001) Disparities in vegetable and fruit consumption: European cases from the north to the south. *Public Health Nutr* **4**, 35–43.
42. Prättälä R, Hakala S, Roskam AJ *et al.* (2009) Association between educational level and vegetable use in nine European countries. *Public Health Nutr* **12**, 2174–2182.
43. Harley K, Eskenazi B & Block G (2005) The association of time in the US and diet during pregnancy in low-income women of Mexican descent. *Paediatr Perinat Epidemiol* **19**, 125–134.
44. Ortiz-Andrellucchi A, Sánchez-Villegas A, Ramírez-García O *et al.* (2009) Calidad nutricional de la dieta en gestantes sanas de Canarias. *Med Clin (Barc)* **133**, 615–621.
45. Cárcel C, Quiles J, Rico B *et al.* (2005) Adecuación de la ingesta nutricional de embarazadas de segundo y tercer trimestre. *Rev Esp Nutr Comunitari* **11**, 133–144.
46. Armitage JA, Taylor PD & Poston L (2005) Experimental models of developmental programming: consequences of exposure to an energy rich diet during development. *J Physiol* **565**, 3–8.
47. Leung BM & Kaplan BJ (2009) Perinatal depression: prevalence, risks, and the nutrition link – a review of the literature. *J Am Diet Assoc* **109**, 1566–1575.
48. Scholl TO, Sowers M, Chen X *et al.* (2001) Maternal glucose concentration influences fetal growth, gestation, and pregnancy complications. *Am J Epidemiol* **154**, 514–520.
49. Farmer G, Russell G, Hamilton-Nicol DR *et al.* (1988) The influence of maternal glucose metabolism on fetal growth, development and morbidity in 917 singleton pregnancies in nondiabetic women. *Diabetologia* **31**, 134–141.
50. Sociedad Española de Ginecología y Obstetricia SEGO (2010) Protocolos Segeo 'Anemia y embarazo'. <http://www.prosego.com> (accessed July 2011).
51. Navarrete-Muñoz EM, Giménez Monzó D, García de La Hera M *et al.* (2010) Ingesta dietética y de suplementos de ácido fólico en mujeres embarazadas de Valencia. *Med Clin (Barc)* **135**, 637–643.
52. Hercberg S, Preziosi P & Galan P (2001) Iron deficiency in Europe. *Public Health Nutr* **4**, 537–545.
53. Casanueva E, Pfeiffer F, Drijanski A *et al.* (2003) Iron and folate status before pregnancy and anemia during pregnancy. *Ann Nutr Metab* **47**, 60–63.
54. Rodríguez-Bernal CL, Rebagliato M & Ballester F (2012) Maternal nutrition and fetal growth: the role of iron status and intake during pregnancy. *Nutr Diet Suppl* **4**, 25–37.
55. Jain SK & Wise R (1995) Relationship between elevated lipid peroxides, vitamin E deficiency and hypertension in preeclampsia. *Mol Cell Biochem* **151**, 33–38.
56. Murcia M, Rebagliato M, Espada M *et al.* (2010) Iodine intake in a population of pregnant women: INMA mother and child cohort study, Spain. *J Epidemiol Community Health* **64**, 1094–1099.
57. Rebagliato M, Murcia M, Espada M *et al.* (2010) Iodine intake and maternal thyroid function during pregnancy. *Epidemiology* **21**, 62–69.
58. Murcia M, Rebagliato M, Iñiguez C *et al.* (2011) Effect of iodine supplementation during pregnancy on infant neurodevelopment at 1 year of age. *Am J Epidemiol* **173**, 804–812.
59. Maillot M, Darmon N & Drewnowski A (2010) Are the lowest-cost healthful food plans culturally and socially acceptable? *Public Health Nutr* **13**, 1178–1185.
60. Cannon G (2002) Nutrition: the new world disorder. *Asia Pac J Clin Nutr* **11**, Suppl. 3, S498–S509.

Appendix 1

Daily intake of selected macronutrients during the first trimester of pregnancy among women of the INMA-Valencia cohort (n 822), 2004–2005

	Mean	SD	P5	P25	P50	P75	P95
Energy (kJ)	9680.5	88.3	6118.3	7941.6	9296.8	11 024.4	14 538.1
Energy (kcal)	2313.7	21.1	1462.3	1898.1	2222.0	2634.9	3474.7
Protein (g)	102.5	25.6	62.7	84.9	100.3	119.5	149.1
Carbohydrates (g)	261.8	83.7	143.9	206.1	249.3	305.7	423.9
Total fat (g)	99.2	30.1	59.8	78.4	95.6	114.9	156.4
SFA (g)	31.1	10.9	16.9	23.8	29.6	36.7	50.6
Trans-fatty acids (g)	1.7	0.9	0.6	1.0	1.4	2.1	3.5
MUFA (g)	46.1	14.3	26.2	35.5	44.9	53.7	71.0
PUFA (g)	14.9	6.3	8.3	11.1	13.3	17.1	27.3
n-3 fatty acids (g)	1.6	0.5	0.9	1.2	1.5	1.8	2.6
n-6 fatty acids (g)	13.2	5.9	7.1	9.5	11.6	15.1	24.8
Dietary fibre (g)	23.7	8.1	11.9	18.0	22.8	28.6	39.0

P, percentile.

Appendix 2

Daily intake of selected micronutrients during the first trimester of pregnancy among women from the INMA-Valencia cohort (n 822), 2004–2005

	Nutrient intakes							Reference values	
	Mean	sd	P5	P25	P50	P75	P95	EAR	UL
From food									
Vitamin A (µg/d)*	1395.6	821.8	557.3	864.6	1217.1	1656.3	2813.4	550	3000
Vitamin D (µg/d)	3.0	1.7	1.0	1.8	2.6	3.8	6.6	10	100
Vitamin C (mg/d)	140.9	76.1	52.4	86.0	123.6	176.9	278.5	70	2000
Vitamin E (mg/d)†	11.1	3.0	7.2	9.1	10.8	12.6	16.3	12	1000¶
Vitamin B ₁₂ (µg/d)	9.7	4.9	4.6	6.6	8.6	11.2	18.8	2.2	ND
Folate (µg/d)	298.1	75.3	192.6	240.3	289.3	346.7	440.7	520	1000¶
Fe (mg/d)	20.7	3.4	15.6	18.3	20.4	22.8	27.0	22	45
Ca (mg/d)	1275.9	342.3	788.4	1025.3	1253.1	1467.9	1891.2	800	2500
Iodine (µg/d)‡	218.1	76.5	98.7	162.3	214.8	269.8	341.9	160	1100
Zn (mg/d)	27.2	4.8	19.8	23.9	26.6	30.0	35.9	9.5	40
From food & supplements									
Vitamin A (µg/d)	1474.8	825.4	623.9	963.3	1289.1	1730.4	2926.6	550	3000
Vitamin D (µg/d)	5.3	3.8	1.3	2.5	4.3	7.0	13.3	10	100
Vitamin C (mg/d)	158.9	79.7	59.9	100.9	143.2	198.3	298.9	70	2000
Vitamin E (mg/d)	14.0	5.0	7.9	10.4	12.9	16.7	23.1	12	1000¶
Vitamin B ₁₂ (µg/d)	12.1	13.2	5.8	8.1	10.4	13.8	23.0	2.2	ND
Folate (µg/d)§	2111.6	2995.4	406.0	644.5	880.1	1214.7	8775.1	520	1000¶
Fe (mg/d)	26.8	18.3	16.3	19.2	21.9	26.1	60.7	22	45
Ca (mg/d)	1309.2	345.0	823.3	1057.4	1285.6	1512.3	1912.9	800	2500
Iodine (µg/d)	262.9	96.9	116.2	192.5	258.2	322.7	435.0	160	1100
Zn (mg/d)	28.1	5.1	20.2	24.6	27.6	31.1	37.9	9.5	40

P, percentile; EAR, Estimated Average Requirement; UL, Upper Limit; ND, not determined.

*Calculated as retinol activity equivalents.

†Calculated as α-tocopherol equivalents.

‡Calculated taking into account iodized salt consumption.

§Calculated as folate equivalents.

||As preformed vitamin A only.

¶Applies to synthetic forms only.