Abstract

Cite this article: (2021). E-Poster Presentation. *European Psychiatry* 64(S1), S183–S608.

Anxiety disorders and somatoform disorders

EPP0001

Associations between social anxiety and avoidance, attachment styles and parental marital status, in late adolescence

B. Rodrigues Maia^{1*}, C. Coelho², M. Marques³ and F. Carvalho⁴

¹Faculty Of Philosophy And Social Sciences, Centre For Philosophical And Humanistic Studies, Portugal, Universidade Católica Portuguesa, Braga, Portugal; ²Faculty Of Philosophy And Social Sciences, Universidade Católica Portuguesa, Braga, Portugal; ³Coimbra Hospital And University Centre, Portugal, Coimbra Hospital and University Centre, Coimbra, Portugal and ⁴Espaço Psicológico – Consultório De Psicologia, Coimbra, Portugal, Espaço Psicológico – Consultório de Psicologia, Coimbra, Portugal, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.484

Introduction: The relation between insecure general attachment and social anxiety has long been established.

Objectives: To explore the associations between social interaction and performance anxiety and avoidance, attachment styles, and parental marital status.

Methods: 146 Portuguese adolescents, with a mean age of 18.99 years old (SD = .848; range: 18–20), filled in the Social Interaction and Performance Anxiety and Avoidance Scale and the Experiences in Close Relationships-Relationship Structures Questionnaire.

Results: Distress/Anxiety was correlated with avoidance attachment to mother and father ($r_s = .17^*$, $p = .04$; $r_s = .18^*$, $p = .03$), to anxious attachment to romantic partner ($r_s = .21^*$, $p = .01$), and to anxious and avoidance attachment to best friend ($r_s = .25^{**}$, $p = .00$; ($r_s = .17^*$, $p = .035$). Avoidance was significantly correlated with avoidance to father and to romantic partner ($r_s = .18^*$, $p = .03$), and to anxious and avoidance attachment to best friend ($r_s = .21^{**}$, $p = .009$; $r_s = .18^*$, $p = .03$). A significant difference was found in avoidance attachment to father $X^2 = 10.246$ (4, $n = 146$), $p = .036$, by parental marital status, with the adolescents with single/divorced parents presenting a higher mean score ($M_d = 111.10$; $M_d = 82.93$) than the other groups.

Conclusions: Distress/anxiety seems to be associated with more close relationships, and a single/divorced status with Avoidance. Longitudinal studies are needed to explore if insecure attachment to parents predicts insecure extra-familial attachment, and to explore the long-term effects of parental marital status.

Keywords: social anxiety and avoidance; attachment styles; parental marital status; late adolescence

EPP0002

Anxiety and depressive symptoms in patients with psoriasis

R. Sallemi, M. Bouhamed, R. Masmoudi, I. Feki and J. Masmoudi*

Psychiatry, Hedi chaker hospital, Sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.485

Introduction: Psoriasis is a chronic inflammatory skin disease that affects approximately 2% of the population. It seems to have a multifactorial aetiology and it can be considered as a psychosomatic disorder.

Objectives: To determine risk factors for anxiety and depression in psoriasis

Methods: Case-control study including 44 subjects with psoriasis and 50 controls without psoriasis. All participants answered the Hospital Anxiety and Depression Scale (HADS) to measure the severity of anxiety and depression

© The Author(s), 2021. Published by Cambridge University Press on behalf of the European Psychiatric Association. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

EUROPEAN PSYCHIATRIC ASSOCIATION