Research Article

A conceptual model of multi-scale formation processes of open-air Middle Paleolithic sites in the arid Negev desert, Israel

Maya Oron^{a,b*} , Joel Roskin^c, Yoav Avni^d, Naomi Porat^d, Emil Aladjem^b, Dmitry Yegorov^b, Jacob Vardi^b and Erella Hovers^a

^aInstitute of Archaeology, The Hebrew University of Jerusalem, Mt. Scopus, Jerusalem 91905, Israel; ^bIsrael Antiquities Authority, Nahman Avigad 1, Jerusalem, 9370726, Israel; ^cDepartment of Geography and Environment, Bar Ilan University, Ramat-Gan 5290002, Israel and ^dGeological Survey of Israel, 32 Yesha'ayahu Leibowitz St. Jerusalem 9692100, Israel

Abstract

Open-air surface accumulations and scatters of material cultural remains often are perceived as less-reliable archaeological archives, where it is difficult to distinguish anthropogenic versus geogenic formation processes or to assess their specific effects on the integrity of archaeological records. Here we analyze the depositional histories of three Middle Paleolithic open-air sites in the Negev desert of Israel, combining archaeological and geomorphological methods to create a conceptual model of multi-scale effects on the archaeological remains. Relying on the long research history in archaeology and geomorphology in the Negev, we show that integration of archaeological and geomorphological methodologies provides nuanced insights to our understanding of the archaeological record. The links established between regional and local geomorphic processes and lithic taphonomy by applying such a multi-scale analysis further allow back-tracking environmental processes from flint taphonomic attributes. Placing each site within the range of regional and local processes of exposure and burial by using informed and critically evaluated data helps to create a robust regional archaeological data base. We suggest that our approach is useful in other arid zone contexts and may have implications for understanding Pleistocene population movements across such regions.

Keywords: Open-air sites, Arid environments, Post-deposition processes, Lithic taphonomy, Middle Paleolithic, Geoarchaeology (Received 28 October 2022; accepted 22 May 2023)

INTRODUCTION

The term "site" is applied to a broad range of archaeological occurrences, from limited artifact surface scatters to thick stratified anthropogenic accumulations (Bar-Yosef and Goren, 1980). One of the confounding factors underlying the undiscriminating use of the term is the effects of post-depositional processes, which often blur distinctions between single events and multiple recurrences of archaeological deposition and accumulations. In open-air sites (OAS), such processes take place at diverse scales—from microscopic local disturbances to large-extent landscape processes.

Because OAS lack physical boundaries, they may present 'lateral stratigraphies,' in which sequential occupations are located side-by-side rather than in vertical sequences. In such cases, the temporal relationship between the remains of the occupations may be difficult to reconstruct. Given this combination of reasons, open-air anthropogenic accumulations of material culture often are perceived as partial, distorted, and potentially time-averaged archaeological records, where the effects of underlying anthropogenic versus geogenic factors are difficult to pry apart. Surface

*Corresponding author email address: mayaoron@gmail.com

Cite this article: Oron M, Roskin J, Avni Y, Porat N, Aladjem E, Yegorov D, Vardi J, Hovers E (2023). A conceptual model of multi-scale formation processes of open-air Middle Paleolithic sites in the arid Negev desert, Israel. *Quaternary Research* 116, 162–180. https://doi.org/10.1017/qua.2023.31

artifact scatters are treated with special caution, given that they may represent either remains of in situ occupations, or of deflated in situ occupations, or are entirely post-depositional scatters caused by geomorphic processes. Therefore, many researchers have viewed surface Paleolithic materials as less reliable, and therefore less informative, archaeological archives compared with sheltered or in situ OA sites, arguably to the degree that their investigation may lead to fallacious interpretations of long-term cultural processes (e.g., Coco and Iovita, 2020; Coco et al., 2020; but see Holdaway and Davies, 2019; Douglass et al., 2023).

Researchers have attempted to evaluate the scientific value of archaeological data extracted from OAS, focusing on the formation of palimpsests and possible ways to disentangle them at evolutionary and anthropological time scales, by using various types of empirical archaeological and geological data to discriminate between human and natural agents (see for example: Stern, 1994; Bailey, 2007; Malinsky-Buller et al., 2011; Vaquero et al., 2012; Hovers et al., 2014, and references therein). Several problems inherent to the use of surface assemblages in Paleolithic research can be overcome by combining refitting analysis and attribute analysis, which focus on lithic technology (Chiotti et al., 2007, 2009), and by relying on landscape-scale geomorphology and site-scale sedimentology (Thompson et al., 2014; Wright et al., 2017; Burrough et al., 2022).

Archaeological thinking about site integrity in arid zones has been especially concerned with landscape stability and with the

© University of Washington. Published by Cambridge University Press, 2023. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



possible outcomes of prolonged exposure of surface archaeological remains to climate-driven geomorphic processes that control landscape evolution (e.g., Cancellieri and di Lernia, 2013). Arid zone surface sites may be highly resilient and preserve highintegrity archaeological signatures where the landscape is stable and relatively uninterrupted biologically or geomorphologically. Such sites, however, may also be prone to winnowing. On the other hand, some arid zone sites, especially those located along highly inclined slopes, may suffer from high erosion rates. Researchers focused their attempts on potential ways to discern single events of human activity (Hallinan and Shaw, 2015) and assess their preservation (Cziesla, 1990a, b) in order to quantify human presence and its effect on desert landforms and desert pavements (Foley, 1981; Foley and Lahr, 2015). Another research focus has been to extract archaeological information at large geographical scales in geomorphologically dynamic areas where landscape evolution caused erosion and artifact movements at a number of spatial scales (e.g., Goldberg and Brimer, 1983; Olszewski et al., 2010; Groucutt et al., 2017; Wright et al., 2017; Hardaker, 2020).

In this study we consider the potential archaeological merits of OAS in arid zones. Based on empirical data from the extensive research history in the Negev desert region in southern Israel, we combine archaeological and geomorphological methods to build a conceptual model of multi-scale effects on the archaeological remains. We show that integration of methodologies from both disciplines provides nuanced insights to our understanding of the archaeological record. By integrating the archaeological and geomorphic data sets in a multi-scale analytical approach, archaeologists may gain better options of interpreting regional effects on human evolution while the artifactual contexts provide high-resolution controls for site-specific geomorphic processes. Such insights then can be incorporated into syntheses and behavioral models on a regional scale.

We present a 'proof-of-concept' study for assessing the feasibility of the model suggested here for future study of OAS. Specifically, we apply the approach to three OAS in the central and northern Negev desert of Israel: Nahal Aqev (unit 7) (henceforth Nahal Aqev), Nahal Yitnan 7, and Giv'at Barne'a. All three OAS are placed within the chronological framework of the Middle Paleolithic (MP) period, yet under three different depositional scenarios. Our multi-scale approach allows us to assess the integrity of the three sites as sources of archaeological data and compare them, relying on geomorphic conditions supported by a plethora of geomorphic and Quaternary studies of the Negev to interpret the target sites (see Enzel et al., 2008; Enzel and Bar-Yosef, 2017). Our investigation focuses on the degree and kind of lithic taphonomic modification (Hiscock, 1985) in relation to variable temporal and spatial scales of landscape processes.

GEOLOGY AND QUATERNARY GEOMORPHIC EVOLUTION OF THE STUDY REGION

The studied sites are situated along a southwest-northeast transect that parallels the northwestern fringes of the Levantine-Syrian Arc fold structures of the northern and central Negev (Fig. 1). The Syrian Arc includes a series of parallel northeast-trending monoclines and synclines exposing limestone, dolomite, chalk, and flint of Upper Cretaceous to Tertiary age. Folding initiated during the late Turonian and continued to the early Eocene (Eyal and Reches, 1983). During the Eocene, thick carbonaceous sediments were deposited in the synclines, forming the sub-horizontal carbonate strata of the Avedat Group in the Avedat Plateau and the hills surrounding the Beer-Sheva valley. The sediments of Early Miocene fluvial conglomerates and sandstones of the Hazeva Formation (Calvo and Bartov, 2001) were deposited on the Oligocene surface of the central and northern Negev, truncating the ridge crests and the top of the geological section deposited in the synclines of the Syrian Arc structures (Avni et al., 2012).

The present elevation of the region has been controlled by differential uplift since the Middle Miocene, influenced by Dead Sea rifting that generated a general uplift of its flanks (Avni, 2017). This tectonic uplift, which ceased during the late Early Pleistocene (ca. 1 Ma), determined the two main ephemeral




Figure 1. (a) Map of the Negev desert showing the 3 studied open-air sites and other MP sites in the region. (b) Soils and rainfall map (after Amit et al., 2011) with sample sites (in red).

channel systems currently in the Negev desert area: one flowing towards the Mediterranean Sea in the northwest and the other into the Dead Sea Rift in the east.

The drainage basins of the central and northern Negev were shaped by phases of fluvial incision and aggradation during the Pliocene and the Pleistocene (Avni et al., 2017; Matmon and Zilberman, 2017). Simultaneously, fluvial, aeolian, weathering, and soil-formation processes were active and shaped the presentday landscape. These include bedrock weathering (Yair, 1983; Wieler et al., 2021), cycles of loess deposition, erosion and redeposition, differential slope runoff generation (Wieler et al., 2016), fluvial incision and deposition, calcrete formation (Vogel and Geyh, 2008) and the formation of Reg soils through incorporation of dust and salt into the clastic sediments, clast shattering, and surface patination. Several fluvial terraces, dated to the last 0.5 Ma, have developed as a result of incision and episodic deposition of fluvial sediments, mainly controlled by climatic fluctuations (Goldberg, 1976, 1986; Porat et al., 2010; Matmon et al., 2016; Avni et al., 2017, 2021). These terraces are mapped and named from the older group of terraces (Q1), deposited during the Middle Pleistocene, to the youngest (Q3) terrace, deposited during the Late Pleistocene glacial phase (Avni and Wieler, 2013; Avni et al., 2017, 2021). The sites under discussion fall within the Q2 depositional phases that covered the Middle-Late Pleistocene transition.

Soil and sediments of the central and northern Negev, which comprise the main substrate for prehistoric sites, are dominated by primary and secondary loess deposits. Primary loess deposits, several meters thick, which originated as desert dust (Yaalon and Dan, 1974), blanketed the highlands and valleys of the central and northern Negev during the Middle and Late Pleistocene (Bruins and Yaalon, 1980; Pye and Tsoar, 1987; Crouvi et al., 2008, 2009; Avni et al., 2021). Most of the currently preserved loess cover in the Negev accumulated during the last major event of loess deposition, dated to ca. 70-20 ka, corresponding to the last glacial period (MIS 4-2) (Crouvi et al., 2008), with the most intense accumulation taking place ca. 40-20 ka. Substantial older loess deposits were eroded during MIS 5 (Lucke et al., 2019; Avni et al., 2021). Remains of earlier loess deposits, predating MIS 4, were preserved as relatively small-scale relicts in hilltop deposits (Crouvi et al., 2008, 2009) and within fluvial terraces deposited along the upper parts of main drainage basins of the central Negev highlands, such as in the upper Zin valley and the upper tributaries of Nahal (Wadi) Arod (Faershtein et al., 2016; Avni et al., 2021). The fine, clay-size fraction of the loess cover in the Negev originates from distal sources in the Sahara Desert, while the coarser, silt-sized grains are winnowed (Roskin et al., 2014) and possibly abraded (Crouvi et al., 2008) from relatively proximal sources, namely the Nile Delta and Sinai-Negev erg (Ben-Israel et al., 2015; Roskin and Tsoar, 2017).

Primary loess was eroded from the slopes and redeposited along the Negev highland valleys as alluvial and colluvial fills (Bowman et al., 1986), that are up to \sim 20 m thick in the northern Negev plains (Zilberman, 1991). Several paleosols with distinct carbonate-rich Bk horizons, evidence of cycles of deposition and pedogenesis, developed in the primary and secondary deposits (Zilberman, 1991; Crouvi et al., 2009; Robins et al., 2021).

Altogether, the loess deposits were pivotal components in the long- and short-term soil formation as well as slope and fluvial sedimentological processes. Most of the soils in the study region, such as calcic Loessial Serozems (Calciorthids) and Reg soils (Dan

and Raz, 1970), are comprised of substantial amounts of aeolian or reworked loess that form an important component of the soils. Archaeological deposits have been affected by these processes and may serve as markers for determining accumulation, fixation, and erosion rates (Lucke and Bäumler, 2021).

THE SITES

To construct a simple model with explicit parameters, we selected three sites—Nahal Agev, Nahal Yitnan 7, and Giv'at Barne'a (Fig. 1a)—as examples for distinctly different depositional scenarios of lithic artifacts on the open landscape. We acknowledge that these are not the only possible scenarios of site formation histories. Still, for the purposes of the present research we chose to focus on these discrete case studies for clarity's sake. The three sites are all situated in a zone that today corresponds to an arid climate receiving an average of 75–100 mm annual precipitation (Fig. 1b; Amit et al., 2011). Each site is situated within the upper reaches of a major drainage basin (Nahal Zin, Nahal Besor, and Wadi Al-Arish, respectively) of the central-northern Negev and the northern Sinai Peninsula. Floods in main ephemeral watercourses occur 1-3 times a year, with some of the large floods lasting for 1-2 days. Flood duration averages seven days per annum (Alexandrov et al., 2003).

Accumulation histories are analyzed and explained in the Results section. We note here that our field observations, archaeological, geomorphic, and sedimentological analyses suggest that each site has undergone distinct landscape evolution processes, therefore each assemblage represents a different accumulation scenario in terms of its nature and duration.

The three sites were subject to analyses on three different scales: (1) the regional scale, where each site was placed in the context of the large geological and geomorphic processes that controlled landscape evolution in the Negev; (2) the site-specific scale of formation processes; and (3) assemblage-scale flint taphonomy. Importantly, the main raw material in all three assemblages is flint extracted from geological formations of Eocene age (Broun, 1967; Benjamini, 1979), exposed in immediate proximity to each of the sites. This reduces (albeit does not remove) the likelihood that lithic taphonomic characteristics are dictated by raw material differences.

Nahal Agev

Nahal Aqev is an in situ archaeological assemblage that was sealed by sediment shorty after its accumulation; therefore, it has been subjected to a single process—continuous burial in an active fluvial-colluvial wadi terrace. The archaeological site was initially excavated in the 1970s (Munday, 1977). During re-excavations in 2015–2016, several archaeological horizons were exposed below the formerly excavated ones. Of these newly excavated horizons, we analyzed unit 7, which is well confined stratigraphically and dated by luminescence methods (Barzilai et al., 2022).

Nahal Yitnan 7

Nahal Yitnan 7 is an assemblage with a complex history of episodic burial by loess sediments and pedogenesis, and partial exposure on the surface mainly due to slope processes. The assemblage was retrieved from a surface collection and from a shallow excavation of a horizon of a dense scatter of flaked flint with maximum depth of 30 cm below the eroded modern surface (Yegorov, 2017).

Giv'at Barne'a

Giv'at Barne'a is a flint quarry and knapping site adjacent to a flint outcrop, at the top of a bedrock-dominated hill. Here, artifacts exposed on the surface and associated by typo-technological traits to several periods from the MP to Holocene, were collected and mapped.

HYPOTHESES

Given the geomorphic history of the study region, we posit that (1) the stratigraphic location of sites in relation to the regional loess cover is chronologically informative and enables their allocation to the time span of MIS 4–2 or to earlier or later time periods (we would thus expect that dates of sites embedded in loess sequences will fall within this time range); (2) the local geomorphic processes at each site are informative about cycles of deposition and erosion within the main periods of loess cover (this information in turn may be useful in creating a more accurate and nuanced geomorphic history); (3) taphonomic effects in lithics, as known from the literature (see Methods section and Supplementary data 4), may be linked to anthropogenic activities or/and to environmental processes (i.e., if the sources of taphonomic modifications are understood, they can be informative about broader scale processes in the geomorphic record). We use this information as a means to build a model of site preservation integrity that can be used to create an informed reconstruction of human settlement, activity, and movement across the region. However, a detailed discussion of the latter is outside the scope of this paper.

MATERIALS AND METHODS

Geomorphic field and laboratory methods

Stratigraphic sections at the local site scale were described by conventional geological and sedimentological field methods and placed within the regional stratigraphic scheme provided by 1:200,000 and 1:50,000 geological maps of the Geological Survey of Israel (Roded, 1996; Zilberman and Avni, 2005; Wdowinski et al., 2012; Avni and Wieler, 2013). In addition, field surveys and recent Google Earth imagery were used for complimentary data on the study sites.

Samples were sieved to finer than 2 mm and analyzed for particle size distribution (PSD) with a Beckman-Coulter LS grain-size analyzer (Fraunhofer.rf780d diffraction). A representative aliquot of \sim 2 ml subsample was treated to dissolve carbonates and disperse fines. Total inorganic carbon (IC) was measured with a Primacs SLC TOC Analyzer. Munsell color was determined in the laboratory on dry samples.

Samples for optically stimulated luminescence (OSL) dating were collected at each site from excavated sediment sections related to the archaeological finds. The sediments were mostly loess and loess derivatives (reworked, mixed with gravel, or with additional local carbonates). The dated very fine-grained quartz sand (90–125 μm) grains originated from distal loess sources and are presumed to have been well bleached at the time of deposition (see Supplementary data 2 for sampling preparation and analytical protocols).

OSL dating followed protocols after Faershtein et al. (2016). Equivalent doses (De) were measured on the purified quartz using the OSL signal and the single aliquot regenerative dose (SAR) protocol (Murray and Wintle, 2000). Eighteen to 34 aliquots (2 mm) were measured for each sample, and the average De and errors were calculated using the central age model (CAM; Galbraith and Roberts, 2012). Dose rates were calculated from concentrations of the radioactive elements U, Th, and K, measured on the additional sample by ICP-MS (U & Th) or ICP-OES (K). Moisture contents were estimated at $5\pm3\%$, as appropriate for this arid to hyperarid region, and the cosmic dose was evaluated from current burial depths.

Lithic taphonomy

The lithic samples

From the perspective of lithic technology, MP assemblages from southern Levantine sites conform to a general classification of Levantine Mousterian—characterized by the conspicuous appearance of Levallois technology (Bar-Yosef, 2006), which is absent or very rare in previous or successive periods in the Levant (Goren-Inbar, 2011; Malinsky-Buller, 2016; Goder-Goldberger, 2020; Zaidner and Weinstein-Evron, 2020). Procedures of the Levallois knapping system resulted in several indicative artifact categories such as Levallois cores, Levallois target items (points, flakes, and blades), and specific core trimming elements (Boëda, 1995). Since Levallois technology is diagnostic of MP knapping and produces some technologically indicative artifacts, we analyzed Levallois items from each site, assuming they fall within the MP even if no absolute dating was available. Hence, samples for lithic taphonomic analysis included all artifacts from Nahal Agev and all Levallois cores and products from the other two assemblages. In Nahal Yitnan 7 the samples include all Levallois products excavated and collected (4.2% of the total assemblage); for Giv'at Barne'a, only a sample (~40% of all artifacts) was counted to techno-typological groups. All Levallois cores and products were analyzed (3% of the counted assemblage).

Taphonomic attributes

There does not appear to be any clear linear timeline for formation of the damages impacting flint items (Burroni et al., 2002). While the intensity of alteration cannot be used as a relative chronological marker, types of impact can serve as indicators of assemblage history and accumulation processes. By contextualizing lithic taphonomic attribute states (e.g., states of breakage, surface alteration), we explored their use as meaningful tools for back-tracking natural processes that affected a site. A sample from each site (see above) was subjected to an attribute analysis focusing on taphonomic variables. The analysis was carried out by simple naked eye observations. After quantification of the results, taphonomic attributes were used for understanding each assemblage in the context of its accumulation history, as well as part of an inter-site comparison.

We focused on five variables (see Supplementary data 4 for further information about known causes of each taphonomic condition and examples).

Item breakage. Unintentional artifact breakage may occur due to rolling and saltation as coarse bedload during flood events, trampling by both anthropogenic and non-anthropogenic agents, and sediment compaction after coverage (McBrearty et al., 1998; Burroni et al., 2002; Hovers, 2003; Eren et al., 2010, 2011; Jennings, 2011). For our analysis, items were categorized as broken or complete.

General surface abrasion and edge damage. Small fractures and the general roughness of an artifact's lateral edges and the ridges on its flaked surfaces are related to mechanical movement in fluvial processes (Dibble et al., 1997; Burroni et al., 2002; Grosman et al., 2011; Hovers et al., 2014; Bustos-Pérez et al., 2019). The degrees of surface and ridge abrasion (fresh, partly abraded, and highly abraded) and of the item's edge damage (none, minimal, significant, and high) were assessed and categorized based on visual inspection.

Patination. In geomorphological studies, the term patina encompasses a varied group of rock-surface alterations compared to its archaeological use (Caux et al., 2018) and includes surface gloss (which we discuss separately). Here, we adopt the archaeological usage, by which 'patination' refers to changes in the color of flint, related to surface alterations caused by sediment chemistry, flint microscopic structure, humidity, and exposure to sun/light (Hurst and Kelly, 1961; Honea, 1964; Rottländer, 1975; Friedman et al., 1995; Howard, 1999, 2002; Burroni et al., 2002). Items were categorized into three conditions related to patination: existence of patina (patinated, not patinated, double patinated), type of coloring identified (desert varnish, white patina, white stains, black/gray coloring), and the patinated face of the artifact (one or both surfaces of the item).

<u>Gloss</u>. Glossy surfaces on flint artifacts, smooth to the touch, are a result of chemical processes and, to a lesser degree, mechanical weathering (Howard, 1999, 2002; Burroni et al., 2002). Items were categorized into three groups according to the degree of gloss apparent on the item surface (none, some, significant).

<u>Pitting.</u> Pitting refers to visible spherical pits in the size range from 2 mm to several cm in diameter that are created on artifact surfaces by thermal (often referred to as potlid scars/craters) and chemical processes and may cause the splitting and breakage of an artifact (Burroni et al., 2002; Knight and Zerboni, 2018, Yegorov et al., 2020). For this analysis items were categorized to three groups according to the number of pits apparent on the item surface (none, some, significant).

Statistical analysis

Chi-square (χ^2) tests were used to test meaningful patterning of taphonomy-related conditions within each assemblage $(\chi^2$ goodness of fit test) and for comparisons between them (χ^2) test of independence). The correlations between each site to the others were checked further using an ordinal correlation test, expressing the similarities or dissimilarities between the distribution of attribute states between each pair of sites in numeric values between -1 and 1. The tests were calculated using Excel, following the χ^2 formulas for the tests and using the excel data analysis correlation tool (see Supplementary data 5 for full information on the variables used and the calculations).

RESULTS

The results for each of the three studied sites are presented in this section, following the three analytical scales we employed.

Nahal Agev

Geological and geomorphological settings

The site of Nahal Aqev is situated within the gorge/canyon of the Aqev wadi, part of the Zin drainage system in the central Negev Highlands (Figs. 1, 2). The wadi fill consists of gravels and redeposited loess organized as a series of fluviatile terraces, accompanied by lateral colluvial deposits situated along the transition zone between the cliffs and the valley bottom. These terraces, which are mapped as Q2 and Q3 terraces, correspond to the Middle Pleistocene and Late Pleistocene deposition phases (Avni and Wieler, 2013; Avni et al., 2017, 2021).

Local context

The site is stratified within the abandoned colluvial body that corresponds with the Q2 terrace surface (Avni et al., 2021), deposited at an elevation of 430 m above mean sea level (amsl) along the western bank of the Ein Aqev Canyon. Eighteen sedimentological




Figure 2. The study area of Nahal Aqev. (a) Location of the site and the current Ein Aqev spring, situated 500 m downstream (view to north). (b) Old and new excavation areas and the geological trench (view to east). The full terrace section is 7 m deep. (c) Southern view of the geological units bearing archaeological horizons, color-shaded for clarity (Unit 7, light blue, 0.5 m scale bar placed on the section). (d) Geological map of the site's vicinity (Avni and Wieler, 2013).

units were identified in the terrace sequence, including three preserved archaeological horizons (Barzilai et al., 2022).

Unit 7 is 30 cm thick and contains abundant knapped flint items (Fig. 2c). Both field observations and technological analyses suggest that the lithic artifacts, horizontally embedded within the matrix, derive from an in situ horizon protected from subsequent erosion by rapid accumulation of fine-grained sediments within the Q2 terrace. There are no indications for sorting, and 60.5% of the retrieved items are <2 cm in size, which suggests minor post-depositional water disturbance (Schick, 1984; Dibble et al., 1997). The abundance of refitted items also suggests rapid burial of the lithics after their initial deposition (Villa, 1982; Dibble et al., 1997; Deschamps and Zilhão, 2018; Bertran et al., 2019).

Unit 7 was dated to 131 ± 23 ka by the post-infrared signal of alkali feldspars, which is consistent with its placement on a Q2 surface (Avni et al., 2021; Barzilai et al., 2022). The feldspars showed very little scatter in De values, indicating adequate bleaching of all grains prior to deposition.

Flint taphonomy

The lithic assemblage of unit 7 comprises 4133 artifacts, among which 1479 are larger than 2 cm and were subjected to a detailed attribute analysis, per practice for MP assemblages (Hovers, 2009). Of the latter, 23 are cores and 1456 are knapping products (debitage and tools, Table 1). A smaller sample (n = 224) was studied for gloss and pitting attributes. The taphonomic analysis did not reveal any significant differences between cores and flaking products (but note the small size of the core sample).

Flint items from Nahal Aqev show minimal surface abrasion, with 98% (n=1449) showing no abrasion and only 2% (n=30) abraded. More than half of the items are broken (57%, n=844). Edge damage is less frequent with 40% (n=530) showing minimal damage, 4% (n=51) having significant damage, and a single item with extensive edge damage. Most of items (71%, n=158) do not show gloss; with relatively low frequencies of moderate (27%, n=60) and significant (3%, n=6) gloss.

None of the items in the sample shows surface pitting (this may be a result of the smaller sample studied). Only 17% of the items (n = 251) in the total sample show patina, among which the dominant types are white stains (39%, n = 98) and black-gray color (38%, n = 96). An additional 22% (n = 55) bear white patina and only 2% (n = 3) show brown patination. Out of the patinated items, 94% (n = 237) bear patina on both faces. Only two items show different colorings on different flaking scars. None of the artifacts shows differential patination of distinct artifact faces.

The χ^2 tests for Nahal Aqev showed that the distribution is significantly different from the null hypothesis for all attributes (see Supplementary data 5 for test results). This suggests meaningful patterning of taphonomy-related conditions, which in turn can be linked to specific environmental conditions at the site.

Nahal Yitnan 7

Geological and geomorphological settings

The Nahal Yitnan 7 site is located at the northern part of a plateau of the Eocene Avedat Group rocks (Figs. 1, 3a, b). The plateau was probably overrun by fluvial deposits (sandstone and conglomerates) of the Early Miocene Hazeva Formation. The site is adjacent to the Nahal Yitnan basin, a tributary of Nahal Beersheva.

The Nahal Yitnan basin drains northwards along an ancient ~300-m-wide valley that runs along the contact between soft

chalk rocks of the Avedat Group and the hard limestone and dolomite rocks of the Judea Group (Fig. 3a, b). The valley was initially infilled with both in-situ and reworked Hazeva Formation clasts. The region was overlain by thick late Quaternary loess deposits that have been significantly eroded.

Today, the area is heavily grazed and tilled by local Bedouins, hence natural vegetation cover is sparse. In the Late Pleistocene, the region was likely covered by shrubs that stabilized the surface and affected soil formation and slopewash dynamics (Kijowska-Strugała et al., 2017).

Local context

The Nahal Yitnan 7 site is located at an elevation of ~450 m amsl on a slope of eroded Eocene chalk that is partially topped by a 20–60 cm-thick calcrete pre-dating loess deposition. Lithic artifacts were both scattered on the surface and buried along and within the southeastern gentle (5–10°) slope of a hill within the drainage area of a first-order tributary of the ephemeral Nahal Yitnan. The surface of the middle and lower parts of the slope exposes a ~40% cover of weathered and angular, cobble-size chalk fragments, and rounded fluvial pebbles on a calcic Loessial Serozem soil (Fig. 3a). The upper part of the slope comprises a 0–50 cm thick reworked loess deposit. A first-order wadi that drains the hillslope to the south has a thicker loess infill (Dan et al., 1976; Yair and Enzel, 1987).

The artifacts at Nahal Yitnan 7, which were systematically collected and excavated in two areas: B and C (Fig. 3; Yegorov, 2017), were assigned to the MP period on the basis of their typotechnological characteristics. To identify the spatial distribution of the artifacts and to see if they derived from buried assemblages, two ~1 m deep trenches (East trenches A and B, respectively; Figs. 4a, S1.2, S1.3) were dug along the slope of the hill. An additional trench (West trench, Figs. 4b, S1.4) was excavated in the first-order wadi. The artifact-bearing horizon was identified at different depths in all three trenches, where it was sampled for sediment analysis and OSL dating.

The trenches reveal distinct catenary changes in the stratigraphic sequence at the site (see Supplementary data 1 for full sedimentological analysis). Particle-size distribution of samples from most of the units shows a general pattern characteristic of primary loess (Crouvi et al., 2009) in the region (see full data in Supplementary data 3), mixed through slope-wash processes with fine-grained sand originating from the underlying Hazeva Formation. The sediments are highly calcic (46–64%) and resemble incipient calcrete, which can precipitate rapidly during intervals of decreased dust (loess) deposition, with contribution from the underlying weathered and eroded chalk deposits.

Approximately 20 cm-thick units of knapped flint artifacts, with differential preservation states, were identified, mainly in the Bk horizons (Fig. 4; Tables S1.1–3). Some artifacts were found in vertical positions. The presence of artifact-bearing sedimentological units in all the trenches demonstrates that lithics were distributed over an area of several hundred square meters. The spatial distribution of lithics on the surface is in accordance with the finds of artifacts in the upper units of the trench sections. Erosion of the loess along with modern, 30-cm-deep tilling (see Fig. 3a) partially contributed to the exposure of the artifacts.

OSL ages from the three stratigraphic sections range from 63 ± 5 ka to 30 ± 2 ka (Fig. 4; Table S2.1). The scatter within each sample, together with the reverse stratigraphic order in the West Trench (Fig. 4b), can be attributed to fluvial–colluvial processes

 Table 1. Breakdown of flint taphonomy attributes for the three assemblages

			Nahal Aqev 7				Nahal Yitnan 7				Giv'at Barne'a			
		Cores		Products		Cores		Products		Cores		Products		
		N	%	N	%	N	%	N	%	N	%	N	%	
Breakage	Not broken	15	65.2	620	42.6	31	93.9	54	46.6	61	79.2	31	46.3	
	Broken	8	34.8	836	57.4	2	6.1	62	53.4	10	13.0	36	53.7	
	Minimal breakage	0	0.0	0	0.0	0	0.0	0	0.0	6	7.8	0	0.0	
	Total	23	100.0	1456	100.0	33	100.0	116	100.0	77	100.0	67	100.0	
Surface Abrasion	Fresh	22	95.7	1427	98.0	12	36.4	67	57.8	25	32.5	24	35.8	
	Partly abraded	1	4.3	28	1.9	20	60.6	48	41.4	46	59.7	37	55.2	
	Highly abraded	0	0.0	1	0.1	1	3.0	1	0.9	6	7.8	6	9.0	
	Total	23	100.0	1456	100.0	33	100.0	116	100.0	77	100.0	67	100.0	
Edge Damage	No breakage	13	61.9	721	55.7	2	6.1	4	3.4	6	7.8	1	1.5	
	Minimal breakage	7	33.3	523	40.4	5	15.2	29	25.0	26	33.8	17	25.4	
	Significant breakage	1	4.8	50	3.9	20	60.6	58	50.0	29	37.7	37	55.2	
	High breakage	0	0.0	1	0.1	6	18.2	25	21.6	16	20.8	12	17.9	
	Total	21	100.0	1295	100.0	33	100.0	116	100.0	77	100.0	67	100.0	
Patina	Not patinated	21	91.3	1206	82.8	2	6.1	32	27.6	2	2.6	0	0.0	
	Patinated	1	4.3	249	17.1	31	93.9	83	71.6	75	97.4	66	98.5	
	Double patina	1	4.3	1	0.1	0	0.0	1	0.9	0	0.0	1	1.5	
	Total	23	100.0	1456	100.0	33	100.0	116	100.0	77	100.0	67	100.0	
Patina	White	0	0.0	55	22.0	22	71.0	50	59.5	15	20.0	18	26.9	
type	White stains	1	50.0	97	38.8	9	29.0	26	31.0	4	5.3	2	3.0	
	Desert varnish	1	50.0	2	0.8	0	0.0	6	7.1	56	74.7	47	70.1	
	Black/gray	0	0.0	96	38.4	0	0.0	2	2.4	0	0.0	0	0.0	
	Total	2	100.0	250	100.0	31	100.0	84	100.0	75	100.0	67	100.0	
Patina side	Flaking surface	1	50.0	3	1.2	2	6.5	8	9.5	7	9.3	1	1.5	
	Preparation surface	0	0.0	11	4.4	0	0.0	1	1.2	0	0.0	0	0.0	
	Both	1	50.0	236	94.4	29	93.5	75	89.3	68	90.7	66	98.5	
	Total	2	100.0	250	100.0	31	100.0	84	100.0	75	100.0	67	100.0	
Gloss	None	16	76.2	142	70.0	0	0.0	13	11.2	16	20.8	39	58.2	
	Some	5	23.8	55	27.1	6	18.2	22	19.0	40	51.9	19	28.4	
	Significant	0	0.0	6	3.0	27	81.8	81	69.8	21	27.3	9	13.4	
	Total	21	100.0	203	100.0	33	100.0	116	100.0	77	100.0	67	100.0	
Pitting	None	21	100.0	203	100.0	32	97.0	114	98.3	56	72.7	50	74.6	
	Some	0	0.0	0	0.0	1	3.0	2	1.7	14	18.2	16	23.9	
	Significant	0	0.0	0	0.0	0	0.0	0	0.0	7	9.1	1	1.5	
	Total	21	100.0	203	100.0	33	100.0	116	100.0	77	100.0	67	100.0	




Figure 3. Nahal Yitnan 7 site. (a) Southeasterly view of the slope with artifact scatter (~50 m long) and trench location (~20 m between East and West trenches). The valley of Nahal Yitnan is in the background. (b) Geological map of the site's vicinity (Roded, 1996; Wdowinski et al., 2012). The site's location is marked by the black and red diamond (c) Developed calcrete exposed uphill from the MP flint scatter (d) Excavation area C where horizon 3 bearing the artifacts was excavated. (e) Section of the excavation in area B showing horizons 4 and 3 with highly calcic loess at the bottom, under the artifact-bearing layer. Scale in (c-e) = 1 m.




Figure 4. The studied sections of the Nahal Yitnan 7 site in the East (a) and West (b) showing the main units: 1 = fractured, horizontally bedded chalk; 2 = calcrete (nari) surface; 3 = bk calcic loess with artifacts dispersed within upper part and with few clasts; 3a = bk1 calcic loess with matrix-supported clasts; 3b = bk2 calcic loess and clasts with artifacts dispersed within central part of unit; 3c = bk3 calcic loess loam with few clasts; 4 = brown tilled loess with dispersed lithics. OSL ages are in in ka. See Supplementary data 1 for full sediment analysis.

that had affected the slope and the sediments that were analyzed for dating (Porat et al., 2009; Oron et al., 2019). Most likely, the MP flint artifacts were originally embedded in loess deposited during the Late Pleistocene on top of the bedrock (Crouvi et al.,

2009). However, the slopes were continuously eroded by slopewash. Dynamic studies of slope wash in the semi-arid fringe of the Negev have shown that pebble- and larger-sized clasts are not transported downslope by slope wash, rather by livestock

activity and that clast angularity was not a significant factor (Ungar et al., 2010). However, these studies related to decadalscale rainfall recurrence. We suggest that the extreme rainfall events such as those with recurrences exceeding 10-1000 years generated several brief clast movements. The movement of eroded loess in such events probably contributed to local reposition and transport of clasts and angular to sub-angular artifacts. These erosional processes led to size-related differential dislocation of artifacts in the lithic assemblage and resulted in underrepresentation of the small (<2 cm) component, which may have been eroded downslope. Notably, the two most scattered OSL samples (with overdispersion values of 43-44%; OD in Table S2.1) contain populations of older grains with ages of 75-77 ka (Table S2.1), suggesting that these grains are remnants of a landscape originally covered with older loess. Thus, human activity predates 63 ± 5 ka, whereas the OSL ages represent continuous slope processes and loess mixing and not the initial deposition of the loess sediment.

Artifact transport by slope wash and redeposition on the slope is supported by sedimentology. While the grain size distribution of the > 2-mm fraction of the upper A horizon resembles loess, it is not a primary and pure loess deposit (as in Crouvi et al., 2009) but locally redeposited loess from slope wash (Table S1.4). The surface cover of thin loess mixed with a dense clast cover testifies to substantial stripping of the primary loess, which in turn explains the loess accumulations within the valleys of the adjacent drainage systems.

The high carbonate content (Table S1.4) of the artifact-rich layers appears to be post-depositional carbonate precipitation. During and following occupation, possibly after a period of erosion, aeolian loess deposition continued. The availability of calcium carbonate from adjacent and underlying bedrock, along with a 30% initial content of carbonates in the loess, led to development of a Bk horizon at the artifact level.

Flint taphonomy

Of the 149 Levallois items in the Nahal Yitnan 7 sample (Table 1), 33 are cores and 116 are knapping products (Levallois debitage and tools). There are no significant differences between cores

and products in the distribution of taphonomic variables, with the exception of surface abrasion (more prominent on cores) and breakage (slightly more frequent in the flake sample).

In general, the Nahal Yitnan 7 sample shows moderate frequencies of surface abrasion, with 53% (n = 89) without any ridge abrasion, 46% (n = 68) partially abraded, and only 1% (n = 2) heavily abraded. More than half of the items (57%, n = 85) are complete, but 96% (n = 143) present edge damage, of which 42% (n = 78) present 'intensive damage' and 21% (n = 31) show 'high damage' (the most intensive damage category). Most items show surface gloss, 72% (n = 108) with significant polish and 19% (n = 28) with a moderate amount. Surface pitting occurs on only 2% (n = 3) of the items.

Seventy-eight percent (n = 114) of the items are patinated. The dominant patina is white (63%, n = 72). An additional 30% (n = 35) bear white stains, 5% (n = 6) show brown patination, and 2% (n = 2) with black/gray patination. In 90% (n = 104) of the cases, patina is apparent on both sides of the item, but only one item bears two different types of patina (on different flaking scars on the same face of the item).

The χ^2 tests for Nahal Yitnan showed that the distribution is significantly different from the null hypothesis for all attributes apart from item breakage, where the distribution of breakage condition may be considered random (see Supplementary 5 for test results).

Giv'at Barne'a

Geological and geomorphic setting

The Giv'at Barne'a site is situated in the northwestern fringe of the central Negev Highlands, within the upper eastern reaches of the Wadi-Al-Arish basin of central and northern Sinai (Figs. 1, 5). The site, which is within the Ramat Barne'a plateau, is 10 km south of and below the Central Negev highlands. The plateau is composed of Middle Eocene chalky limestone interbedded with massive limestone units, forming the Matred/Nahal Yeter Formation. These units are primary runoff-generating rock formations (Wieler et al., 2016), suggesting that the slopes have been undergoing significant slopewash. Pliocene to Early Pleistocene conglomerates of the Ahuzam Formation deposited






Figure 5. Giv'at Barne'a area. (a) Slope from the upper area on top of the hill (right) to the lower area (left). (b) Geological map of the area near the site (marked with black/red diamond) within the Ramat Barne'a plateau (Zilberman and Avni, 2005). (c) Lower excavation area. (d) Cores on the surface.

along the drainage basins form prominent alluvial terraces rising \sim 20–30 m above the active channels (Zilberman and Avni, 2005). During this long period of surface exposure, the plateau underwent significant weathering, loess deposition, and erosion. In adjacent regions of the Negev, soil formation occurred mainly in topographic lows and colluvial environments (Amit et al., 2006; Enzel et al., 2008; Roskin et al., 2013). The soils on the stable carbonate surfaces are mainly brown lithosols and in the colluvial deposits mostly Loessial Serozems (Dan et al., 1970).

Local context

The Giv'at Barne'a site is situated on a small ($\sim 150 \times 100$ m) geomorphologically stable, flat-topped hill (Fig. S1.5). The upper slope comprises sub-horizontal, well-fractured, well-pitted chalky limestone layers in the form of ledges that contain flint nodules (5–50 cm in diameter). The ledges are separated by several meterwide horizontal intervals of flat bedrock. The middle and lower northern slope of the hill comprises softer limey chalk. The middle slope is partly covered by loess-dominated fill topped by cobble- to small boulder-size, pitted carbonate clasts, originating upslope.

The moderate (3–5°) lower part of the hillslopes has a colluvium-like morphology, with an unsorted mixture of pitted and angular carbonate pebbles and cobbles creating an incipient desert pavement-like cover. The degree of soil maturity and stability suggests that the soil is no older than Late Pleistocene in age (Table S1.5; Dan et al., 1982; Amit and Gerson, 1986).

Flint appears mainly as knapped items. Natural flint fragments appear in varying concentrations on the surface.

Scattered knapped items were collected from the top 5 cm of the surface in two selected areas of the Giv'at Barne'a site: one on the hilltop and the other at the northern colluvial slope of the hill, where the artifacts are part of an incipient desert pavement clast cover, whereby dust accumulated below the exposed clasts (Amit and Gerson, 1986). Based on their typo-technological characteristics, items in these assemblages belong to several archaeological periods, although there is no clear differentiation in patination.

We focused on the diagnostic Middle Paleolithic Levallois items. Some of the Levallois lithic artifacts in the hilltop area were found adjacent to each other and could be refitted, which indicates only restricted movement of material in this part of the site. Post-depositional movement of artifacts contributing to the desert pavement on the colluvial slope may have been minimal, given the low slope angle and lack of sorting by artifact size.

The sediment on the lower colluvial slope was dated by OSL to determine the age of dust buildup beneath the artifacts (Fig. 6;




Figure 6. OSL-dated section of the lower colluvial slope pedosediment (A and B soil horizons) at the Giv'at Barne'a site. Ages are in ka.

Fig. S1.6). The OSL samples show high over-dispersion values (43% and 49%; Table S2.2), indicating significant mixture or partial bleaching possibly due to downslope transport. OSL ages of 21 ± 2 ka (horizon A) and 32 ± 5 ka (B horizon) represent gradual Late Pleistocene accumulation of dust. The age range of the soil profile, $\sim 37-19$ ka, falls within pre-LGM times when there was significant loess deposition (Crouvi et al., 2009) and fixation (Kidron et al., 2014; Lucke and Bäumler, 2021) in the region. As a desert pavement develops in a stable arid environment, the OSL ages postdate the overlying lithics and provide a timeline for down-profile loess accumulation, perhaps during periods of relatively higher precipitation. The rate of deposition decreased in the Holocene when primary and secondary loess availability decreased (Avni et al., 2017) and the climate gradually aridified (Baruch and Goring-Morris, 1997; Robins et al., 2022; Vardi et al., 2023).

The hilltop flint outcrop adjacent to the excavated areas has been used as a raw material source. During flint extraction, blocks were removed from a limestone ledge at the upper and western part of the hillslope. Loess-like dust in cracks and underneath the displaced limestone blocks was also dated by OSL to identify secondary loess deposition post-dating the knapping activities. Three reworked loess samples in the rock fissures were dated to 1.6–0.4 ka (Fig. S1.7; Table S2.2). These young Late Holocene ages, which indicate erosion cycles and refill of the loess, exemplify the dynamic erosive character of the Middle–Late Holocene that inhibits loess fixation on hilltops (Lucke et al., 2019). The location of the studied quarry is on a western-facing slope that was exposed to incoming cyclonic rain fronts (Enzel et al., 2008), which may have enabled relatively intensive processes of erosion.

The hill most likely was covered with loess during parts of the Late Pleistocene because primary loess sections with ages ranging from >90 ka to ca. 14 ka are preserved on hilltops in the Negev highlands (Crouvi et al., 2008; Crouvi, 2009). The runoff-generating rock formations at the site led to loess erosion that slightly reorganized the spatial and vertical positions of the lithics. Because no older and overlying loess sediment was preserved on the upper parts of the hill, the time of site-use during the MP cannot be ascertained.

Flint taphonomy

The sample consists of 144 items attributed to the Levallois production system (Table 1). Of these, 77 are cores and 67 are knapping products (Levallois debitage and tools). Seventy-three items were collected from the hilltop surface and 71 from the colluvial slope. There are no significant differences between cores and products in the distribution of taphonomic variables, with the exception of breakage and of surface polish that are slightly more frequent in the debitage compared to cores (Table 1).

The Giv'at Barne'a MP artifacts show high frequencies of abrasion, with only 34% (n=49) showing no ridge abrasion, 58% (n=83) partially abraded, and 8% (n=12) heavily abraded. Most items are complete (64%, n=93), but 95% (n=137) show edge damage (46%, n=66 with significant damage and 19%, n=27 categorized as high damage).

Many items bear surface gloss, 41% (n = 59) showing a moderate amount and 21% (n = 30) with significant polish. Twenty-six percent (n = 38) of the items in the sample show also pitting on the surface. Almost all items are patinated (98%, n = 142) and the dominant patina is brown (72%, n = 103), with 23% (n = 3 3) having white patina and 4% (n = 6) showing white stains. Patina is apparent on both sides of the item in 95% (n = 133) of

the items. Only one item bears two different types of patina (on different flaking scars on the same face of the item).

Several differences are observed between the lithics of the two areas. Forty percent (n = 15) of the items in the colluvial slope show no ridge abrasion, as opposed to only 28% (n = 9) at the hill-top. The lower area is also characterized by more items with white coloring (34%, n = 17), as opposed to only 7% (n = 2) on the hilltop. For the other taphonomic attributes, no differences were observed between the two areas. The χ^2 tests for Giva't Barne'a showed that the distribution is significantly different from the null hypothesis for all attributes (see Supplementary data 5 for test results).

Inter-assemblage comparison of flint taphonomy

Inter-assemblage differences were tested statistically using χ^2 tests under the null hypothesis of similar distribution of taphonomic variables among the three sites. The results show that the three assemblages differ significantly (in statistical terms) with regards item breakage, edge damage, and patina types. For the other variables, the sample size was not large enough to obtain valid results, so they were tested again combining some of the categories, still following questions of differential preservation between sites (see Supplementary data 5 for full tests results).



Figure 7. Taphonomy variables on core and knapping products: (a) surface abrasion, (b) item breakage, (c) edge damage.

The three assemblages differ in the distribution of preservation states (Fig. 7). Giva't Barne'a appears to have undergone more artifact surface abrasion in general, as indicated by higher frequencies of abraded items and of highly abraded pieces in particular, while Nahal Aqev items are mostly unabraded; Nahal Yitnan presents more variable results. These differences are expressed in the correlation test results, showing high correlation between Giva't Barne'a and Nahal Yitnan and a weak correlation between Giva't Barne'a and Nahal Agev. Broken items are somewhat more frequent in Nahal Aqev compared to the two other sites, yet there is a fairly strong correlation between the sites, with the strongest correlation between Giv'at Barne'a and Nahal Yitnan (Fig. 7b; Supplementary data 5). Edge-damage counts show similar results to surface abrasion (Fig. 7c; Supplementary data 5), Nahal Aqev being distinguished from the others by higher frequencies of artifacts without edge damage, while Giv'at Barne'a and Nahal Yitnan show a high correlation value.

Some degree of artifact surface gloss was detected in all three assemblages (Fig. 8a). In comparison to the two other case studies, gloss in the Nahal Yitnan assemblage is more frequent, more intensive, and it appears on all cores and on most knapping products. The correlation between Nahal Aqev and Giva't Barne'a is positive, whereas it is strongly negative between these two sites and Nahal Yitnan.

Pitting is rare overall in the three assemblages (Fig. 8b). It is more frequent in Giv'at Barne'a while completely absent from the Nahal Aqev assemblage, but these differences are not statistically valid due to the very low frequencies.

The distributions of patination states in the assemblages track the patterns of surface abrasion and edge damage (Fig. 9a). Nahal Aqev items are less patinated, whereas all items in Giv'at Barne'a and most items in Nahal Yitnan are patinated. Accordingly, there is a high correlation value between Giv'at Barne'a and Nahal Yitnan and very low correlation values between each of these sites and Nahal Aqev.

The distributions of patina types, however, show a more complex pattern (Fig. 9b). The most dominant patina type differs from one assemblage to the other as manifested in a very weak statistical correlation between Nahal Aqev and Nahal Yitnan and negative correlation values for Giva't Barne'a and the other sites. In Nahal Aqev, white stains and black/gray patina are common on the knapping products but do not appear on the cores. The white patina and desert varnish are most frequent in all the products in the assemblages from Nahal Yitnan and Giv'at Barne'a, respectively. In all three assemblages, patination always appears on both faces of the items (Fig. 9c) with the same coloring. Indeed, distribution of this specific taphonomic variable does not differ statistically among the assemblages and it shows strong correlation between all sites. In the few cases where patination appears on only one face of the core, it is on the flaking surface.

The corelation test results are consistent with the understanding that the differences do not conform to a unified pattern. While all




Figure 8. Taphonomy variables on core and knapping products: (a) surface gloss, (b) pitting.




Figure 9. Coloring variables on core and knapping products: (a) patination, (b) Type of patina, (c) patination side on each item.

three sites show some similarity in item breakage, pitting, and patination side and differ in the distribution of patina types, some attributes show stronger similarities between varying pairing of the sites. Specifically, Nahal Yitnan and Giv'at Barne'a show stronger corelations for surface abrasion, edge damage, and patination, while Nahal Aqev and Giv'at Barne'a show a stronger correlation due to the lower frequencies of gloss compared to Nahal Yitnan.

In summary, the results of the flint taphonomy analysis show that the three sites differ from one another in most variables. Importantly, all sites are located within a limestone-dominated bedrock in areas that were strongly affected by Late Pleistocene loess accumulation and erosion. We suggest that differences in

site accumulation histories and post-depositional processes are manifested in the flint taphonomy. The combination of the results of the different scales of analysis and tying geomorphological processes affecting accumulation and preservation at each site with the taphonomy of the lithic assemblage, may also help work in reverse and relate specific observed taphonomic variables to specific regional or local geomorphic processes (see discussion).

DISCUSSION

The geomorphic, geochronological, and sedimentological analyses, at regional and local scales suggest three scenarios of burial and exposure histories for the three MP assemblages that serve as case studies. Despite some recognized intra- and inter-exposure variations in raw material properties, their common Eocene origin suggests that knapped items are less susceptible to taphonomic effects stemming from differences in raw material. This enables focus on the potential environmental drivers of modification and transformation.

Nahal Aqev represents a rapid accumulation of the archaeological horizon, followed by rapid burial shortly after anthropogenic activity. This is a diametrically different depositional history from the two other sites. At Giv'at Barne'a, the context is that of a long, almost continuous process of surface exposure, where reworked loess gradually accumulated beneath the artifacts. The depositional context of Nahal Yitnan 7 is complex, indicating a last glacial history of loess accumulation followed by episodes of exposure (of different degrees) due to active slope processes and erosion.

Several aspects of the analyses of lithic taphonomy seem to tie in with the geomorphic processes and may be helpful in backtracking sedimentation cover and local fluvial processes. Taphonomic aspects such as surface abrasion of items, edge damage, pitting, and patination seem to co-vary with exposure duration. On the other hand, attributes such as artifact surface gloss are potentially indicators of localized slopewash, therefore sedimentological processes (represented in this study by the Nahal Yitnan case study) and are not necessarily attributed to long exposure of artifacts on the surface. Edge damage can certainly be a result of trampling during human occupation and, possibly, due to the use of the items. The much lower frequencies of edge damage in Nahal Aqev compared to the other two sites may suggest that long exposure and slope processes have more effect on edge damage than human behavior.

According to previous studies, artifact breakage is affected by the nature of sediments and cycles of coverage and exposure, trampling (by both human and non-human agents) when fully or partially exposed on the surface, and sediment compaction (see also Supplementary data 4), and thus may depict a more complex picture compared to other taphonomic variables. Nahal Aqev showed slightly higher breakage frequencies than the other sites, maybe due to burning and sediment compaction under a 3-4 m burial depth. The breakage at Giv'at Barne'a probably was due mainly to trampling and exposure to the sun since the artifacts were at the surface, while at Nahal Yitnan fluvial and slope processes are the more likely causes. Despite their different accumulation processes, the three analyzed assemblages show similar distributions of breakage, with nearly equal frequencies of broken items. This equifinality of anthropogenic and natural processes is of importance, because it clarifies that this taphonomic variable is a less-reliable proxy of depositional history.

The diversity and complexity of coloring documented in this study and the various inferred processes align with the observations from previous studies (e.g., Hurst and Kelly, 1961; Honea, 1964; Rottländer, 1975; Friedman et al., 1995; Howard, 1999, 2002; Burroni et al., 2002; Goldsmith et al., 2012, 2014; Dorn, 2013; Caux et al., 2018). Our results reaffirm the conclusion that patination should not be considered an a priori indication for higher levels of post-depositional damage or disturbance, and thus a less well-preserved site. The correlation between patina and preservation or exposure versus sediment coverage is not straightforward. Importantly, it is the type of patina (rather than its existence per se) that speaks to the natural or anthropogenic formation processes, both during and after site accumulation. In the arid environment of the Negev desert, coloring of

flint surfaces sometimes points to long exposure on the landscape, leading mostly to the formation of desert varnish. In other cases, coloring is related to formation of the archaeological assemblages in a geomorphologically active environment, where white coloring, which is related to alkaline carbonate-rich sediment, will be more frequent at the expense of desert varnish. As in Nahal Yitnan 7, white patina is expected to occur with abrasion and gloss from the slope and fluvial activities. When the white coloring is associated with low rates of abrasion and edge damage, it may likely indicate sediment coverage of a well-preserved horizon, and not necessarily exposure.

Black and gray coloring is strongly associated with direct exposure to fire, and in most cases was caused by human activity. Unlike controlled heating, the exposure of flint item to direct fire probably is an unintentional side-effect of the use of fire on-site, similar to breakage or edge damage in some cases. Black coloring can be an indicator for fire-related activity on site, even when hearths are not recognized. On the other hand, the absence or low frequencies of black and gray coloring from sites such as Giva't Barne'a and Nahal Yitnan may result from re-patination due to exposure on the surface or chemical processes, and therefore cannot disprove fire-related activities.

The links between lithic taphonomy indicators and geomorphic processes are summarized in Table 2. Bearing in mind the complexity and variety of processes that can influence each attribute, as well as the non-linear accumulation of some of them, we merely point out the directionality (increase/decrease) for the processes recognized in the analyzed sites based on the differences observed and the correlation tests.

Another issue arising from our data pertains to the use of luminescence dating in OAS in arid environments. The current study highlights some limitations but also the potential of the method as a tool in understanding site-formation processes. Again, this is related to the different burial and exposure histories for the MP archaeological artifacts in the sample sites. In Nahal Aqev, a dated sedimentological sequence provides the full chronological framework for human occupation. On the other hand, OSL dating at Giv'at Barne'a could not contribute to identifying the timing of human activity. No sediment was deposited with and above the artifacts; dust only accumulated under the artifacts, yet based on their typo-technological characteristics, this accumulation postdates the items. Finally, in the more complex accumulation scenario of Nahal Yitnan, OSL ages, not directly applicable to the time of the human occupation, provide a minimal age for occupation and help explain and understand redeposition processes of artifacts.

How information is mined from OAS with different degrees of integrity speaks to the frame of mind of researchers as much as it does to the quality of the information gleaned from the sites themselves. Archaeologists typically prefer to study the bestpreserved sites (Nahal Aqev in our sample), because these require simpler explanatory scenarios and provide more reliable information about human activities and behaviors. Perhaps counterintuitively, sites such as Giv'at Barne'a allow an approach based on parsimony. Because site history presents a more linear trend of long-term exposure, there is less variability of formation process. This in turn calls for fewer steps and assumptions in analyzing the sites and the lithic remains in them. In particular cases (e.g., when flat areas form part of the site), assemblages might be more complete—to the degree that items can be refitted and will offer reliable information about aspects such as lithic technology and activity areas, even if the time frame remains

Table 2. Summary of the links between lithic taphonomy and observed geomorphic processes. Empty cells represent cases where cause/effect cannot be determined based on our sample.

	Surface exposure	Human activity during occupation	Fluvial processes	Sediment cover
Breakage	Increase	Increase	Increase	Increase
Surface abrasion	Decrease	_	Decrease	_
Edge damage	Increase	Increase	Increase	_
Patina	Increase	Increase**	_	Increase
White	Decrease*	_	_	Increase*
Desert varnish	Increase	_	_	Decrease
Black/gray	_	Increase	_	_
Gloss	Increase	_	Increase	Increase*
Pitting	Increase	Increase**	_	_

^{*}Depending on sediment chemistry; **Unevenly in specific areas of the site, due to burning.

vague. In such cases, the occurrence of refits, combined with their lateral and vertical distributions can be used as a quantitative indicator for the intensity of post-depositional processes (Cziesla, 1990a,b). Otherwise, it is conceptually easier to limit analyses of such assemblages to very general conclusions (e.g., "hominins were active on this landscape at some point or points in time during the MP"), or to inflate the inferential significance of the archaeological contents of the sites because it is difficult to test empirically the underlying hypotheses.

Our analysis suggests that 'intermediate' sites such as Nahal Yitnan 7, with cycles of accumulation and exposure, may contribute environmental information and dating constraints that are missing from long-exposure sites, and can be used in a nuanced manner to infer broader trends in the human occupation of a region. While refitting success is unlikely in such contexts and the assemblages cannot be used, for example, in spatial analyses, the assemblages can be mined for technological information (see Hovers et al., 2014), timing and length of the occupation, and be discussed in the context of broader regional schemes.

In the current case study, the whole assemblage of Nahal Aqev can be included in the technological and typological analysis, can be assigned confidently to the MP, and its OSL dating places it within ca. 130 ka or older (Barzilai et al., 2022). In Giv'at Barne'a, one may recognize and discuss the nature of the site as a workshop, but due to the mixture with knapping products from several periods, we cannot include non-Levallois items in the analysis and reconstruct the whole reduction sequence. In contrast, at the site of Nahal Yitnan, despite its complex depositional history, one can identify the better-preserved areas and include all lithic items in the technological analysis (e.g., technologically non-diagnostic items), such that the emerging understanding of the characteristics of the assemblage is based on a reliable dataset. While taking into consideration that the site probably represents more than one MP occupation event and may not contain a complete assemblage, it is possible to rule out mixture with other periods based on the taphonomic characteristics. Moreover, due to the geomorphic reconstruction of the loess cover-erosion cycles, the site can be tied to locally and




Figure 10. A schematic illustration for the conceptual model.

regionally dated sequences and the temporal placement of the assemblage can be further constrained.

Integration of the three levels of analysis (i.e., examining the associations among regional, local, and site-specific depositional histories) allowed us to clarify the potential role of each such scale in understanding arid region OAS, in which lithic artifacts are the most common, and often the only, indications of human presence on the landscape. Such insights for the Negev MP are significant because technological traditions in this area have been argued to be linked to those observed in several neighboring areas (e.g., Goder-Goldberger et al., 2016; Barzilai et al., 2022). The Levantine MP has often been characterized through a three-phase sequence of technological changes, as evinced in the long sequence of Tabun Cave (e.g., Bar-Yosef, 1998). More recently, it was argued that the tri-partite Tabun model of technological change may more accurately describe MP technological variability in the Mediterranean ecological zone, but is less compelling for the arid areas because of the different ecological affordances and settlement systems in such areas (Hovers, 2009). Clarifying the temporal span and integrity of the assemblages from OAS in the arid zone can now address such questions with better-informed empirical data.

Our work also allows us to construct a conceptual model for arid region OAS (Fig. 10), drawing attention to the variety of potential relationships among environment, human behavior, and the archaeological record in arid conditions. The use of these artifacts as indicators for environmental effects has great value. At the site level, it allows us to assess accumulation history, but for regional research it can also be used as a tool for coherent integration of data from different sites into one data set. The understanding of where each site is placed within the range of exposure and preservation possibilities in the specific region allows, in turn, augmentation of the regional archaeological data base by using informed and critically evaluated data. Thus, we can mine the regional data base for information that otherwise would be overlooked or considered unreliable, or would be considered as introducing undesirable 'noise,' rank them differentially and choose the relevant assemblage from each site. The three model sites presented in this paper are part of a larger sample of 11 MP assemblages from the Negev, all analyzed in a similar way. In turn, the way that each one of the project assemblages can contribute to the regional research is assessed and the data is integrated and used to refine our knowledge of the human presence in the Negev during the MP.

Finally, we suggest that our conceptual model bears implications for areas beyond the Negev desert, for example Arabia (Hilbert et al., 2016; Groucutt et al., 2021) and north and northeastern Africa (Olszewski et al., 2010; Foley and Lahr, 2015), where much cultural material is known from surface collections. The kind of analysis that we used in our 'proof of concept' study may help in obtaining more information and in reaching a more informed evaluation of such records. Such additional insights may bear on the potential contribution of similar records to the ongoing discussion of Late Pleistocene dispersals and contacts among modern human populations and their contemporaries in arid areas, in which such regional records play a role.

Supplementary Material. The supplementary material for this article can be found at https://doi.org/10.1017/qua.2023.31

Acknowledgments. The excavations in Nahal Yitnan 7 and Giv'at Barne'a were conducted on behalf of the Israel Antiquities Authority, which granted

us permission to use all figures of the excavations and finds from these sites. Artifacts were photographed by C. Amit and D. Gazit. The authors thank O. Barzilai and E. Boaretto for allowing the use of unpublished data from Nahal Aqev. Sedimentological analysis was conducted at the Sedimentology Laboratories at the University of Haifa. Maya Oron's PhD project has been supported by the IAA scholarship program and a research grant from the Ruth Amiran Fund for archaeological research in Israel, awarded by The Institute of Archaeology, The Hebrew University of Jerusalem.

Competing Interests. The authors declare no competing interests.

REFERENCES

- Alexandrov, Y., Laronne, J.B., Reid, I., 2003. Suspended sediment transport in flash floods of the semiarid northern Negev, Israel. *International Association of Hydrological Sciences*, Publication 278, 346–352.
- Amit, R., Enzel, Y., Sharon, D., 2006. Permanent Quaternary hyperaridity in the Negev, Israel, resulting from regional tectonics blocking Mediterranean frontal systems. *Geology* 34, 509–512.
- Amit, R., Gerson, R., 1986. The evolution of Holocene reg (gravelly) soils in deserts: an example from the Dead Sea region. *CATENA* 13, 59–79.
- Amit, R., Simhai, O., Ayalon, A., Enzel, Y., Matmon, A., Crouvi, O., Porat, N., McDonald, E., 2011. Transition from arid to hyper-arid environment in the southern Levant deserts as recorded by early Pleistocene cummulic Aridisols. *Quaternary Science Reviews* 30, 312–323.
- Avni, Y., 2017. Tectonic setting and physiography setting of the Levant. In: Enzel, Y., Bar-Yosef, O. (Eds.), Quaternary of the Levant (Environment, Climate Change and Humans). Cambridge University Press, Cambridge, UK; New York, NY, pp. 3–16.
- Avni, Y., Faershtein, G., Porat, N., 2017. Studies of stream terraces in the Negev Highlands and their relationship with the Levant alluvial chronologist. In: Enzel, Y., Bar-Yosef, O. (Eds.), Quaternary of the Levant (Environment, Climate Change and Humans). Cambridge University Press, Cambridge, UK, New York, pp. 457–469.
- Avni, Y., Oron, M., Cohen-Sasson, E., Porat, N., Barzilai, O., 2021. Chrono-sequences of alluvial terraces and fossilized water bodies as a predictive model for detecting Lower and Middle Palaeolithic sites in the Negev desert, Israel. *Quaternary Science Reviews* 268, 107114. https://doi.org/10.1016/j.quascirev.2021.107114.
- Avni, Y., Segev, A., Ginat, H., 2012. Oligocene regional denudation of the northern Afar dome: pre- and syn-breakup stages of the Afro-Arabian plate. *The Geological Society of America Bulletin* 124, 1871–1897.
- Avni, Y., Wieler, N., 2013. Geological Map of Israel (1:50,000): Sede Boqer Sheet (18-IV). The Geological Survey of Israel, Jerusalem, Israel.
- Bailey, G., 2007. Time perspectives, palimpsests and the archaeology of time. Journal of Anthropological Archaeology 26, 198–223.
- Bar-Yosef, O., 2006. Between observations and models. In: Hovers, E., Kuhn, S.L. (Eds.), Transitions Before The Transition: Evolution and Stability in the Middle Paleolithic and Middle Stone Age. Springer, New York, pp. 305–325.
- Bar-Yosef, O., 1998. The chronology of the Middle Paleolithic in the Levant. In: Akazawa, T., Aoki, K., Bar-Yosef, O. (Eds.), Neandertals and Modern Humans in Western Asia. Plenum Press, New York, pp. 39–56.
- Bar-Yosef, O., Goren, N., 1980. Afterthoughts following prehistoric surveys in the Levant. *Israel Exploration Journal* 30, 1–16.
- Baruch, U., Goring-Morris, N., 1997. The arboreal vegetation of the Central Negev Highlands, Israel, at the end of the Pleistocene: evidence from archaeological charred wood remains. Vegetation History and Archaeobotany 6, 249–259.
- Barzilai, O., Oron, M., Porat, N., White, D., Timms, R., Blockley, S., Zular, A., et al., 2022. Expansion of eastern Mediterranean Middle Paleolithic into the desert region in early marine isotopic stage 5. Scientific Reports 12, 4466. https://doi.org/10.1038/s41598-022-08296-9.
- Ben-Israel, M., Enzel, Y., Amit, R., Erel, Y., 2015. Provenance of the various grain-size fractions in the Negev loess and potential changes in major dust sources to the Eastern Mediterranean. *Quaternary Research* 83, 105–115.
- Benjamini, C., 1979. The Nahal Yeter Formation—a new Eocene formation from the Northwestern Negev, Israel. *Israel Journal of Earth Sciences* 28, 164–166.

- Bertran, P., Todisco, D., Bordes, J.-G., Discamps, E., Vallin, L., 2019. Perturbation assessment in archaeological sites as part of the taphonomic study: a review of methods used to document the impact of natural processes on site formation and archaeological interpretations. *PALEO. Revue d'Archéologie Préhistorique* 30, 52–75.
- Boëda, E., 1995. Levallois: a volumetric construction, methods, a technique. In: Dibble H.L., Bar-Yosef, O. (Eds.), The Definition and Interpretation of Levallois Technology, Monographs in World Prehistory. Prehistory Press, Madison, Wisconsin, pp. 41–65.
- Bowman, D., Karnieli, A., Issar, A., Bruins, H.J., 1986. Residual colluvio-aeolian aprons in the Negev Highlands (Israel) as palaeo-climatic indicator. *Palaeogeography, Palaeoclimatology, Palaeoecology* **56**, 89–101.
- Braun, M., 1967. Type section of 'Avedat Group, Eocene, Formation in the Negev, southern Israel. Stratigraphic Sections 4. Geological Survey of Israel.
- Bruins, H.J., Yaalon, D.H., 1980. Stratigraphy of the Netivot section in the desert loess of the Negev (Israel). In: Pesci, M. (Ed.), Studies on Loess: Papers Published in Co-Operation of the INQUA Loess Commission Together with the UNESCO IGCP Magnetostratigraphic Project 128. Akademiai Kiado, Budapest, pp. 161–169.
- Burroni, D., Donahue, R.E., Pollard, A.M., Mussi, M., 2002. The surface alteration features of flint artefacts as a record of environmental processes. *Journal of Archaeological Science* 29, 1277–1287.
- Burrough, S.L., Thomas, D.S., Allin, J.R., Coulson, S.D., Mothulatshipi, S.M., Nash, D.J., Staurset, S., 2022. Lessons from a lakebed: unpicking hydrological change and early human landscape use in the Makgadikgadi basin, Botswana. *Quaternary Science Reviews* 291, 107662. https://doi.org/10.1016/j.quascirev.2022.107662.
- Bustos-Pérez, G., Díaz, S., Baena, J., 2019. An experimental approach to degrees of rounding among lithic artifacts. *Journal of Archaeological Method and Theory* 26, 1243–1275.
- Calvo, R., Bartov, Y., 2001. Hazeva Group, southern Israel: new observations, and their implications for its stratigraphy, paleogeography, and tectonosedimentary regime. *Israel Journal of Earth Sciences* 50, 71–100.
- Cancellieri, E., di Lernia, S., 2013. Middle Stone Age human occupation and dispersals in the Messak plateau (SW Libya, central Sahara). *Quaternary International* 300, 142–152.
- Caux, S., Galland, A., Queffelec, A., Bordes, J.G., 2018. Aspects and characterization of chert alteration in an archaeological context: a qualitative to quantitative pilot study. *Journal of Archaeological Science: Reports* 20, 210–219.
- Chiotti, L., Dibble, H.L., Olszewski, D.I., McPherron, S.P., Schurmans, U.A., 2009. Middle Palaeolithic lithic technology from the western High Desert of Egypt. *Journal of Field Archaeology* 34, 307–318.
- Chiotti, L., Olszewski, D.I., Dibble, H.L., McPherron, S.R., Schurmans, U., Smith, J.R., 2007. Paleolithic Abydos: reconstructing individual behaviors across the high desert landscape. In: Hawass, Z., Richards, J.E. (Eds.), The Archaeology and Art of Ancient Egypt: Essays in Honor of David B. O'Connor. Supreme Council of Antiquities Press (distributed by American University in Cairo Press), Cairo, pp. 169–183.
- Coco, E., Holdaway, S., Iovita, R., 2020. The effects of secondary recycling on the technological character of lithic assemblages. *Journal of Paleolithic Archaeology* 3, 453–474.
- Coco, E., Iovita, R., 2020. Time-dependent taphonomic site loss leads to spatial averaging: implications for archaeological cultures. *Humanities and Social Sciences Communications* 7, 136. https://doi.org/10.1057/s41599-020-00635-3.
- Crouvi, O., Amit, R., Enzel, Y., Porat, N., Sandler, A., 2008. Sand dunes as a major proximal dust source for late Pleistocene loess in the Negev desert, Israel. *Quaternary Research* 70, 275–282.
- Crouvi, O., Amit, R., Porat, N., Gillespie, A.R., McDonald, E.V., Enzel, Y., 2009. Significance of primary hilltop loess in reconstructing dust chronology, accretion rates, and sources: an example from the Negev desert, Israel. *Journal of Geophysical Research: Earth Surface* 114, F02017. https://doi.org/10.1029/2008JF001083.
- Cziesla, E., 1990a. On refitting of stone artefacts. In: Cziesla, E., Eickhoff, S., Arts, N., Winter, D. (Eds.), The Big Puzzle: International Symposium on Refitting Stone Artefacts, Studies in Modern Archaeology. Holos, Bonn, Germany, pp. 9–44.

Cziesla, E., 1990b. Artefact production and spatial distribution on the open air site 80/14 (Western Desert, Egypt). In: Cziesla, E., Eickhoff, S., Arts, N., Winter, D. (Eds.), The Big Puzzle: International Symposium on Refitting Stone Artefacts, Studies in Modern Archaeology. Holos, Bonn, Germany, pp. 583–610.

- Dan, J., Raz, Z., 1970. The Soil Association Map of Israel (scale 1:250,000).Volcani Institute of Agricultural Research, Bet Dagan.
- Dan, J., Yaalon, D.H., Koyumdjiski, H., Raz, Z., 1970. The soil association map of Israel (1:1,000 000). Israel Journal of Earth Sciences 21, 29–49.
- Dan, J., Yaalon, D.H., Koyumdjiski, H., Raz, Z., 1976. The Soils of Israel (with 1:500,000 map). Volcani Center, Bet-Dagan Pamphlet 159, 30 pp.
- Dan, J., Yaalon, D.H., Moshe, R., Nissim, S., 1982. Evolution of Reg soils in southern Israel and Sinai. *Geoderma* 28, 173–202.
- Deschamps, M., Zilhão, J., 2018. Assessing site formation and assemblage integrity through stone tool refitting at Gruta da Oliveira (Almonda karst system, Torres Novas, Portugal): a Middle Paleolithic case study. PLoS One 13, e0192423. https://doi.org/10.1371/journal.pone.0192423.
- Dibble, H.L., Chase, P.G., McPherron, S.P., Tuffreau, A., 1997. Testing the reality of a "living floor" with archaeological data. *American Antiquity* 62, 629–651.
- Dorn, R.I., 2013. Rock coatings. In: Pope, G.A. (Ed.), Treatise on Geomorphology. Academic Press, San Diego, pp. 70–97.
- **Douglass, M.J., Wandsnider, L., Holdaway, S.J.**, 2023. Surface artifact scatters, data collection, and significance: case studies from Australia and the United States. *Advances in Archaeological Practice* **11**, 29–41.
- Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B., Sharon, D., 2008. The climatic and physiographic controls of the eastern Mediterranean over the Late Pleistocene climates in the southern Levant and its neighboring deserts. Global and Planetary Change 60, 165–192.
- Enzel, Y., Bar-Yosef, O., 2017. Quaternary of the Levant: Environments, Climate Change, and Humans. Cambridge University Press, Cambridge, UK.
- Eren, M.I., Boehm, A.R., Morgan, B.M., Anderson, R., Andrews, B., 2011.
 Flaked stone taphonomy: a controlled experimental study of the effects of sediment consolidation on flake edge morphology. *Journal of Taphonomy* 9, 201–217.
- Eren, M.I., Durant, A., Neudorf, C., Haslam, M., Shipton, C., Bora, J., Korisettar, R., Petraglia, M., 2010. Experimental examination of animal trampling effects on artifact movement in dry and water saturated substrates: a test case from South India. *Journal of Archaeological Science* 37, 3010–3021.
- Eyal, Y., Reches, Z., 1983. Tectonic analysis of the Dead Sea Rift region since the Late-Cretaceous based on mesostructures. *Tectonics* 2, 167–185.
- Faershtein, G., Porat, N., Avni, Y., Matmon, A., 2016. Aggradation-incision transition in arid environments at the end of the Pleistocene: an example from the Negev Highlands, southern Israel. *Geomorphology* 253, 289–304.
- Foley, R., 1981. Off-site archaeology: an alternative approach for the short sited. In: Hodder, I., Isaac, G.L., Hammond, N. (Eds.), Pattern of the Past: Studies in the Honour of David Clarke. Cambridge University Press, Cambridge, UK, pp. 157–184.
- Foley, R.A., Lahr, M.M., 2015. Lithic landscapes: Early human impact from stone tool production on the central Saharan environment. *PloS ONE* 10, e0116482. https://doi.org/10.1371/journal.pone.0116482.
- Friedmann, E., Goren-Inbar, N., Rosenfeld, A., Marder, O., Burian, F., 1995. Hafting during Mousterian times—further indication. *Journal of the Israel Prehistoric Society* 26, 8–31.
- Galbraith, R.F., Roberts, R.G., 2012. Statistical aspects of equivalent dose and error calculation and display in OSL dating: an overview and some recommendations. *Quaternary Geochronology* 11, 1–27.
- Goder-Goldberger, M., 2020. The Middle to Upper Palaeolithic transition as seen from Far'ah II and Boker Tachtit, Israel. In: Leplongeon, A., Goder-Goldberger, M., Pleurdeau, D. (Eds.), Not Just a Corridor: Human Occupation on the Nile Valley and Neighbouring Regions Between 75,000 and 15,000 Years Ago. Publications Scientifiques du Muséum National d'Histoire Naturelle, Paris, pp. 221–237.
- Goder-Goldberger, M., Gubenko, N., Hovers, E., 2016. "Diffusion with modifications": Nubian assemblages in the central Negev highlands of Israel and their implications for Middle Paleolithic inter-regional interactions. *Quaternary International* 408, 121–139.

- Goldberg, P., 1986. Late Quaternary environmental history of the southern Levant. *Geoarchaeology* 1, 225–244.
- Goldberg, P., 1976. Upper Pleistocene geology of the Avdat/Aqev area. In: Marks, A.E. (Ed.), Prehistory and Paleoenvironments of the Central Negev, Israel, Vol. I, The Avdat/Aqev Area. Part 1. Southern Methodist University Press, Dallas, Texas, pp. 25–56.
- Goldberg, P., Brimer, B., 1983. Late Pleistocene geomorphic surfaces and environmental history of the Avdat/Havarim area, Nahal Zin. In: Marks, A.E. (Ed.), Prehistory and Paleoenvironments of the Central Negev, Israel, Vol III, The Avdat/Aqev Area. Part 3. Southern Methodist University (SMU) Press, Dallas, Texas, pp. 1–13.
- Goldsmith, Y., Enzel, Y., Stein, M., 2012. Systematic Mn fluctuations in laminated rock varnish developed on coeval early Holocene flint artifacts along a climatic transect, Negev desert, Israel. *Quaternary Research* 78, 474–485.
- Goldsmith, Y., Stein, M., Enzel, Y., 2014. From dust to varnish: geochemical constraints on rock varnish formation in the Negev desert, Israel. Geochimica et Cosmochimica Acta 126, 97–111.
- Goren-Inbar, N., 2011. Culture and cognition in the Acheulian industry: a case study from Gesher Benot Ya'aqov. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 1038–1049.
- Grosman, L., Sharon, G., Goldman-Neuman, T., Smikt, O., Smilansky, U., 2011. Studying post depositional damage on Acheulian bifaces using 3-D scanning. *Journal of Human Evolution* 60, 398–406.
- Groucutt, H.S., Scerri, E.M., Amor, K., Shipton, C., Jennings, R.P., Parton, A., Clark-Balzan, L., Alsharekh, A., Petraglia, M.D., 2017. Middle Palaeolithic raw material procurement and early stage reduction at Jubbah, Saudi Arabia. Archaeological Research in Asia 9, 44–62.
- Groucutt, H.S., White, T.S., Scerri, E.M., Andrieux, E., Clark-Wilson, R., Breeze, P.S., Armitage, S.J., *et al.*, 2021. Multiple hominin dispersals into Southwest Asia over the past 400,000 years. *Nature* **597**, 376–380. doi:10.1038/s41586-021-03863-y
- Hallinan, E., Shaw, M., 2015. A new Middle Stone Age industry in the Tanwka Karoo, Northern Cape Province, South Africa. Antiquity Project Gallery 89, 344. http://antiquity.ac.uk/projgall/hallinan344.
- Hardaker, T., 2020. A geological explanation for occupation patterns of ESA and early MSA humans in Southwestern Namibia? An interdisciplinary study. Proceedings of the Geologists' Association 131, 8–18.
- Hilbert, Y.H., Crassard, R., Rose, J.I., Geiling, J.M., Usik, V.I., 2016. Technological homogeneity within the Arabian Nubian Complex: comparing chert and quartzite assemblages from central and southern Arabia. *Journal of Lithic Studies* 3, 411–437.
- Hiscock, P., 1985. The need for a taphonomic perspective in stone artefact analysis. Queensland Archaeological Research 2, 82–97.
- Holdaway, S.J., Davies, B., 2019. Surface stone artifact scatters, settlement patterns, and new methods for stone artifact analysis. *Journal of Paleolithic Archaeology* 3, 612–632.
- **Honea, K.H.,** 1964. The patination of stone artifacts. *Plains Anthropologist* **9**, 14–17.
- Hovers, E., 2009. The Lithic Assemblages of Qafzeh Cave. Oxford University Press, Oxford, UK.
- Hovers, E., 2003. Treading carefully: site formation processes and Pliocene lithic technology. In: Martínez-Moreno, J., Torcal, R.M., de la Torre Sainz, I. (Eds.), Oldowan: Rather More than Smashing Stones: First Hominid Technology Workshop. Universitat Autònoma de Barcelona, Centre d'Estudis del Patrimoni Arqueològic de la Prehistòria, Barcelona, pp. 145–164
- Hovers, E., Ekshtain, R., Greenbaum, N., Malinsky-Buller, A., Nir, N., Yeshurun, R., 2014. Islands in a stream? Reconstructing site formation processes in the late Middle Paleolithic site of 'Ein Qashish, northern Israel. Quaternary International 331, 216–233.
- **Howard, C.D.**, 1999. River patina on flint artifacts: features and genesis. *Plains Anthropologist* **44**, 293–295.
- Howard, C.D., 2002. The gloss patination of flint artifacts. Plains Anthropologist 47, 283–287.
- Hurst, V.J., Kelly, A.R., 1961. Patination of cultural flints: flint artifacts can be dated by cortical changes in mineralogy and texture. Science 134, 251–256.

- Jennings, T.A., 2011. Experimental production of bending and radial flake fractures and implications for lithic technologies. *Journal of Archaeological Science* 38, 3644–3651.
- Kidron, G.J., Zohar, M., Starinsky, A., 2014. Spatial distribution of dust deposition within a small drainage basin: implications for loess deposits in the Negev desert. Sedimentology 61, 1908–1922.
- Kijowska-Strugała, M., Wiejaczka, Ł., Lekach, J., Bucała-Hrabia, A., 2017. Diversification of the hydromorphological state and the habitat quality of streams in the Negev desert (Israel). *Environmental Earth Sciences* **76**, 99. https://doi.org/10.1007/s12665-016-6347-1.
- Knight, J., Zerboni, A., 2018. Formation of desert pavements and the interpretation of lithic-strewn landscapes of the central Sahara. *Journal of Arid Environments* 153, 39–51.
- Lucke, B., Bäumler, R., 2021. Holocene dust dynamics archives in archaeological ruins in arid and semi-arid environments in the Southern Levant. Arabian Journal of Geosciences 14, 2663. https://doi.org/10.1007/s12517-021-08884-5
- Lucke, B., Roskin, J., Vanselow, K.A., Bruins, H.J., Abu-Jaber, N., Deckers, K., Lindauer, S., et al., 2019. Character, rates, and environmental significance of Holocene dust accumulation in archaeological Hilltop Ruins in the southern Levant. *Geosciences* 9, 190. https://doi.org/10.3390/geosciences9040190.
- Malinsky-Buller, A., 2016. The muddle in the Middle Pleistocene: the Lower–Middle Paleolithic transition from the Levantine perspective. *Journal of World Prehistory* 29, 1–78.
- Malinsky-Buller, A., Hovers, E., Marder, O., 2011. Making time: 'Living floors', 'palimpsests' and site formation processes—a perspective from the open-air Lower Paleolithic site of Revadim Quarry, Israel. *Journal of Anthropological Archaeology* 30, 89–101.
- Matmon, A., Elfassi, S., Hidy, A., Geller, Y., Porat, N., Team, A., 2016.

 Controls on aggradation and incision in the NE Negev, Israel, since the Middle Pleistocene. *Geomorphology* 261, 132–146.
- Matmon, A., Zilberman, E., 2017. Landscape evolution along the Dead Sea fault and its margins. In: Enzel, Y., Bar-Yosef, O. (Eds.), *Quaternary of the Levant (Environment, Climate Change and Humans)*. Cambridge University Press, Cambridge, UK, New York, pp. 17–30.
- McBrearty, S., Bishop, L., Plummer, T., Dewar, R., Conard, N., 1998. Tools underfoot: human trampling as an agent of lithic artifact edge modification. *American Antiquity* 63, 108–129.
- Munday, F.C., 1977. Nahal Aqev (D35): a stratified, open-air Mousterian occupation in the Avdat/Aqev area. In: Marks, A.E. (Ed.), *Prehistory and Paleoenvironments of the Central Negev, Israel, Vol. II, The Avdat/Aqev Area, Part 2 and the Har Harif.* Southern Methodist University Press, Dallas, Texas, pp. 35–60.
- Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiation Measurements* 32, 57–73.
- Olszewski, D.I., Dibble, H.L., McPherron, S.P., Schurmans, U.A., Chiotti, L., Smith, J.R., 2010. Nubian complex strategies in the Egyptian high desert. *Journal of Human Evolution* 59, 188–201.
- Oron, M., Groman-Yaroslavski, I., Lavi, R., Porat, N., Roskin, J., 2019. Mitzpe Ramon: a flint quarry and blade production workshop dated to the early Late Neolithic in the Central Negev Highlands, Israel. *Mitekufat Haeven: Journal of the Israel Prehistoric Society* **49**, 100–118.
- Porat, N., Amit, R., Enzel, Y., Zilberman, E., Avni, Y., Ginat, H., Gluck, D., 2010. Abandonment ages of alluvial landforms in the hyperarid Negev determined by luminescence dating. *Journal of Arid Environments* 74, 861–869.
- Porat, N., Duller, G.A., Amit, R., Zilberman, E., Enzel, Y., 2009. Recent faulting in the southern Arava, Dead Sea transform: evidence from single grain luminescence dating. *Quaternary International* 199, 34–44.
- Pye, K., Tsoar, H., 1987. The mechanics and geological implications of dust transport and deposition in deserts with particular reference to loess formation and dune sand diagenesis in the northern Negev, Israel. *Geological Society, London, Special Publications* 35, 139–156.
- Robins, L., Greenbaum, N., Yu, L., Bookman, R., Roskin, J., 2021. High-resolution portable-OSL analysis of vegetated linear dune construction in the margins of the northwestern Negev dunefield (Israel) during

the late Quaternary. Aeolian Research 50, 100680. https://doi.org/10.1016/j. aeolia.2021.100680.

- Robins, L., Roskin, J., Yu, L., Bookman, R., Greenbaum, N., 2022. Aeolian-fluvial processes control landscape evolution along dunefield margins of the northwestern Negev (Israel) since the late Quaternary. *Quaternary Science Reviews* 285, 107520. https://doi.org/10.1016/j.quascirev.2022.107520.
- Roded, R., 1996. Geological Map of Israel (1:50,000): Dimona Sheet (19-1). The Geological Survey of Israel, Jerusalem, Israel.
- Roskin, J., Katra, I., Agha, N., Goring-Morris, A.N., Porat, N., Barzilai, O., 2014. Rapid anthropogenic response to short-term aeolian-fluvial palaeoenvironmental changes during the Late Pleistocene–Holocene transition in the northern Negev desert, Israel. *Quaternary Science Reviews* 99, 176–192.
- Roskin, J., Katra, I., Porat, N., Zilberman, E., 2013. Evolution of Middle to Late Pleistocene sandy calcareous paleosols underlying the northwestern Negev desert dunefield (Israel). *Palaeogeography, Palaeoclimatology*, *Palaeoecology* 387, 134–152.
- Roskin, J., Tsoar, H., 2017. Late Quaternary chronologies of the Northern Sinai–Northwestern Negev dunefield and their palaeoclimatic and palaeoenvironmental implications. In: Enzel, Y., Bar-Yosef, O. (Eds.), Quaternary of the Levant (Environment, Climate Change and Humans). Cambridge University Press, Cambridge, UK, New York, pp. 505–522.
- Rottländer, R., 1975. The formation of patina on flint. Archaeometry 17, 106–110.
- Schick, K.D., 1984. Processes of Palaeolithic Site Formation: an Experimental Study. Ph.D. dissertation, University of California, Berkeley, California.
- **Stern, N.**, 1994. The implications of time-averaging for reconstructing the land-use patterns of early tool-using hominids. *Journal of Human Evolution* **27**, 89–105.
- Thompson, J.C., Mackay, A., De Moor, V., Gomani-Chindebvu, E., 2014.
 Catchment survey in the Karonga district: a landscape-scale analysis of provisioning and core reduction strategies during the Middle Stone Age of northern Malawi. African Archaeological Review 31, 447–478.
- Ungar, E.D., Stavi, I., Lavee, H., Sarah, P., 2010. Effects of livestock traffic on rock fragment movement on hillsides in a semiarid patchy rangeland. *Land Degradation and Development* 21, 92–99.
- Vaquero, M., Chacón, M.G., García-Antón, M.D., Soler, B.G. de, Martínez, K., Cuartero, F., 2012. Time and space in the formation of lithic assemblages: the example of Abric Romaní Level J. Quaternary International 247, 162–181.
- Vardi, J., Yegorov, D., Degen-Eisenberg, D., Boaretto, E., Langgut, D., Avni, Y., Caracuta, V., 2023. The utilization and extinction of Juniper trees from the Negev desert (Israel)—data from a late 6th–5th millennia site of Har Harif. *Journal of Arid Environments* 210, 104906. https://doi. org/10.1016/j.jaridenv.2022.104906.

- Villa, P., 1982. Conjoinable pieces and site formation processes. American Antiauity 47, 276–290.
- Vogel, J.C., Geyh, M.A., 2008. Radiometric dating of hillslope calcrete in the Negev desert, Israel. South African Journal of Science 104, 493–495.
- Wdowinski, S., Sneh, A., Avni, Y., 2012. Geological Map of Israel (1:50,000): Tel Malhata Sheet (15-III). The Geological Survey of Israel, Jerusalem, Israel.
- Wieler, N., Avni, Y., Rosensaft, M., 2016. The significance of the geological strata on desert runoff agriculture: indications for stable desert environment over the last 1600 years in southern Israel. *Journal of Arid Environments* 135, 147–163.
- Wieler, N., Gini, T.E., Gillor, O., Angel, R., 2021. Estimating the growth rate in desert biological rock crusts by integrating archaeological and geological records. *Biogeosciences Discussions* 2021. https://doi.org/10.5194/bg-2020-467.
- Wright, D.K., Thompson, J.C., Schilt, F., Cohen, A.S., Choi, J.-H., Mercader, J., Nightingale, S., et al., 2017. Approaches to Middle Stone Age landscape archaeology in tropical Africa. *Journal of Archaeological Science* 77, 64–77.
- Yaalon, D.H., Dan, J., 1974. Accumulation and distribution of loess-derived deposits in the semi-desert and desert fringe areas of Israel. Zeitschrift für Geomorphologie, Supplementband 20, 91–105.
- Yair, A., 1983. Hillslope hydrology water harvesting and areal distribution of some ancient agricultural systems in the northern Negev desert. *Journal of Arid Environments* 6, 283–301.
- Yair, A., Enzel, Y., 1987. Relationship between annual rainfall and sediment yield in arid and semi-arid areas: the case of the northern Negev. In: Ahnert, F. (Ed.), Geomorphological Models: Theoretical and Empirical Aspects. Catena Supplement 10, 121–135.
- Yegorov, D., 2017. Nahal Yitnan. *Hadashot Arkheologiyot* 129. https://www.hadashot-esi.org.il/report_detail_eng.aspx?id=25178&mag_id=125.
- Yegorov, D., Marder, O., Khalaily, H., Milevski, I., Rosen, S.A., 2020. Heat treated or not heat treated: archaeological and experimental interpretation of flint assemblage from the Middle Pre-Pottery Neolithic B site of Yiftahel. *Journal of Archaeological Science: Reports* 29, 102090. https://doi.org/10.1016/j.jasrep.2019.102090.
- Zaidner, Y., Weinstein-Evron, M., 2020. The emergence of the Levallois technology in the Levant: a view from the early Middle Paleolithic site of Misliya Cave, Israel. *Journal of Human Evolution* 144, 102785. https://doi.org/10.1016/j.jhevol.2020.102785.
- Zilberman, E., 1991. Landscape Evolution in the Central, Northern and Northwestern Negev during the Neogene and the Quaternary. *Geological Society of Israel Report* GSI/45/90, 1–108. [in Hebrew]
- Zilberman, E., Avni, Y., 2005. Geological Map of Israel (1:50,000), Har Hamran Sheet (21-I). The Geological Survey of Israel, Jerusalem, Israel.