

ARTICLE

Maximal chordal subgraphs

Lior Gishboliner and Benny Sudakov

Department of Mathematics, ETH, Zürich, Switzerland

Corresponding author: Lior Gishboliner; Email: lior.gishboliner@math.ethz.ch

(Received 24 May 2022; revised 25 January 2023; accepted 22 March 2023; first published online 2 May 2023)

Abstract

A chordal graph is a graph with no induced cycles of length at least 4. Let $f(n, m)$ be the maximal integer such that every graph with n vertices and m edges has a chordal subgraph with at least $f(n, m)$ edges. In 1985 Erdős and Laskar posed the problem of estimating $f(n, m)$. In the late 1980s, Erdős, Gyárfás, Ordman and Zalcstein determined the value of $f(n, n^2/4 + 1)$ and made a conjecture on the value of $f(n, n^2/3 + 1)$. In this paper we prove this conjecture and answer the question of Erdős and Laskar, determining $f(n, m)$ asymptotically for all m and exactly for $m \leq n^2/3 + 1$.

Keywords: chordal subgraph

2020 MSC Codes: Primary: 05C35

1. Introduction

One of the central questions in extremal combinatorics can be formulated as follows. Given a graph G and a property \mathcal{P} , what is the maximal subgraph of G one can find which satisfies this property. The study of this problem goes back to the work of Turán in 1941, whose theorem states that the largest subgraph of the n -vertex complete graph with no clique of size $k + 1$ is the complete k -partite graph with sides as equal as possible. This graph is called the Turán graph. We denote it by $T_k(n)$ and its size by $t_k(n)$. Turán's theorem is the starting point of extremal graph theory and has inspired extensive research. One such research direction studies which other (more elaborate) structures must appear in a graph with more than $t_k(n)$ edges. For example, a series of works determined how many $(k + 1)$ -cliques must exist in a graph with $t_k(n) + a$ edges (for a suitable range of a) [7, 11, 16]. Other examples are results on finding many $(k + 1)$ -cliques which share one or more vertices [5, 7, 8, 15], and results on finding $(k + 1)$ -cliques with large degree sum [1, 3, 4, 12].

In this paper we study the Turán type problem for chordal graphs. A graph is called chordal if it contains no induced cycle of length at least 4. Chordal graphs are one of the most studied classes in graph theory and have numerous applications, for example in semidefinite optimisation (see the survey [17]) and evolutionary trees (see [6]). In 1985, Erdős and Laskar [10] asked to determine the maximum integer $f(n, m)$ such that every graph with n vertices and m edges contains a chordal subgraph with at least $f(n, m)$ edges. To put this question under the umbrella of classical extremal graph theory, one needs to consider equivalent definitions of chordal graphs. It is well known that a graph is chordal if and only if it can be constructed from a single-vertex graph by repeatedly adding a vertex and connecting it to a clique of the current graph¹ (this is

¹A related fact is that a graph is chordal if and only if it has a tree-decomposition in which the bags are cliques. So chordal graphs can be thought of as 'trees of cliques'.



called a perfect elimination ordering), see [13, Chapter 4]. So if G is a triangle-free graph, then every chordal subgraph of G must be a forest. More generally, if G has no cliques of size $k + 1$, then every chordal subgraph of G has at most $(k - 1)(n - k + 1) + \binom{k-1}{2} = (k - 1)n - \binom{k}{2}$ edges. In particular, this bound applies to k -partite graphs. Another way of proving this bound for k -partite graphs is to observe that if G is k -partite with parts V_1, \dots, V_k and H is a chordal subgraph of G , then $e_H(V_i, V_j) \leq |V_i| + |V_j| - 1$ for every $i < j$ (because a chordal subgraph of a bipartite graph must be a forest). Hence, $e(H) \leq \sum_{i < j} (|V_i| + |V_j| - 1) = (k - 1)n - \binom{k}{2}$.

The above discussion shows that if $m \leq t_k(n)$ then $f(n, m) \leq (k - 1)n - \binom{k}{2}$. It is natural to guess that the value of $f(n, m)$ “jumps” as m increases from $t_k(n)$ to $m = t_k(n) + 1$, because at this point the graph must contain $(k + 1)$ -cliques. Erdős and Laskar [10] proved that this is indeed the case for $k = 2$, showing that $f(n, t_2(n) + 1) \geq (1 + \epsilon)n$. In the late 80’s, Erdős, Gyárfás, Ordman and Zalcstein [9] determined the value of $f(n, t_2(n) + 1)$ exactly for even n , showing that $f(n, \frac{n^2}{4} + 1) = \frac{3n}{2} - 1$. This bound is achieved by the graph $T_2(n) + e$, obtained by adding an edge to the Turán graph $T_2(n)$. It is natural to conjecture that for every k and n , the value of $f(n, t_k(n) + 1)$ is determined by $T_k(n) + e$, which is the graph obtained by adding an edge to a largest class of $T_k(n)$. It is not hard to check that the largest chordal subgraph of $T_k(n) + e$ has $kn - \lceil \frac{n}{k} \rceil + 2 - \binom{k+1}{2}$ edges. So we get the following conjecture.

Conjecture 1.1. $f(n, t_k(n) + 1) = kn - \lceil \frac{n}{k} \rceil + 2 - \binom{k+1}{2}$.

The authors of [9] only studied Conjecture 1.1 in the cases $k = 2, 3$, although they very likely had the full conjecture in mind. For $k = 3$, they proved that $f(n, t_3(n) + 1) \geq 7n/3 - 6$ and asked to determine $f(n, t_3(n) + 1)$. This question was later mentioned again in the problem survey of Gyárfás [14]. Answering this question, we resolve Conjecture 1.1 for the case $k = 3$.

Theorem 1.2. $f(n, t_3(n) + 1) = 3n - \lceil \frac{n}{3} \rceil - 4$.

Our next result proves Conjecture 1.1 asymptotically for every k . In fact, we go a step further and determine $f(n, m)$ asymptotically for every value of m , answering the question of Erdős and Laskar.

Theorem 1.3. Let $k, n \geq 1$ and $t_k(n) + 1 \leq m \leq t_{k+1}(n)$. Set $a = m - t_k(n)$. Then

$$f(n, m) = (k - 1/k)n + \sqrt{2(k + 1)a/k} - \binom{k + 1}{2} - O(\sqrt{n}).$$

The construction giving the upper bound in Theorem 1.3 is to take an (unbalanced) complete k -partite graph with $k - 1$ smaller classes of the same size and one bigger class, and to add a balanced complete bipartite graph inside the bigger class. One then needs to optimise the sizes of the classes and the size of the complete bipartite graph so as to minimise the size of chordal subgraphs. It is best to take the $k - 1$ smaller classes of size $\frac{n-r}{k}$, the bigger class of size $\frac{n+(k-1)r}{k}$ and the complete bipartite of size $r \times r$, where $r := \sqrt{\frac{2ka}{k+1}}$. See Figure 1, and see Section 2 for the details.

For $k = 1, 2$, we can go a step further and determine $f(n, m)$ exactly. This is done in the following two theorems.

Theorem 1.4. Let $n \geq 1$ and $m \leq t_2(n)$. Then $f(n, m) = \min\{r : t_2(r) \geq m\} - 1$.

For $n \geq 1$ and $m \geq t_2(n) + 1$, let $g_2(n, m)$ be the minimum of $2n - t + r$, taken over all pairs $t, r \geq 0$ satisfying $t(n - t) + t_2(r) \geq m$.

Theorem 1.5. Let $n \geq 1$ and $t_2(n) + 1 \leq m \leq t_3(n)$. Then $f(n, m) = g_2(n, m) - 3$.

The extremal construction for Theorem 1.5 is given by taking a $t \times (n - t)$ complete bipartite graph and placing a complete bipartite graph with r vertices inside the side of size t .




Figure 1. The construction showing optimality of Theorem 1.3.

1.1. Proof ideas

Recall that a graph is chordal if and only if it can be obtained from the one-vertex graph by repeatedly adding *simplicial vertices*, that is, vertices whose neighbourhood is a clique. In particular, adding simplicial vertices to a chordal graph keeps it chordal. We will often use this fact (implicitly) to claim that certain graphs are chordal.

Let us first recall the argument used by Erdős, Gyárfás, Ordman and Zalcstein [9] to prove Conjecture 1.1 for $k = 2$ (and n even). Let G be a graph with n vertices and $n^2/4 + 1$ edges, and let x, y, z be a triangle in G . We need to show that G has a chordal subgraph H with at least $3n/2 - 1$ edges. If $d(x) + d(y) + d(z) \geq 3n/2 + 2$, then take H to be the subgraph consisting of all edges touching x, y, z . Suppose now that $d(x) + d(y) + d(z) \leq 3n/2 + 1$. Then by averaging, we can assume without loss of generality that $d(x) + d(y) \leq n$. Deleting x, y , we get a graph with at least $n^2/4 + 1 - (n - 1) \geq (n - 2)^2/4 + 1$ edges. By induction, this graph contains a chordal subgraph H' with at least $3(n - 2)/2 - 1$ edges. Adding the edges xy, xz, yz gives the required chordal subgraph H .

Our proof of Theorem 1.5 is also based on this inductive argument, but with two key differences. First, we need a relation between $g_2(n, m)$ and $g_2(n', m')$ (for $n' = n - 2$, say), so that the induction can be carried through when deleting vertices. And second, it turns out that the induction scheme of deleting two vertices does not work to give the correct bound on $f(n, m)$ for all m in the range of Theorem 1.5. Instead, we sometimes need to delete just one vertex and then add two edges when adding the vertex back. To this end, we need to know that the deleted vertex has two neighbours which form an edge in the chordal subgraph H' that we find using induction. To guarantee this, we strengthen the induction hypothesis to say that not only does G contain a chordal subgraph with the correct number of edges, but that any given triangle in G can be included in such a chordal subgraph.

The idea of strengthening the induction hypothesis is also used in the proof of Theorem 1.2. Here we show that every K_4 can be included in a chordal subgraph with the correct number of edges. This proof has a more involved case analysis. It would be interesting to find a shorter proof.

The proof of Theorem 1.3 is based on induction as well. Here, instead of deleting only a few vertices, we delete a large number of vertices. To give the general idea, we sketch first the proof in the case $m = t_3(n) + 1$. So let G be a graph with n vertices $t_3(n) + 1$ edges. We need to show that G has a chordal subgraph H with at least $\frac{8n}{3} - 6 - C\sqrt{n}$ edges. Let us assume first that $e(G) \geq t_3(n) + 2n$. By a theorem of Faudree [12] (see also [1, 3]), there is a triangle $x, y, z \in V(G)$ with $d(x) + d(y) + d(z) \geq 6e(G)/n \geq 2n + 12$. In particular, x, y, z have at least 12

common neighbours. Let w_1, \dots, w_7 be seven of them. If $d(x) + d(y) + d(z) + d(w_i) \geq \frac{8n}{3} - C\sqrt{n}$ for some i , then take H to be the subgraph consisting of all edges touching x, y, z, w_i . This H is chordal and $e(H) = d(x) + d(y) + d(z) + d(w_i) - 6$, so we are done. Suppose then that $d(x) + d(y) + d(z) + d(w_i) \leq \frac{8n}{3} - C\sqrt{n}$ for every i . In particular, $d(w_i) \leq \frac{2n}{3} - C\sqrt{n}$. Assume that $d(x) \geq d(y) \geq d(z)$, so that $d(x) \geq \frac{2n}{3}$ and hence $d(y) + d(z) + d(w_i) \leq 2n - C\sqrt{n}$ for each i . Delete y, z, w_1, \dots, w_7 to get a graph G' on $n - 9$ vertices. It is easy to see that $e(G')$ is well above $t_3(n - 9) + 1$. So by the induction hypothesis, there exists a chordal subgraph H' of G' with $e(H') \geq \frac{8(n-9)}{3} - 6 - C\sqrt{n}$ edges. Now add back the vertices y, z, w_1, \dots, w_7 , and add to H' the edges of the triangle x, y, z and the edges between x, y, z and w_1, \dots, w_7 . This is a total of 24 edges. So $e(H) = e(H') + 24 \geq \frac{8n}{3} - 6 - C\sqrt{n}$, as required. It is also easy to see that H is chordal (if we add the new vertices in the order y, z, w_1, \dots, w_7 , then we always add a simplicial vertex). The number 7 was chosen here so that the number of edges added would be large enough for the induction to carry through. But the key point is that such a number must exist. Indeed, each w_i contributes 3 edges to H . On the other hand, the term $\frac{8n}{3}$ suggests that it is enough to add $\frac{8}{3}$ edges per vertex on average. So by adding 3 edges per vertex, we are gaining over the required bound.

It now remains to handle the case that $e(G) \leq t_3(n) + 2n$. Here we proceed as follows. If the minimum degree of the graph is at least $\frac{2n}{3} - \sqrt{n}$, then take a 4-clique x, y, z, w and take H to be the subgraph consisting of edges touching x, y, z, w . Else, delete a vertex of minimum degree and continue with the remaining graph. After $O(\sqrt{n})$ steps, we get a graph with $n' = n - O(\sqrt{n})$ vertices and at least $t_3(n') + 2n'$ edges, so we can apply the first case.

To prove the general case of Theorem 1.3 we find a $(k - 1)$ -clique x_1, \dots, x_{k-1} and a forest F inside $N(x_1, \dots, x_{k-1})$ such that F has few components. We delete $V(F)$ and x_2, \dots, x_{k-1} and apply induction to find a chordal subgraph H' . We then add to H' the edges of the clique x_1, \dots, x_{k-1} , the edges of F , and the edges between $V(F)$ and x_1, \dots, x_{k-1} . Note that when adding back the vertices of F one by one, most vertices contribute k edges: one edge in F and $k - 1$ edges to x_1, \dots, x_{k-1} (this fails once for each connected component of F , and this is why we want the number of components to be small). On the other hand, the main term in Theorem 1.3 is $(k - 1/k)n$, which suggests that each vertex adds $k - 1/k$ edges on average. So again we are gaining over the required bound (at least if we ignore the second term $\sqrt{2(k + 1)a/k}$ for the moment). A somewhat lengthy calculation shows that this argument indeed works for any value of a .

The rest of this short paper is organised as follows. Theorem 1.3 is proved in Section 2, Theorems 1.4–1.5 in Section 3 and Theorem 1.2 in Section 4.

2. Proof of Theorem 1.3

In this section we prove Theorem 1.3. We begin with the upper bound. Here we use the following construction. For simplicity, assume that n is divisible by $k, k + 1$. For general n the construction is essentially the same (and, since we are only interested in an approximate result, we are allowed a small error due to divisibility issues). Fix $k \geq 1$ and $m \leq t_{k+1}(n) = \frac{kn^2}{2(k+1)}$, so that $a := m - t_k(n) \leq \frac{kn^2}{2(k+1)} - \frac{(k-1)n^2}{2k} = \frac{n^2}{2k(k+1)}$. Set $r := \sqrt{\frac{2ka}{k+1}} \leq \frac{n}{k+1}$. Consider a complete k -partite graph with sides X, Y_1, \dots, Y_{k-1} such that $|X| = \frac{n+(k-1)r}{k}$ and $|Y_i| = \frac{n-r}{k}$ for every $1 \leq i \leq k - 1$. Place an $r \times r$ complete bipartite graph with sides A, B inside X . This is possible as $2r \leq \frac{n+(k-1)r}{k}$. The resulting graph G has

$$\begin{aligned}
 e(G) &= (k - 1) \cdot \frac{n + (k - 1)r}{k} \cdot \frac{n - r}{k} + \binom{k - 1}{2} \left(\frac{n - r}{k}\right)^2 + r^2 = \frac{(k - 1)n^2}{2k} + \frac{(k + 1)r^2}{2k} \\
 &= \frac{(k - 1)n^2}{2k} + a = t_k(n) + a = m.
 \end{aligned}$$

Let H be a chordal subgraph of G . We have $e_H(A, B) \leq |A| + |B| - 1 = 2r - 1$, $e_H(A, Y_i) \leq |A| + |Y_i| - 1$, $e_H(X \setminus A, Y_i) \leq |X| - |A| + |Y_i| - 1$ and $e_H(Y_i, Y_j) \leq |Y_i| + |Y_j| - 1$, because each of these bipartite graphs is induced in G , so its intersection with H is a forest. So

$$\begin{aligned} e(H) &\leq 2r - 1 + (k - 1)|X| - 2(k - 1) + 2 \sum_{i=1}^{k-1} |Y_i| + \sum_{1 \leq i < j \leq k-1} (|Y_i| + |Y_j| - 1) \\ &= kn - |X| + 2r - \binom{k + 1}{2} = (k - 1/k)n + \frac{(k + 1)r}{k} - \binom{k + 1}{2} \\ &= (k - 1/k)n + \sqrt{2(k + 1)a/k} - \binom{k + 1}{2}, \end{aligned}$$

giving the upper bound on $f(n, m)$ for Theorem 1.3. We now prove the lower bound, which we restate for convenience as follows.

Theorem 2.1. *For every $k \geq 1$ there is $C = C(k)$ such that the following holds. Let $n, a \geq 1$, and let G be a graph with n vertices and at least $t_k(n) + a$ edges. Then G has a chordal subgraph with at least $(k - 1/k)n + \sqrt{2(k + 1)a/k} - C\sqrt{n} - \binom{k + 1}{2}$ edges.*

For the proof of Theorem 2.1 we need two lemmas. The following lemma uses an argument originally used by Edwards [3, 4] and Faudree [12] (see also [1]) to find cliques with a large degree sum.

Lemma 2.2. *Let $k, n, a \geq 1$ and let G be a graph with n vertices and at least $\frac{(k-1)n^2}{2k} + a$ edges. Consider the following process: for $i = 1, 2, \dots$, take x_i to be a vertex of maximum degree among all vertices in $N(x_1, \dots, x_{i-1})$. Then this process continues for at least k steps, and $N(x_1, \dots, x_{k-1})$ contains at least a edges.*

Proof. We prove the lemma by induction on k . The base case $k = 1$ is trivial. Let $k \geq 2$. By the induction hypothesis, the process continues for at least $k - 1$ steps. It remains to show that $N(x_1, \dots, x_{k-1})$ contains at least a edges, because this would also imply that $N(x_1, \dots, x_{k-1}) \neq \emptyset$ and hence the process continues for at least k steps. For $1 \leq i \leq k - 1$, let S_i be the set of vertices which are adjacent to x_1, \dots, x_{i-1} but not adjacent to x_i . In particular, S_1 is just the set of vertices not adjacent to x_1 and $x_i \in S_i$ for all i . Then $V(G) = S_1 \cup \dots \cup S_{k-1} \cup N(x_1, \dots, x_{k-1})$. Put $S := S_1 \cup \dots \cup S_{k-1}$, $N := N(x_1, \dots, x_{k-1})$, $s_i := |S_i|$, $s = |S|$ and $d_i := d(x_i)$. Note that $s_i \leq n - d_i$. Also, all vertices in S_i have degree at most d_i . We have

$$e(N, S) + 2e(S) = \sum_{v \in S} d(v) \leq \sum_{i=1}^{k-1} s_i \cdot d_i \leq \sum_{i=1}^{k-1} s_i(n - s_i). \tag{1}$$

Since $e(G) = e(N) + e(S) + e(N, S)$, we have $2e(S) = 2e(G) - 2e(N) - 2e(N, S)$. Plugging this into (1) and rearranging, we get

$$e(N) \geq e(G) - \frac{1}{2}e(N, S) - \frac{1}{2} \cdot \sum_{i=1}^{k-1} s_i(n - s_i). \tag{2}$$

We have $e(N, S) \leq |N| \cdot |S| = (n - s)s$. Also, by Cauchy–Schwarz,

$$\sum_{i=1}^{k-1} s_i(n - s_i) = ns - \sum_{i=1}^{k-1} s_i^2 \leq ns - \frac{s^2}{k - 1}.$$

Plugging this into (2) gives

$$e(N) \geq e(G) - \frac{1}{2}(n-s)s - \frac{1}{2} \left(ns - \frac{s^2}{k-1} \right) = e(G) - ns + \frac{ks^2}{2(k-1)}.$$

The maximum of $ns - \frac{ks^2}{2(k-1)}$ is obtained at $s = \frac{(k-1)n}{k}$ and equals $\frac{(k-1)n^2}{2k}$. Hence, $e(N) \geq e(G) - \frac{(k-1)n^2}{2k} \geq a$, as required. \square

Lemma 2.3. *Let G be a graph with n vertices and a edges. Let $s \geq 1$ and suppose that $a \geq 2s^2$. Then G contains a forest F with s vertices and at least $s - 1 - \frac{sn}{a}$ edges.*

Proof. Let C_1, \dots, C_m be the connected components of G with $|C_1| \geq \dots \geq |C_m|$. Let $\ell \geq 1$ be the minimal integer satisfying $|C_1| + \dots + |C_\ell| \geq s$. If $\ell \leq 1 + \frac{sn}{a}$ then take F to be a forest contained in $C_1 \cup \dots \cup C_\ell$ having s vertices and ℓ connected components. Suppose now by contradiction that $\ell > 1 + \frac{sn}{a}$. Set $r = |C_1| + \dots + |C_{\ell-1}|$. Then $r < s$ and $|C_{\ell-1}| \leq \frac{r}{\ell-1}$. We have $e(G) \leq \binom{r}{2} + \sum_{i=\ell}^m \binom{|C_i|}{2}$. By convexity, the sum $\sum_{i=\ell}^m \binom{|C_i|}{2}$ is maximised when all except maybe one of the $|C_i|$'s are equal to their maximal value, which is $\frac{r}{\ell-1}$. So

$$\begin{aligned} e(G) &\leq \binom{r}{2} + \left\lceil \frac{n-r}{r/(\ell-1)} \right\rceil \cdot \binom{r/(\ell-1)}{2} \leq \binom{r}{2} + \left(1 + \frac{n-r}{r/(\ell-1)} \right) \cdot \binom{r/(\ell-1)}{2} \\ &\leq \binom{r}{2} + \binom{r/(\ell-1)}{2} + \frac{nr}{2(\ell-1)} \\ &< 2 \binom{s}{2} + \frac{sn}{2sn/a} \leq a, \end{aligned}$$

where the last inequality uses $a \geq 2s^2$. We got a contradiction to $e(G) = a$. \square

We are now ready to prove Theorem 2.1. An overview of the proof can be found in Section 1.1.

Proof of Theorem 2.1. The proof is by induction on n . Fix constants $k \ll c \ll c_1 \ll C$, to be chosen implicitly later. Suppose first that $a \leq (ck+1)^2n$. In this case we proceed as follows. If $\delta(G) \geq \lfloor \frac{(k-1)n}{k} \rfloor - c_1\sqrt{n}$, then take a $(k+1)$ -clique $x_1, \dots, x_{k+1} \in V(G)$ and take H to consist of all edges that touch x_1, \dots, x_{k+1} . Then H is chordal and

$$\begin{aligned} e(H) &= \sum_{i=1}^{k+1} d(x_i) - \binom{k+1}{2} \geq (k+1) \cdot \left(\frac{(k-1)n}{k} - 2c_1\sqrt{n} \right) - \binom{k+1}{2} \\ &= (k-1/k)n - 2(k+1)c_1\sqrt{n} - \binom{k+1}{2} \geq (k-1/k)n + \sqrt{2(k+1)a/k} - C\sqrt{n} - \binom{k+1}{2}, \end{aligned}$$

where the last inequality holds as $C \gg c_1, c$ and $a \leq (ck+1)^2n$. Suppose now that there is $v \in V(G)$ with $d(v) \leq \lfloor \frac{(k-1)n}{k} \rfloor - c_1\sqrt{n}$. Let $G' = G - v$. Then

$$e(G') \geq t_k(n) + a - \left\lfloor \frac{(k-1)n}{k} \right\rfloor + c_1\sqrt{n} = t_k(n-1) + a + c_1\sqrt{n}.$$

By the induction hypothesis with parameter $a' = a + c_1\sqrt{n}$, G' contains a chordal subgraph H' with

$$e(H') \geq (k-1/k)(n-1) + \sqrt{2(k+1)a'/k} - C\sqrt{n} - \binom{k+1}{2}.$$

As $(k - 1/k)(n - 1) \geq (k - 1/k)n - k$, it suffices to show that $\sqrt{2(k + 1)a'/k} \geq \sqrt{2(k + 1)a/k} + k$. Squaring and plugging in the value of a' , we get

$$2(k + 1)/k \cdot (a + c_1\sqrt{n}) = 2(k + 1)a'/k \geq 2(k + 1)a/k + 2k\sqrt{2(k + 1)a/k} + k^2.$$

Cancelling the term $2(k + 1)a/k$ from both sides and rearranging, we see that it is enough to have

$$c_1\sqrt{n} \geq \frac{k^2}{k + 1}\sqrt{2(k + 1)a/k} + \frac{k^3}{2(k + 1)},$$

which holds because $a \leq (ck + 1)^2n$ and $c_1 \gg c$.

For the rest of the proof we assume that $a \geq (ck + 1)^2n$. Note that $e(G) \geq t_k(n) + a \geq \frac{(k-1)n^2}{2k} + \frac{a}{2}$ because $t_k(n) \geq \frac{(k-1)n^2}{2k} - O_k(1)$ and $a \geq c \gg k$. Let x_1, \dots, x_{k-1} be as in Lemma 2.2 and put $N = N(x_1, \dots, x_{k-1})$. By Lemma 2.2 we have $e(N) \geq \frac{a}{2}$. Also, the choice of x_1, \dots, x_{k-1} in Lemma 2.2 implies that $d(y) \leq d(x_{k-1}) \leq \dots \leq d(x_1)$ for every $y \in N$. For convenience, we set

$$d_0 := \frac{(k - 1)n}{k} + \sqrt{\frac{2a}{k(k + 1)}} - c\sqrt{n}.$$

Note that $d_0 \geq \frac{(k-1)n}{k}$ by our assumption that $a \geq (ck + 1)^2n$.

Claim 2.4. *If the statement of the theorem does not hold, then $G[N]$ contains a forest F with $v(F) = \lfloor \sqrt{n} \rfloor$, $e(F) \geq v(F) - 1 - \frac{2n^{3/2}}{a}$, and*

$$\sum_{y \in V(F)} d(y) \leq v(F) \cdot d_0 + n/k. \tag{3}$$

Proof. We consider two cases. Suppose first that there is $x_k \in N$ such that $d(x_k) \geq d_0$. Then $d(x_i) \geq d_0$ for every $1 \leq i \leq k - 1$. Hence, $d(x_1) + \dots + d(x_k) \geq k \cdot d_0$. This means that x_1, \dots, x_k have at least

$$kd_0 - (k - 1)n = \sqrt{\frac{2ka}{k + 1}} - ck\sqrt{n} \geq \sqrt{a} - ck\sqrt{n} \geq \sqrt{n}$$

common neighbours, where the last inequality holds by the assumption $a \geq (ck + 1)^2n$. Take F to be the star whose centre is x_k and whose leaves are $\lfloor \sqrt{n} \rfloor - 1$ common neighbours of x_1, \dots, x_k . Let $y \in N(x_1, \dots, x_k)$. If $d(y) \geq d_0$ then

$$d(x_1) + \dots + d(x_k) + d(y) \geq (k + 1)d_0 = (k - 1/k)n + \sqrt{2(k + 1)a/k} - (k + 1)c\sqrt{n},$$

and then the subgraph consisting of all edges touching $\{x_1, \dots, x_k, y\}$ is a chordal graph with at least $(k - 1/k)n + \sqrt{2(k + 1)a/k} - (k + 1)c\sqrt{n} - \binom{k+1}{2}$ edges, so the assertion of the theorem holds. Hence, we may assume that $d(y) \leq d_0$ for every $y \in N(x_1, \dots, x_k)$. This means that

$$\sum_{v \in V(F)} d(v) \leq d(x_k) + (v(F) - 1) \cdot d_0 \leq n/k + v(F) \cdot d_0,$$

as required by the claim. Also, F has the right number of edges, as $e(F) = v(F) - 1$.

The second case is that $d(y) \leq d_0$ for every $y \in N$. Since $e(N) \geq \frac{a}{2} \geq 2n$, we can apply Lemma 2.3 to $G[N]$ with parameters $\frac{a}{2}$ and $s = \lfloor \sqrt{n} \rfloor$ to obtain a forest F with $\lfloor \sqrt{n} \rfloor$ vertices and at least $v(F) - 1 - \frac{2n^{3/2}}{a}$ edges. All vertices in F have degree at most d_0 , so (3) holds. \square

We continue with the proof of the theorem. Let F be the forest given by Claim 2.4. Let G' be the graph obtained from G by deleting the $t := k - 2 + v(F)$ vertices $T := \{x_2, \dots, x_{k-1}\} \cup V(F)$. By (3), we have

$$\sum_{v \in T} d(v) \leq d(x_2) + \dots + d(x_{k-1}) + v(F) \cdot d_0 + n/k \leq (v(F) + k - 2) \cdot d_0 + n = t \cdot d_0 + n,$$

where the second inequality uses that $d_0 \geq \frac{(k-1)n}{k}$. As $e(G) \geq t_k(n) + a \geq \frac{(k-1)n^2}{2k} - O_k(1) + a$, we have that

$$\begin{aligned} e(G') &\geq e(G) - t \cdot d_0 - n \geq \frac{(k-1)n^2}{2k} - O_k(1) + a - t \cdot d_0 - n \\ &= \frac{(k-1)(n-t)^2}{2k} + \frac{(k-1)nt}{k} - \frac{(k-1)t^2}{2k} - O_k(1) + a - t \cdot d_0 - n \\ &= \frac{(k-1)(n-t)^2}{2k} - O_k(1) + a - \frac{(k-1)t^2}{2k} - t \cdot \sqrt{\frac{2a}{k(k+1)}} + ct\sqrt{n} - n \\ &\geq t_k(n-t) + a - \frac{(k-1)t^2}{2k} - t \cdot \sqrt{\frac{2a}{k(k+1)}} + \frac{c}{2}t\sqrt{n}, \end{aligned}$$

where the last inequality uses that $t \geq \lfloor \sqrt{n} \rfloor$ and $c \gg k$, so that $\frac{c}{2}t\sqrt{n} \geq O_k(1) + n$. Set

$$a' := a - \frac{(k-1)t^2}{2k} - t \cdot \sqrt{\frac{2a}{k(k+1)}} + \frac{c}{2}t\sqrt{n},$$

so that $e(G') \geq t_k(n-t) + a'$. We have $a' \geq 1$ because $t \leq \sqrt{n} + k - 2$, $a \geq c^2n$ (say) and $c \gg k$. By the induction hypothesis, G' contains a chordal subgraph H' of size at least

$$e(H') \geq (k-1/k) \cdot (n-t) + \sqrt{2(k+1)a'/k} - C\sqrt{n-t} - \binom{k+1}{2}.$$

Let H be the subgraph of G obtained by adding to H' the edges of the clique x_1, \dots, x_{k-1} , the edges between x_1, \dots, x_{k-1} and $V(F)$, and the edges of F . Then H is chordal. To complete the proof, it suffices to verify that

$$e(H) \geq (k-1/k)n + \sqrt{2(k+1)a/k} - C\sqrt{n} - \binom{k+1}{2}.$$

By the definition of H , we have

$$e(H) = e(H') + \binom{k-1}{2} + (k-1) \cdot v(F) + e(F).$$

Note that

$$(k-1) \cdot v(F) + e(F) \geq k \cdot v(F) - 1 - \frac{2n^{3/2}}{a} = k \cdot (t-k+2) - 1 - \frac{2n^{3/2}}{a},$$

and therefore,

$$\binom{k-1}{2} + (k-1) \cdot v(F) + e(F) \geq tk - \frac{(k+1)(k-2)}{2} - 1 - \frac{2n^{3/2}}{a}.$$

For convenience, set $h := \frac{(k+1)(k-2)}{2} + 1 + \frac{2n^{3/2}}{a}$. Then,

$$e(H) \geq (k-1/k) \cdot (n-t) + \sqrt{2(k+1)a/k} - C\sqrt{n-t} - \binom{k+1}{2} + tk - h. \tag{4}$$

So it remains to verify that the right-hand side of (4) is at least as large as

$$(k - 1/k)n + \sqrt{2(k + 1)a/k} - C\sqrt{n} - \binom{k + 1}{2}.$$

Cancel the terms $(k - 1/k)n, \binom{k+1}{2}$ which appear in both expressions. Also, we may drop the terms $C\sqrt{n - t}, C\sqrt{n}$. After rearranging, we get the inequality

$$\sqrt{2(k + 1)a'/k} \geq \sqrt{2(k + 1)a/k} - \frac{t}{k} + h.$$

By squaring and plugging in the value of a' , we get:

$$\begin{aligned} & \frac{2(k + 1)}{k} \cdot \left(a - \frac{(k - 1)t^2}{2k} - t \cdot \sqrt{\frac{2a}{k(k + 1)}} + \frac{c}{2}t\sqrt{n} \right) \geq \\ & \frac{2(k + 1)a}{k} + \frac{t^2}{k^2} + h^2 - \frac{2t}{k} \cdot \sqrt{\frac{2(k + 1)a}{k}} + 2h \cdot \sqrt{\frac{2(k + 1)a}{k}} - \frac{2th}{k}. \end{aligned} \tag{5}$$

Both sides of the inequality (5) have the terms $\frac{2(k+1)}{k}a$ and $-\frac{2t}{k} \cdot \sqrt{\frac{2(k+1)a}{k}}$. We can also drop the negative term $\frac{2th}{k}$ on the right-hand side. After rearranging, we get the inequality

$$\frac{2(k + 1)}{k} \cdot \frac{c}{2}t\sqrt{n} \geq t^2 + h^2 + 2h \cdot \sqrt{\frac{2(k + 1)a}{k}}. \tag{6}$$

We have $t \leq \sqrt{n} + k$ and $h \leq O_k(1) + \frac{2n^{3/2}}{a} \leq O_k(1) + \sqrt{n}$, so $t^2, h^2 \leq O_k(n)$. Also,

$$h\sqrt{a} \leq \left(O_k(1) + \frac{2n^{3/2}}{a} \right) \cdot \sqrt{a} \leq O_k(n) + \frac{2n^{3/2}}{\sqrt{a}} = O_k(n),$$

as $n \leq a \leq n^2$. So the right-hand side of (6) is $O_k(n)$. On the other hand, the left-hand side is larger than $\frac{cn}{2}$ because $t \geq \lfloor \sqrt{n} \rfloor \geq \sqrt{n}/2$. So (6) holds because $c \gg k$, as required. \square

3. Proof of Theorems 1.4 and 1.5

Proof of Theorem 1.4. For the upper bound, let $r \geq 1$ be the minimal integer satisfying $t_2(r) \geq m$ and take G to be $T_2(r)$ with $n - r$ isolated vertices. Then $e(G) \geq m$, but every chordal subgraph of G has at most $r - 1$ edges. For the lower bound, we prove by induction on the number of vertices that every graph G with m edges has a chordal subgraph H with at least $g_1(m) - 1$ edges, where $g_1(m) := \min_{r: t_2(r) \geq m} r$. For $m = 0$, the assertion is trivial. Suppose $m \geq 1$ and let $xy \in E(G)$. Fix r such that $g_1(m) = r$. If $d(x) + d(y) \geq r$ then take H to be the subgraph consisting of all edges touching x, y . This graph is chordal and has $d(x) + d(y) - 1$ edges. Suppose now that $d(x) + d(y) \leq r - 1$; without loss of generality, $d(x) \leq \lfloor \frac{r-1}{2} \rfloor$. Let $G' = G - x$. Then $e(G') = m - d(x) \geq m - \lfloor \frac{r-1}{2} \rfloor$. We claim that $e(G') > t_2(r - 2)$. Indeed, if $e(G') \leq t_2(r - 2)$ then $m \leq t_2(r - 2) + \lfloor \frac{r-1}{2} \rfloor = t_2(r - 1)$, in contradiction to the choice of r . So $g_1(e(G')) \geq r - 1$. By the induction hypothesis, G' contains a chordal subgraph H' with at least $r - 2$ edges. Now, $H' + \{xy\}$ is a chordal subgraph of G with at least $r - 1$ edges, as required. \square

The rest of this section is dedicated to proving Theorem 1.5. Recall that for $n \geq 1$ and $m \geq t_2(n) + 1$, we define $g_2(n, m) := \min_{t,r} (2n - t + r)$, where the minimum is taken over all integers $t, r \geq 0$ satisfying $t(n - t) + t_2(r) \geq m$. We start by proving the upper bound in Theorem 1.5. First we claim that $g_2(n, m) \leq 2n$. Indeed, for $t = r = \lceil \frac{2m}{3} \rceil$ we have $t(n - t) + t_2(r) = t_3(n) \geq m$, so $g_2(n, m) \leq 2n - t + r = 2n$, as required. Now take t, r such that $t(n - t) + t_2(r) \geq m$ and

$g_2(n, m) = 2n - t + r$. Since $g_2(n, m) \leq 2n$, we have $t \geq r$. Take a complete bipartite graph with sides X of size t and Y of size $n - t$ and add a copy of $T_2(r)$ with sides A, B inside X . The resulting graph G has $t(n - t) + t_2(r) \geq m$ edges. Let H be a chordal subgraph of G . Then $e_H(A, B) \leq |A| + |B| - 1 = r - 1$, $e_H(A, Y) \leq |A| + |Y| - 1$ and $e_H(X \setminus A, Y) \leq |X| - |A| + |Y| - 1$, because each of these bipartite graphs is induced in G , so its intersection with H is a forest. Overall, we got that $e(H) \leq r - 1 + |X| + 2|Y| - 2 = 2n - t + r - 3 = g_2(n, m) - 3$. This shows that $f(n, m) \leq g_2(n, m) - 3$, as required. To prove the lower bound in Theorem 1.5, we prove the following stronger claim.

Theorem 3.1. *Let G be a graph with n vertices and $m \geq t_2(n) + 1$ edges, and let x, y, z be a triangle in G . Then G has a chordal subgraph with at least $g_2(n, m) - 3$ edges which contains the edges xy, xz, yz .*

We need the following facts about the numbers $g_2(n, m)$.

Lemma 3.2. *In the definition of $g_2(n, m)$, we may assume that $-\frac{1}{2} \leq 2t - n - \frac{r}{2} \leq \frac{1}{2}$.*

Proof. Fix t, r which achieve the minimum in the definition of $g_2(n, m)$; so $t(n - t) + t_2(r) \geq m$ and $g_2(n, m) = 2n - t + r$. Put $h(t, r) := 2t - n - \frac{r}{2}$. If $h(t, r) \in \{-\frac{1}{2}, 0, \frac{1}{2}\}$ then we are done. If not, we try replacing (t, r) with $(t - 1, r - 1)$ or $(t + 1, r + 1)$. Suppose first that $h(t, r) \geq 1$. Replace (t, r) with $(t - 1, r - 1)$. We have $2n - (t - 1) + (r - 1) = 2n - t + r$ and $(t - 1)(n - t + 1) + t_2(r - 1) = t(n - t) + t_2(r) - n + 2t - \lfloor \frac{r}{2} \rfloor - 1 \geq t(n - t) + t_2(r) \geq m$, where the penultimate inequality uses $h(t, r) \geq 1$. So $(t - 1, r - 1)$ also achieves the minimum in the definition of $g_2(n, m)$. Moreover, $h(t - 1, r - 1) = 2(t - 1) - n - \frac{r - 1}{2} = h(t, r) - \frac{3}{2}$. So as long as $h(t, r) \geq 1$, we can replace (t, r) with $(t - 1, r - 1)$ and decrease h by $\frac{3}{2}$. At the last step, we decrease h to be in $\{-\frac{1}{2}, 0, \frac{1}{2}\}$.

Similarly, suppose that $h(t, r) \leq -1$. Replace (t, r) with $(t + 1, r + 1)$. We have $2n - (t + 1) + (r + 1) = 2n - t + r$ and $(t + 1)(n - t - 1) + t_2(r + 1) = t(n - t) + t_2(r) + n - 2t + \lceil \frac{r}{2} \rceil - 1 \geq t(n - t) + t_2(r) \geq m$, where the penultimate inequality uses $h(t, r) \leq -1$. So $(t + 1, r + 1)$ also achieves the minimum in the definition of $g_2(n, m)$. Also, $h(t + 1, r + 1) = 2(t + 1) - n - \frac{r + 1}{2} = h(t, r) + \frac{3}{2}$. So as long as $h(t, r) \leq -1$, we can replace (t, r) with $(t + 1, r + 1)$ and increase h by $\frac{3}{2}$. At the last step, we increase h to be in $\{-\frac{1}{2}, 0, \frac{1}{2}\}$. □

Lemma 3.3. *For $n \geq 1$ and $m \geq t_2(n) + 2$, it holds that $g_2(n, m - 1) \geq g_2(n, m) - 1$.*

Proof. Let t, r be such that $t(n - t) + t_2(r) \geq m - 1$ and $2n - t + r = g_2(n, m - 1)$. Since $m - 1 \geq t_2(n) + 1$ we have $r \geq 2$. Thus $t(n - t) + t_2(r + 1) \geq m$, so $g_2(n, m) \leq 2n - t + (r + 1) = g_2(n, m - 1) + 1$, as required. □

Lemma 3.4. *The following holds for every $n \geq 3$.*

1. *If $m \geq t_2(n) + 1$ then $m - n + 1 \geq t_2(n - 2) + 1$ and $g_2(n - 2, m - n + 1) \geq g_2(n, m) - 3$.*
2. *If $m \geq t_2(n) + 2$ then $m - n \geq t_2(n - 2) + 1$ and $g_2(n - 2, m - n) \geq g_2(n, m) - 4$.*

Proof. The first part in both items follows from $t_2(n - 2) = t_2(n) - n + 1$. Let t, r such that $t(n - 2 - t) + t_2(r) \geq m - n + 1$ and $g_2(n - 2, m - n + 1) = 2(n - 2) - t + r$. We have $(t + 1)(n - 1 - t) + t_2(r) = t(n - 2 - t) + n - 1 + t_2(r) \geq m$. Hence, $g_2(n, m) \leq 2n - (t + 1) + r = g_2(n - 2, m - n + 1) + 3$. This proves the first item. For the second item, $g_2(n - 2, m - n) \geq g_2(n - 2, m - n + 1) - 1 \geq g_2(n, m) - 4$, where the first equality uses Lemma 3.3 since $m - n + 1 \geq t_2(n - 2) + 2$. □

Lemma 3.5. *Let $n \geq 1$ and $m \geq t_2(n) + 1$, and let $d \geq 0$ be an integer satisfying $3d \leq g_2(n, m) - 1$. Then $m - d \geq t_2(n - 1) + 1$ and $g_2(n - 1, m - d) \geq g_2(n, m) - 2$.*

Lemma 3.5 is used in the proof of Theorem 3.1 in case one of the vertices x, y, z has degree at most d . We then delete this vertex, apply induction and then add the vertex back with its

two incident edges on the triangle. The induction carries through thanks to the bound $g_2(n - 1, m - d) \geq g_2(n, m) - 2$ from Lemma 3.5. The reason for the assumption $3d \leq g_2(n, m) - 1$ is that if $d(x) + d(y) + d(z) \geq g_2(n, m)$ then the graph consisting of all edges touching x, y, z is a chordal graph with at least $g_2(n, m) - 3$ edges, as required by Theorem 3.1. So we may always assume that one of x, y, z has degree d with $3d \leq g_2(n, m) - 1$.

Proof of Lemma 3.5. First we show that $m - d > t_2(n - 1)$. Set $a := m - t_2(n)$. Since $3d \leq g_2(n, m) - 1$, it is enough to show that

$$g_2(n, m) < 3(m - t_2(n - 1)) + 1 = 3 \cdot \lfloor \frac{n}{2} \rfloor + 3a + 1. \tag{7}$$

For $a = 1, g_2(n, m) = n + \lfloor \frac{n}{2} \rfloor + 2 < 3 \cdot \lfloor \frac{n}{2} \rfloor + 4$, and the last expression equals the right-hand side of (7). Suppose now that $a \geq 2$. Set $k = \lceil \sqrt{\frac{a}{3}} \rceil$, so that $3k^2 \geq a$. Set $t := \lfloor \frac{n}{2} \rfloor + k$ and $r := 4k$. Then $t_2(r) = 4k^2$ and

$$t(n - t) = \left(\lfloor \frac{n}{2} \rfloor + k \right) \left(\lceil \frac{n}{2} \rceil - k \right) \geq t_2(n) - k^2,$$

so $t(n - t) + t_2(r) \geq t_2(n) + 3k^2 \geq t_2(n) + a = m$. Hence, $g_2(n, m) \leq 2n - t + r = 2n - \lfloor \frac{n}{2} \rfloor + 3k \leq 3 \cdot \lfloor \frac{n}{2} \rfloor + 2 + 3k$. So to prove (7), it suffices to show that $3k + 1 < 3a$. As $k \leq \sqrt{\frac{a}{3}} + 1$, it suffices to show that $3 \left(\sqrt{\frac{a}{3}} + 1 \right) < 3a - 1$. Rearranging and squaring, we get the inequality $9a^2 - 27a + 16 > 0$, which holds for all $a \geq 3$. For $a = 2$ we simply note that $k = 1$ and so $3k + 1 = 4 < 6 = 3a$.

Now we show that $g_2(n - 1, m - d) \geq g_2(n, m) - 2$. Fix t, r that achieve the minimum in the definition of $g_2(n, m)$; so $t(n - t) + t_2(r) \geq m$ and

$$g_2(n, m) = 2n - t + r = 3t - 2 \cdot \left(2t - n - \frac{r}{2} \right). \tag{8}$$

By Lemma 3.2, we may assume that $-\frac{1}{2} \leq 2t - n - \frac{r}{2} \leq \frac{1}{2}$. Fix also t', r' such that $t'(n - 1 - t') + t_2(r') \geq m - d$ and $g_2(n - 1, m - d) = 2(n - 1) - t' + r'$. Suppose by contradiction that $g_2(n - 1, m - d) \leq g_2(n, m) - 3$. Then $2(n - 1) - t' + r' \leq 2n - t + r - 3$, so $t' - t \geq r' - r + 1$. Put $c := t' - t$, so that $r' \leq r + c - 1$. Then $t_2(r') \leq t_2(r + c - 1)$. So we can write

$$\begin{aligned} m - d &\leq t'(n - 1 - t') + t_2(r') \leq (t + c)(n - t - c - 1) + t_2(r + c - 1) \\ &= t(n - t) + cn - (2c + 1)t - (c + 1)c + t_2(r + c - 1). \end{aligned} \tag{9}$$

Since (t, r) achieves the minimum in the definition of $g_2(n, m)$, we must have $m - 1 \geq t(n - t) + t_2(r - 1)$, so $t(n - t) \leq m - 1 - t_2(r - 1)$. Plugging this into (9) and rearranging, we get

$$d \geq t + c(2t - n) + (c + 1)c + 1 - t_2(r + c - 1) + t_2(r - 1).$$

Note that

$$t_2(r + c - 1) - t_2(r - 1) \leq \frac{(r + c - 1)^2}{4} - \frac{(r - 1)^2 - 1}{4} = \frac{cr}{2} + \frac{(c - 1)^2}{4}.$$

So we get

$$d \geq t + c \left(2t - n - \frac{r}{2} \right) + (c + 1)c + 1 - \frac{(c - 1)^2}{4}. \tag{10}$$

We now complete the proof by considering the three possible values of $2t - n - \frac{r}{2}$. Suppose first that $2t - n - \frac{r}{2} = 0$. Then $g_2(n, m) = 3t$ by (8). By (10), we have $d \geq t + (c + 1)c + 1 - \frac{(c - 1)^2}{4} = t + \frac{3}{4}(c + 1)^2 \geq t = g_2(n, m)/3$, in contradiction to our assumption on d .

Suppose now that $2t - n - \frac{r}{2} = \frac{1}{2}$. Then $g_2(n, m) = 3t - 1$ by (8). By (10), we have $d \geq t + \frac{c}{2} + (c + 1)c + 1 - \frac{c^2 - 2c + 1}{4} = t + \frac{3c^2}{4} + 2c + \frac{3}{4} > t - 1$, where the last inequality holds for every c . So $d \geq t$ and hence $3d \geq 3t \geq g_2(n, m)$, a contradiction.

Finally, suppose that $2t - n - \frac{r}{2} = -\frac{1}{2}$. Then $g_2(n, m) = 3t + 1$ by (8). By (10), we have $d \geq t - \frac{c}{2} + (c + 1)c + 1 - \frac{c^2 - 2c + 1}{4} = t + \frac{3c^2}{4} + c + \frac{3}{4} > t$, where the last inequality holds for every c . So $d \geq t + 1$ and hence $3d \geq 3t + 3 \geq g_2(n, m)$, a contradiction. \square

Proof of Theorem 3.1. The proof is by induction on n . The base cases $n = 1, 2$ are trivial because for these n there is no graph on n vertices with $t_2(n) + 1$ edges. The case $n = 3$ is also easy to verify. So from now on let $n \geq 4$. Let x, y, z be a triangle in G . We consider several cases. After dealing with each case, we will assume in all subsequent cases that this case does not hold.

Case 1: $d(x) + d(y) + d(z) \geq g_2(n, m)$. In this case, take H to be the graph consisting of all edges touching x, y, z . This graph is chordal and clearly contains the edges of the triangle x, y, z . Also, $e(H) = d(x) + d(y) + d(z) - 3 \geq g_2(n, m) - 3$, as required.

Case 2: There are distinct $u, v \in \{x, y, z\}$ such that $d(u) + d(v) \leq n$. Without loss of generality, suppose that $u = x, v = y$. Let $G' = G - \{x, y\}$. Then $e(G') = e(G) - d(x) - d(y) + 1 \geq m - n + 1$. By the induction hypothesis (applied to any arbitrary triangle in G'), G' contains a chordal subgraph H' with $e(H') \geq g_2(n - 2, m - n + 1) - 3 \geq g_2(n, m) - 6$, by Lemma 3.4. Let $H := H' + \{xy, xz, yz\}$. Then H is chordal, contains the edges of the triangle x, y, z , and satisfies $e(H) = e(H') + 3 \geq g_2(n, m) - 3$.

We claim that if cases 1-2 do not hold then $m \geq t_2(n) + 2$. Indeed, suppose by contradiction that $m = t_2(n) + 1$. We have $g_2(n, t_2(n) + 1) = 2n - \lfloor \frac{n}{2} \rfloor + 2$. Since case 1 does not hold, $d(x) + d(y) + d(z) \leq g_2(n, t_2(n) + 1) - 1 \leq \frac{3n}{2} + 1$. But then there are distinct $u, v \in \{x, y, z\}$ with $d(u) + d(v) \leq \frac{2}{3} \cdot (\frac{3n}{2} + 1) \leq n + \frac{2}{3}$. Hence $d(u) + d(v) \leq n$, contradicting that case 2 does not hold. So $m \geq t_2(n) + 2$.

Case 3: There are distinct $u, v \in \{x, y, z\}$ such that the edge uv is on exactly one triangle (namely the triangle x, y, z). Without loss of generality, suppose that $u = x, v = y$. Since case 2 does not hold, we have $d(x) + d(y) \geq n + 1$. On the other hand, since xy is on exactly one triangle, it must be that $d(x) + d(y) = n + 1, N(x) \cap N(y) = \{z\}$ and every vertex in $V(G) \setminus \{x, y, z\}$ is adjacent to exactly one of x, y . Let $G' = G - \{x, y\}$. Then $e(G') = e(G) - d(x) - d(y) + 1 = m - n$. By Lemma 3.4, $e(G') \geq t_2(v(G')) + 1$, which means that G' contains a triangle. By the induction hypothesis (applied to an arbitrary triangle in G'), G' contains a chordal subgraph H' with $e(H') \geq g_2(n - 2, m - n) - 3 \geq g_2(n, m) - 7$, by Lemma 3.4. So it is enough to show that G contains a chordal graph H which contains the edges of the triangle x, y, z and satisfies $e(H) = e(H') + 4$. Suppose first that z is isolated in H' . We claim that there exists $w \in N_G(z) \setminus \{x, y\}$. Indeed, if not, then $d_G(z) = 2$. Also, as $d(x) + d(y) = n + 1$, we have $d(x) \leq \lfloor \frac{n+1}{2} \rfloor$ or $d(y) \leq \lfloor \frac{n+1}{2} \rfloor$. Suppose this holds for y . Then $d(y) + d(z) \leq \lfloor \frac{n+1}{2} \rfloor + 2 \leq n$ for $n \geq 4$, so case 2 holds, contradiction. This proves our claim that there exists $w \in N_G(z) \setminus \{x, y\}$. Now take $H = H' + \{zw, xy, xz, yz\}$. By adding the vertices in the order z, x, y , we always add a simplicial vertex. Hence, H is chordal.

Suppose now that z is not isolated in H' , and let $w \in V(G) \setminus \{x, y, z\}$ such that $zw \in E(H')$. As mentioned above, w is adjacent in G to either x or y ; without loss of generality it is adjacent to x . Take $H = H' + \{xw, xy, xz, yz\}$. By adding to H' first x and then y , we always add a simplicial vertex. Hence, H is chordal.

Case 4: Cases 1-3 do not hold. Since case 1 does not hold, we have $d(x) + d(y) + d(z) \leq g_2(n, m) - 1$. Assume that $d := d(x) \leq d(y) \leq d(z)$; then $3d \leq g_2(n, m) - 1$. Since case 3 does not hold, there exists $w \in V(G) \setminus \{x\}$ which is a common neighbour of y, z . Set $G' = G - \{x\}$. We have $e(G') = m - d$. By the induction hypothesis, G' has a chordal subgraph H' which contains the

edges of the triangle y, z, w and satisfies $e(H') \geq g_2(n - 1, m - d) - 3 \geq g_2(n, m) - 5$, by Lemma 3.5. Let $H = H' + \{xy, xz\}$. Then H is chordal, contains the edges of the triangle x, y, z , and satisfies $e(H) = e(H') + 2 \geq g_2(n, m) - 3$, as required. \square

4. Proof of Theorem 1.2

For convenience, put $g_3(n) := 3n - \lceil \frac{n}{3} \rceil + 2$. We prove Theorem 1.2 in the following stronger form.

Theorem 4.1. *Let G be a graph with n vertices and $t_3(n) + 1$ edges, and let $X = \{x_1, x_2, x_3, x_4\}$ be a 4-clique in G . Then G has a chordal subgraph with at least $g_3(n) - 6$ edges which contains the edges of the clique X .*

We need some simple facts on the numbers $t_3(n)$ and $g_3(n)$.

Lemma 4.2. *For every $n \geq 5$, it holds that $t_3(n) - t_3(n - 1) = \lfloor \frac{2n}{3} \rfloor$, $t_3(n) - t_3(n - 2) = \lfloor \frac{4n}{3} \rfloor - 1$, $t_3(n) - t_3(n - 3) = 2n - 3$, $t_3(n) - t_3(n - 4) = g_3(n) - 7$.*

Proof. For $i = 1, 2, 3$, $T_3(n - i)$ is obtained from $T_3(n)$ by deleting one vertex from each of the i largest classes of $T_3(n)$. For $i = 1$, the deleted vertex has degree $\lfloor \frac{2n}{3} \rfloor$. For $i = 2$, the sum of degrees of the deleted vertices is $\lfloor \frac{4n}{3} \rfloor$, and these vertices are adjacent. For $i = 3$, the sum of degrees of the deleted vertices is $2n$, and they form a triangle. Finally, $t_3(n) - t_3(n - 4) = t_3(n) - t_3(n - 1) + t_3(n - 1) - t_3(n - 4) = \lfloor \frac{2n}{3} \rfloor + 2(n - 1) - 3 = 3n - \lceil \frac{n}{3} \rceil - 5 = g_3(n) - 7$. \square

Lemma 4.3. *For every $n \geq 5$, it holds that $g_3(n) - g_3(n - 1) = 3$ if $n \equiv 0, 2 \pmod{3}$ and $g_3(n) - g_3(n - 1) = 2$ if $n \equiv 1 \pmod{3}$. Hence, $g_3(n) - g_3(n - 2) \leq 6$, $g_3(n) - g_3(n - 3) = 8$ and $g_3(n) - g_3(n - 4) \leq 11$.*

Proof. $g_3(n) - g_3(n - 1) = 3 - \lceil \frac{n}{3} \rceil + \lceil \frac{n-1}{3} \rceil$, and it is easy to see that $\lceil \frac{n}{3} \rceil - \lceil \frac{n-1}{3} \rceil$ equals 0 if $n \equiv 0, 2 \pmod{3}$ and equals 1 if $n \equiv 1 \pmod{3}$. \square

We also need the following simple lemma saying that in a graph with $t_3(n) + 1$ edges, we can find two 4-cliques sharing 3 vertices. This fact is originally due to Dirac [2]. For completeness, we include a proof.

Lemma 4.4. *A graph G with $n \geq 5$ vertices and $t_3(n) + 1$ edges has two 4-cliques sharing 3 vertices.*

Proof. The proof is by induction on n . For the base case $n = 5$, take a 4-clique and notice that the remaining vertex must send at least 3 edges to this 4-clique as $t_3(5) + 1 = 9$. Let $n \geq 6$. Take a 4-clique x_1, \dots, x_4 . If there is a vertex outside of x_1, \dots, x_4 which has three neighbours in x_1, \dots, x_4 then we are done. Else, $d(x_1) + \dots + d(x_4) \leq 2n + 4$, so there is $i \in [4]$ such that $d(x_i) \leq \lfloor \frac{n}{2} \rfloor + 1 \leq \lfloor \frac{2n}{3} \rfloor$, where the last inequality holds for all $n \geq 6$. So $G' := G - x_i$ has at least $t_3(n - 1) + 1$ edges, and we can apply induction. \square

Proof of Theorem 4.1. The proof is by induction on n . The claim is trivial if $n \leq 4$ so suppose that $n \geq 5$. We proceed by a sequence of claims.

Claim 4.5. *We may assume that there exists $y \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to at least 3 of the vertices x_1, \dots, x_4 .*

Proof of Claim 4.5. Suppose that there is no such y . We will show that the assertion of Theorem 4.1 holds. For $0 \leq j \leq 2$, let A_j be the set of vertices in $V(G) \setminus \{x_1, \dots, x_4\}$ which are adjacent to j of the vertices x_1, \dots, x_4 , and let $a_j = |A_j|$. Then $a_0 + a_1 + a_2 = n - 4$ and $a_1 + 2a_2 = d(x_1) + \dots + d(x_4) - 12$. So

$$d(x_1) + \dots + d(x_4) = n + 8 + a_2 - a_0. \tag{11}$$




Figure 2. Illustrations to proofs of claims. The red edges are added after applying induction. The dashed edges are deleted.

We consider three cases:

Case 1: $0 < a_2 \leq n - 5$. Then $d(x_1) + \dots + d(x_4) \leq 2n + 3$ by (11). Let G' be the graph obtained from G by deleting x_2, x_3, x_4 and all edges touching x_1 . Then $v(G') = n - 3$ and $e(G') = e(G) - (d(x_1) + \dots + d(x_4)) + 6 \geq e(G) - 2n + 3 = t_3(n) + 1 - 2n + 3 = t_3(n - 3) + 1$. In particular, $n - 3 \geq 4$ (else G' cannot have more than $t_3(n - 3)$ edges). By the induction hypothesis (applied to an arbitrary 4-clique in G'), G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$. Take $z \in A_2$ and assume without loss of generality that z is adjacent to x_1, x_2 . Let $H = H' + \{x_1z, x_2z\} + \{x_ix_j : 1 \leq i < j \leq 4\}$. Then $e(H) = e(H') + 8 \geq g_3(n - 3) + 8 - 6 = g_3(n) - 6$. Also, H is chordal: adding x_1z to H' keeps it chordal because x_1 is isolated in H' , and then by adding x_2, x_3, x_4 in this order we always add a simplicial vertex.

Case 2: $a_2 = n - 4$. Then every vertex outside of $\{x_1, \dots, x_4\}$ is adjacent to two of the vertices x_1, \dots, x_4 . By (11), $d(x_1) + \dots + d(x_4) = 2n + 4$. Pick an arbitrary $v \in V(G) \setminus \{x_1, \dots, x_4\}$ and suppose without loss of generality that v is adjacent to x_1 . Let G' be the graph obtained from G by deleting x_2, x_3, x_4 and all edges touching x_1 except for x_1v . Then $v(G') = n - 3$ and $e(G') = e(G) - (d(x_1) + \dots + d(x_4)) + 7 = e(G) - 2n + 3 = t_3(n - 3) + 1$. Pick an arbitrary 4-clique y_1, \dots, y_4 in G' . By the induction hypothesis, G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$ such that H' contains the edges of the clique y_1, \dots, y_4 . Each y_i has two neighbours in x_1, \dots, x_4 . By pigeonhole, two of the y_i 's have a common neighbour. Without loss of generality, we may assume that y_1, y_2 are both adjacent to x_k for some $k \in [4]$. Suppose that y_1 is also adjacent to x_ℓ . Let $H := H' - x_1v + \{x_ky_1, x_ky_2, x_\ell y_1\} + \{x_ix_j : 1 \leq i < j \leq 4\}$, see Figure 2(a). Then $e(H) \geq e(H') + 8$ (since we add 9 edges and delete at most 1), so $e(H) \geq g_3(n - 3) + 8 - 6 = g_3(n) - 6$. Also, $H' - x_1v$ is chordal because x_1 is a leaf or an isolated vertex in H' . Now observe that H is chordal: adding first x_k , then x_ℓ and then the other two vertices among x_1, \dots, x_4 , we add a simplicial vertex at each step.

Case 3: $a_2 = 0$. Then there are at most $n - 4$ edges between x_1, \dots, x_4 and $V(G) \setminus \{x_1, \dots, x_4\}$. Also, $d(x_1) + \dots + d(x_4) \leq n + 8$ by (11). Let G' be the graph obtained from G by deleting x_2, x_3, x_4 and all edges touching x_1 . Then $v(G') = n - 3$ and $e(G') \geq e(G) - (n + 2) = t_3(n) + 1 - (n + 2) = t_3(n - 3) + 1 + n - 5$. Note that $n \geq 8$ because else $e(G) \leq \binom{n-4}{2} + n - 4 + 6 < t_3(n)$ (where the last inequality holds for $n \leq 7$), a contradiction. So $v(G') \geq 5$. By Lemma 4.4, there are distinct $v_1, v_2, w_1, w_2, w_3 \in V(G')$ such that v_i, w_1, w_2, w_3 is a 4-clique for $i = 1, 2$. Let G'' be the graph obtained from G' by deleting all edges touching v_1 . Then $v(G'') = n - 3$ and $e(G'') \geq e(G') - (n - 5)$ because v_1 is not adjacent in G' to x_1 or to itself, leaving $n - 5$ potential neighbours. So $e(G'') \geq t_3(n - 3) + 1$. By the induction hypothesis, G'' contains a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$, such that H' contains the edges of the clique v_2, w_1, w_2, w_3 .

Let $H = H' + \{v_1 w_i : 1 \leq i \leq 3\} + \{x_i x_j : 1 \leq i < j \leq 4\}$. Then $e(H) = e(H') + 9 \geq g_3(n) - 6$, and it is easy to check that H is chordal. \square

Claim 4.6. *We may assume that for every $i \in [4]$, the following holds. If $d(x_i) \geq \lfloor \frac{2n}{3} \rfloor + 1$, then there is $y \in V(G) \setminus \{x_1, \dots, x_4\}$ such that y is adjacent to x_i and to at least two vertices in $X \setminus \{x_i\}$.*

Proof of Claim 4.6. Without loss of generality, $i = 1$. Suppose that $d(x_1) \geq \lfloor \frac{2n}{3} \rfloor + 1$ but there is no y as above. We will show that the assertion of Theorem 4.1 holds. For $0 \leq j \leq 3$, let A_j be the set of vertices $z \in V(G) \setminus \{x_2, x_3, x_4\}$ which are adjacent to exactly j of the vertices x_2, x_3, x_4 , and let $a_j = |A_j|$. Note that $x_1 \in A_3$. By Claim 4.5, we may assume that there exists $z \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to three of the vertices x_1, \dots, x_4 . By assumption, these three vertices cannot include x_1 , so we must have $z \in A_3$. Hence, $A_3 \setminus \{x_1\} \neq \emptyset$.

We have $a_0 + a_1 + a_2 + a_3 = n - 3$ and $a_1 + 2a_2 + 3a_3 = d(x_2) + d(x_3) + d(x_4) - 6$. So $a_2 + 2a_3 = d(x_2) + d(x_3) + d(x_4) - n - 3 + a_0$. By assumption, there is no $y \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to x_1 and belongs to $A_2 \cup A_3$. Hence, $d(x_1) \leq n - a_2 - a_3 = 2n - d(x_2) - d(x_3) - d(x_4) + 3 + a_3 - a_0$. So we get

$$d(x_1) + \dots + d(x_4) \leq 2n + 3 + a_3 - a_0. \tag{12}$$

Pick a set of edges F as follows:

- Suppose first that $a_0 = 0$. We claim that $A_1 \cup A_2 \neq \emptyset$. Indeed, if $A_1 \cup A_2 = \emptyset$ then $A_3 = V(G) \setminus \{x_2, x_3, x_4\}$. But then $N(x_1) = \{x_2, x_3, x_4\}$ so $d(x_1) = 3$. On the other hand, $d(x_1) \geq \lfloor \frac{2n}{3} \rfloor + 1$ by assumption. So $\lfloor \frac{2n}{3} \rfloor \leq 2$, which is false for every $n \geq 5$. Now pick an arbitrary $v \in A_1 \cup A_2$, and suppose without loss of generality that v is adjacent to x_4 . Take F to be the set of all edges between x_4 and $N(x_4) \setminus (A_3 \cup \{x_2, x_3, v\})$. Note that $|F| = d(x_4) - a_3 - 3$.
- If $a_0 \geq 1$ then F is the set of all edges between x_4 and $N(x_4) \setminus (A_3 \cup \{x_2, x_3\})$. Then $|F| = d(x_4) - a_3 - 2$.

In both cases we have $|F| \leq d(x_4) - a_3 - 3 + a_0$. Let G' be the graph obtained from G by deleting x_1, x_2, x_3 and all edges in F . So $v(G') = n - 3$. To count the deleted edges, note that there are $d(x_1) + d(x_2) + d(x_3) - 3$ edges touching x_1, x_2, x_3 , and that the edges in F do not touch x_1, x_2, x_3 . So the number of deleted edges is

$$d(x_1) + d(x_2) + d(x_3) - 3 + |F| \leq d(x_1) + \dots + d(x_4) - a_3 + a_0 - 6 \leq 2n - 3,$$

where the last inequality uses (12). So $e(G') \geq e(G) - 2n + 3 = t_3(n) + 1 - 2n + 3 = t_3(n - 3) + 1$. By the induction hypothesis, G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$. Define a subgraph H'' of G' as follows:

- If there is $z \in A_3 \setminus \{x_1\}$ such that $x_4 z \in E(H')$, then $H'' = H'$;
- Else, pick any $z \in A_3 \setminus \{x_1\}$ and set $H'' = H' - x_4 v + x_4 z$.

We claim that $e(H'') \geq e(H')$. In the first item this is obvious. Suppose we are in the second item. By the definition of F , every neighbour of x_4 in G' belongs to $\{v\} \cup A_3 \setminus \{x_1\}$. Since we are in the second item, the only possible neighbour of x_4 in H' is v . So indeed $e(H'') \geq e(H')$. Also, H'' is chordal; in the second case, this is because x_4 is a leaf in both H' and H'' .

By the definition of H'' , there is $z \in A_3 \setminus \{x_1\}$ such that $x_4 z \in E(H'')$. Now put $H = H'' + \{x_2 z, x_3 z\} + \{x_i x_j : 1 \leq i < j \leq 4\}$, see Figure 2(b). Then $e(H) = e(H'') + 8 \geq g_3(n - 3) + 8 - 6 = g_3(n) - 6$. Also, H is chordal: adding the vertices x_2, x_3, x_1 in this order, we always add a simplicial vertex. This proves Claim 4.6. \square

We now continue with the proof of the theorem. For $i \in [4]$, we say that x_i is *deletable* if $d(x_i) \leq \lfloor \frac{2n}{3} \rfloor$, and there is $y \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to x_j for all $j \in [4] \setminus \{i\}$.

Claim 4.7. *We may assume that there is no deletable x_i .*

Proof. Suppose that x_i is deletable. Without loss of generality, assume that $i = 1$. Let $G' = G - x_1$. Then $e(G') = e(G) - d(x_1) \geq t_3(n) + 1 - \lfloor \frac{2n}{3} \rfloor = t_3(n - 1) + 1$. Let y be a common neighbour of x_2, x_3, x_4 . By the induction hypothesis, G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 1) - 6$, such that H' contains the edges of the clique x_2, x_3, x_4, y . Take $H = H' + \{x_1x_2, x_1x_3, x_1x_4\}$. Then H is chordal and $e(H) = e(H') + 3 \geq g_3(n - 1) + 3 - 6 \geq g_3(n) - 6$. \square

Claim 4.8. We may assume that $d(x_1) + \dots + d(x_4) \leq g_3(n) - 1$.

Proof. Else, take H to be the subgraph of G consisting of all edges which touch x_1, \dots, x_4 . Then H is chordal and $e(H) = d(x_1) + \dots + d(x_4) - 6 \geq g_3(n) - 6$. \square

Claim 4.9. We may assume that there is no 5-clique containing x_1, \dots, x_4 .

Proof. Suppose that x_1, \dots, x_4, y is a 5-clique. If $d(x_i) \leq \lfloor \frac{2n}{3} \rfloor$ then x_i is deletable. Assuming no x_i is deletable, we have $d(x_1) + \dots + d(x_4) \geq 4 \cdot (\lfloor \frac{2n}{3} \rfloor + 1) \geq 2n + \lfloor \frac{2n}{3} \rfloor + 2 = g_3(n)$, so we are done by Claim 4.8. \square

Claim 4.10. We may assume that for every $i \in [4]$, $\sum_{j \neq i} d(x_j) \leq 2n$.

Proof. Suppose that $d(x_1) + d(x_2) + d(x_3) \geq 2n + 1$ (the proof for each other i is symmetric). By Claim 4.8, we may assume that $d(x_4) \leq g_3(n) - 1 - (2n + 1) = n - \lceil \frac{n}{3} \rceil = \lfloor \frac{2n}{3} \rfloor$. By Claim 4.7, we may assume that x_4 is not deletable. Then x_1, x_2, x_3 have no common neighbour except for x_4 . But as $d(x_1) + d(x_2) + d(x_3) \geq 2n + 1$, this means that every vertex in $V(G) \setminus \{x_1, \dots, x_4\}$ is adjacent to exactly two of the vertices x_1, x_2, x_3 . Let $G' = G - \{x_1, \dots, x_4\}$. Then $v(G') = n - 4$ and $e(G') = e(G) - (d(x_1) + \dots + d(x_4)) + 6 \geq e(G) - g_3(n) + 7 = t_3(n) + 1 - g_3(n) + 7 = t_3(n - 4) + 1$ by Lemma 4.2. Fix an arbitrary 4-clique w_1, \dots, w_4 in G' . By the induction hypothesis, G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 4) - 6$ such that H' contains the edges of the clique w_1, \dots, w_4 . Each w_i is adjacent to two of the vertices x_1, x_2, x_3 . By the pigeonhole principle and by symmetry, we may assume that each of w_1, w_2 is adjacent to x_1, x_2 . Moreover, w_3 must be adjacent to x_1 or x_2 ; say to x_1 . Take $H = H' + \{x_1w_1, x_1w_2, x_1w_3, x_2w_1, x_2w_2\} + \{x_i x_j : 1 \leq i < j \leq 4\}$, see Figure 2(c). Then $e(H) = e(H') + 11 \geq g_3(n - 4) + 11 - 6 \geq g_3(n) - 6$. Also, H is chordal: adding the vertices x_1, x_2, x_3, x_4 in this order, we always add a simplicial vertex. This proves Claim 4.10. \square

By Claim 4.5, we may assume that there is $y_0 \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to at least 3 of the vertices x_1, \dots, x_4 ; say to x_2, x_3, x_4 . By Claim 4.7, we may assume that x_i is not deletable for any $i \in [4]$. Since x_1 is not deletable, $d(x_1) \geq \lfloor \frac{2n}{3} \rfloor + 1$. By Claim 4.6, there is $z_0 \in V(G) \setminus \{x_1, \dots, x_4\}$ which is adjacent to x_1 and to at least two of the vertices x_2, x_3, x_4 ; without loss of generality, z_0 is adjacent to x_3, x_4 . Since x_2 is not deletable, we have $d(x_2) \geq \lfloor \frac{2n}{3} \rfloor + 1$. Also, we may assume that $y_0 \neq z_0$ because else we would have a 5-clique containing x_1, \dots, x_4 .

Claim 4.11. We may assume that $d(x_i) \leq \lfloor \frac{2n}{3} \rfloor$ for each $i = 3, 4$.

Proof. By Claim 4.10, we may assume that $d(x_i) \leq 2n - d(x_1) - d(x_2) \leq 2n - 2 \cdot (\lfloor \frac{2n}{3} \rfloor + 1) \leq \lfloor \frac{2n}{3} \rfloor$. \square

If $d(y_0) \leq \lfloor \frac{2n}{3} \rfloor$ then we can delete y_0 , apply induction to the remaining graph to find a chordal subgraph H' with at least $g_3(n - 1) - 6$ edges which contains the edges of the clique x_1, \dots, x_4 and then add the edges x_2y_0, x_3y_0, x_4y_0 to H' to conclude the proof. So suppose that $d(y_0) \geq \lfloor \frac{2n}{3} \rfloor + 1$. Let $v \notin \{x_3, x_4\}$ be a common neighbour of x_2 and y_0 . Such a vertex v exists because $d(x_2) + d(y_0) \geq 2 \cdot (\lfloor \frac{2n}{3} \rfloor + 1) \geq n + 3$, where the last inequality holds for every $n \geq 5$. Note that $v \neq x_1$ because x_1 is not adjacent to y_0 , as otherwise x_1, \dots, x_4, y_0 would be a 5-clique. Similarly, $v \neq z_0$ because v is adjacent to x_2 and z_0 is adjacent to x_1, x_3, x_4 (so otherwise x_1, \dots, x_4, z_0 would be a 5-clique).

Suppose first that x_2, y_0, v have no common neighbour. Then $d(x_2) + d(y_0) + d(v) \leq 2n$. Let $G' = G - \{x_2, y_0, v\}$. Then $e(G') = e(G) - d(x_2) - d(y_0) - d(v) + 3 \geq t_3(n) + 1 - 2n + 3 =$




Figure 3. Applying induction after deleting certain vertices. The added edges are red.

$t_3(n - 3) + 1$. By the induction hypothesis, G' contains a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$ such that H' contains the edges of the clique x_1, x_3, x_4, z_0 . Take $H = H' + \{x_2x_1, x_2x_3, x_2x_4, y_0x_2, y_0x_3, y_0x_4, vx_2, vy_0\}$, see Figure 3(a). Then $e(H) = e(H') + 8 \geq g_3(n - 3) + 8 - 6 = g_3(n) - 6$. Also, H is chordal: adding the vertices x_2, y_0, v in this order, we always add a simplicial vertex.

Suppose now that x_2, y_0, v have a common neighbour w . First, suppose that $w \in \{x_3, x_4\}$, say $w = x_3$. Let $G' = G - \{x_1, x_4\}$. We have $d(x_1) + d(x_4) \leq 2n - d(x_2) \leq 2n - \lfloor \frac{2n}{3} \rfloor - 1 \leq \lfloor \frac{4n}{3} \rfloor$, where the first inequality is by Claim 4.10. So $e(G') = e(G) - d(x_1) - d(x_4) + 1 \geq t_3(n) + 1 - \lfloor \frac{4n}{3} \rfloor + 1 = t_3(n - 2) + 1$. By the induction hypothesis, G' has a chordal subgraph H' with $e(H') \geq g_3(n - 2) - 6$ such that H' contains the edges of the clique x_2, y_0, v, x_3 . Take $H = H' + \{x_4x_2, x_4x_3, x_4y_0, x_1x_2, x_1x_3, x_1x_4\}$, see Figure 3(b). Then $e(H) = e(H') + 6 \geq g_3(n - 2) + 6 - 6 \geq g_3(n) - 6$. Also, H is chordal: adding x_4 and then x_1 , we always add a simplicial vertex. Now suppose that $w \notin \{x_3, x_4\}$. Let $G' = G - \{x_1, x_3, x_4\}$. By Claims 4.10 and 4.11, we have $d(x_1) + d(x_3) \leq 2n - d(x_2) \leq \frac{4n}{3}$ and $d(x_4) \leq \frac{2n}{3}$, so $d(x_1) + d(x_3) + d(x_4) \leq 2n$. Hence, $e(G') = e(G) - d(x_1) - d(x_3) - d(x_4) + 3 \geq t_3(n) + 1 - 2n + 3 = t_3(n - 3) + 1$. By the induction hypothesis, G' has a chordal subgraph H' with $e(H') \geq g_3(n - 3) - 6$ such that H' contains the edges of the clique x_2, y_0, v, w . Now take $H = H' + \{x_3y_0, x_4y_0\} + \{x_ix_j : 1 \leq i < j \leq 4\}$, see Figure 3(c). Then $e(H) = e(H') + 8 \geq g_3(n - 3) + 8 - 6 = g_3(n) - 6$. Also, H is chordal: adding the vertices x_3, x_4, x_1 in this order, we always add a simplicial vertex. This completes the proof of the theorem. \square

5. Concluding remarks

In this paper we study the maximal value $f(n, m)$ such that every graph with n vertices and m edges has a chordal subgraph with at least $f(n, m)$ edges. We determine this function asymptotically for all m and exactly for $m \leq t_3(n) + 1$. Our results suggest the following conjecture on the value of $f(n, m)$ for general m .

Conjecture 5.1. *Let $k \geq 2, n \geq 1$ and $t_k(n) + 1 \leq m \leq t_{k+1}(n)$. Then*

$$f(n, m) = \min_{t,r} (kn - t + r) - \binom{k + 1}{2},$$

where the minimum is taken over all $t, r \geq 0$ satisfying $t_{k-1}(n - t) + t(n - t) + t_2(r) \geq m$.

It seems very likely that some progress on this conjecture can be achieved using our techniques together with a careful case analysis, but it would be interesting to find a proof which avoids such a case analysis as much as possible.

Note added:

Zachary Hunter (personal communication) very recently improved the error term in Theorem 1.3 from $O(\sqrt{n})$ to $O(1)$ in the special case $m = t_k(n) + 1$, proving that $f(n, t_k(n) + 1) = (k - 1/k)n - O_k(1)$.

Acknowledgements

The authors would like to thank the anonymous referee for a careful reading of the paper and useful suggestions.

Funding

Research supported in part by SNSF grant 200021_196965

References

- [1] Bollobás, B. and Nikiforov, V. (2005) The sum of degrees in cliques. *Electron. J. Combin.* **12**.
- [2] Dirac, G. (1963) Extensions of Turán's theorem on graphs. *Acta Math. Hung.* **14**(3-4) 417–422.
- [3] Edwards, C. (1976) Complete subgraphs with largest sum of vertex degrees. In *Combinatorics (Proc. Fifth Hungarian Colloq.)*. Keszthely, pp. 293–306, **1**.
- [4] Edwards, C. (1977) The largest vertex degree sum for a triangle in a graph. *Bull. London Math. Soc.* **9**(2) 203–208.
- [5] Edwards, C. (1977) A lower bound for the largest number of triangles with a common edge. *Unpublished manuscript*.
- [6] Enright, J. and Kondrak, G. (2011) The application of chordal graphs to inferring phylogenetic trees of languages.
- [7] Erdős, P. (1962) On a theorem of Rademacher-Turán. *Illinois J. Math.* **6**(122-127) 1–3.
- [8] Erdős, P., Furedi, Z., Gould, R. J. and Gunderson, D. S. (1995) Extremal graphs for intersecting triangles. *J. Comb. Theory Ser. B* **64**(1) 89–100.
- [9] Erdős, P., Gyárfás, A., Ordman, E. T. and Zalcstein, Y. (1989) The size of chordal, interval and threshold subgraphs. *Combinatorica* **9**(3) 245–253.
- [10] Erdős, P. and Laskar, R. (1985) A note on the size of a chordal subgraph. *Congress. Numer.* **48** 81–86.
- [11] Erdős, P. (1969) On the number of complete subgraphs and circuits contained in graphs. *Časopis pro pěstování matematiky* **94**(3) 290–296.
- [12] Faudree, R. J. (1992) Complete subgraphs with large degree sums. *J. Graph Theory* **16**(4) 327–334.
- [13] Golombic, M. C. (2004) *Algorithmic Graph Theory and Perfect Graphs*. Elsevier.
- [14] Gyárfás, A. (2013) Problems and memories. *arXiv preprint arXiv:1307.1768*.
- [15] Khadžiivanov, N. G. and Nikiforov, V. S. (1979) Solution of a problem of P. Erdős about the maximum number of triangles with a common edge in a graph. *Doklady Bolgarskoj Akademii Nauk. Comptes Rendus de l'Académie Bulgare des Sciences* **32**(10) 1315–1318, (In Russian).
- [16] Lovász, L. and Simonovits, M. (1983) On the number of complete subgraphs of a graph II. In *Studies in Pure Mathematics*, Birkhauser, Basel, pp. 459–495.
- [17] Vandenberghe, L. and Andersen, M. S. (2015) Chordal graphs and semidefinite optimization. *Found. Trends Optim.* **1**(4) 241–433.