

## THE COMMON LIMIT OF A QUADRUPLE SEQUENCE AND THE HYPERGEOMETRIC FUNCTION $F_D$ OF THREE VARIABLES

TAKAYUKI KATO AND KEIJI MATSUMOTO

*Dedicated to Professor Masaaki Yoshida on his sixtieth birthday*

**Abstract.** We study a quadruple sequence and express its common limit by Lauricella's hypergeometric function  $F_D(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, 1; z_1, z_2, z_3)$  of three variables. We give a functional equation of  $F_D$ , which is the key to get our expression of the common limit.

### §1. Introduction

For two positive real numbers  $a_0$  and  $b_0$  with  $a_0 \geq b_0$ , the double sequence  $\{a_n\}$  and  $\{b_n\}$  given as

$$a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n},$$

has a common limit, which is called the arithmetic-geometric mean  $M(a_0, b_0)$  of  $a_0$  and  $b_0$ . It is shown by C. F. Gauss that  $M(a_0, b_0)$  can be expressed by the hypergeometric function:

$$\frac{a_0}{M(a_0, b_0)} = \frac{1}{M(1, x)} = F\left(\frac{1}{2}, \frac{1}{2}, 1; 1 - x^2\right),$$

where  $x = b_0/a_0$ .

In this paper, we study a quadruple sequence  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$  and  $\{d_n\}$

---

Received August 7, 2007.

Revised June 24, 2008.

Accepted February 2, 2009.

2000 Mathematics Subject Classification: Primary 26A18; Secondary 33C65.

given by

$$(a_0, b_0, c_0, d_0) = (a, b, c, d), \quad a \geq b \geq c \geq d \geq 0,$$

$$(1) \quad a_{n+1} = \frac{a_n + b_n + c_n + d_n}{4}, \quad b_{n+1} = \frac{\sqrt{(a_n + d_n)(b_n + c_n)}}{2},$$

$$c_{n+1} = \frac{\sqrt{(a_n + c_n)(b_n + d_n)}}{2}, \quad d_{n+1} = \frac{\sqrt{(a_n + b_n)(c_n + d_n)}}{2}.$$

We can easily see that it has a common limit  $\mu(a, b, c, d)$ . Our main theorem is the expression of  $\mu(a, b, c, d)$  by Lauricella’s hypergeometric function  $F_D$  of three variables:

$$\frac{1}{\mu(1, x_1, x_2, x_3)} = F_D\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}; 1; 1 - x_1^2, 1 - x_2^2, 1 - x_3^2\right)^2.$$

The key for our main theorem is Proposition 1, which is the functional equation of the hypergeometric function  $F_D$  corresponding to the property

$$\frac{a_0}{a_1} \mu\left(1, \frac{b_0}{a_0}, \frac{c_0}{a_0}, \frac{d_0}{a_0}\right) = \mu\left(1, \frac{b_1}{a_1}, \frac{c_1}{a_1}, \frac{d_1}{a_1}\right).$$

It turns out that

$$\mu\left(1, \frac{b_n}{a_n}, \frac{c_n}{a_n}, \frac{d_n}{a_n}\right) F_D\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}; 1; 1 - \left(\frac{b_n}{a_n}\right)^2, 1 - \left(\frac{c_n}{a_n}\right)^2, 1 - \left(\frac{d_n}{a_n}\right)^2\right)^2$$

is independent of  $n$ . This fact implies our main theorem. In order to show Proposition 1, we prepare some essential facts for integrable Pfaffian systems in Section 3 and give the integrable Pfaffian system with respect to  $F_D(\alpha, \beta, \gamma; z)$  of three variables in Fact 4.

C. W. Borchardt considers in [B] the quadruple sequence

$$a_{n+1} = \frac{a_n + b_n + c_n + d_n}{4}, \quad b_{n+1} = \frac{\sqrt{a_n b_n} + \sqrt{c_n d_n}}{2},$$

$$c_{n+1} = \frac{\sqrt{a_n c_n} + \sqrt{b_n d_n}}{2}, \quad d_{n+1} = \frac{\sqrt{a_n d_n} + \sqrt{b_n c_n}}{2},$$

with positive initial terms  $a_0, b_0, c_0, d_0$ . Its common limit  $B(a_0, b_0, c_0, d_0)$  is expressed in terms of period integrals of a hyperelliptic curve  $C$  of genus 2. J. F. Mestre studies the expression of fixed points under the hyperelliptic involution on  $C$  by the initial terms and shows that  $B(a_0, b_0, c_0, d_0)$  can be expressed by the arithmetic-geometric mean  $M(a_0, c_0)$  when  $a_0 = b_0$  and  $c_0 = d_0$ , refer to [M].

J. M. Borwein and P. B. Borwein consider in [BB] two double sequences

$$a_{n+1} = \frac{a_n + 2b_n}{3}, \quad b_{n+1} = \sqrt[3]{b_n \frac{a_n^2 + a_n b_n + b_n^2}{3}},$$

and

$$a_{n+1} = \frac{a_n + 3b_n}{4}, \quad b_{n+1} = \sqrt{b_n \frac{a_n + b_n}{2}};$$

they express their common limits  $M_3(a_0, b_0)$  and  $M_4(a_0, b_0)$  by  $F(\frac{1}{3}, \frac{2}{3}, 1; z)$  and  $F(\frac{1}{4}, \frac{3}{4}, 1; z)$ , respectively. We remark that the expression of  $M_4(a_0, b_0)$  can be obtained by our main theorem as a special case  $b_0 = c_0 = d_0$ .

As a generalization of  $M_3(a_0, b_0)$ , K. Koike and H. Shiga give a triple sequence and express its common limit by Appell’s hypergeometric function  $F_1(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 1; z_1, z_2)$  of two variables  $z_1, z_2$ , refer to [KS1]. They study an extension of the arithmetic-geometric mean and give its expression by Appell’s hypergeometric function  $F_1$  with different parameters in [KS2].

For other studies related to the arithmetic-geometric mean, refer to [MM].

**Acknowledgment.** The authors express their gratitude to Professor K. Ohara for informing of the correct Pfaffian system with respect to the hypergeometric function  $F_D$  obtained by the system [O].

**§2. The quadruple sequence**

LEMMA 1. *The quadruple sequence  $\{a_n\}, \{b_n\}, \{c_n\}$  and  $\{d_n\}$  given as (1) satisfies*

$$a \geq a_{n-1} \geq a_n \geq b_n \geq c_n \geq d_n \geq d_{n-1} \geq d$$

for any  $n \in \mathbb{N}$ . It has a common limit, which is denoted by  $\mu(a, b, c, d)$ .

*Proof.* We assume  $a_n \geq b_n \geq c_n \geq d_n \geq 0$  for  $n \in \mathbb{N}$ . Then we have

$$\begin{aligned} (4a_{n+1})^2 - (4b_{n+1})^2 &= (a_n - b_n - c_n + d_n)^2 \geq 0, \\ (2b_{n+1})^2 - (2c_{n+1})^2 &= (a_n - b_n)(c_n - d_n) \geq 0, \\ (2c_{n+1})^2 - (2d_{n+1})^2 &= (a_n - d_n)(b_n - c_n) \geq 0, \\ a_{n+1} &= \frac{a_n + b_n + c_n + d_n}{4} \leq \frac{a_n + a_n + a_n + a_n}{4} = a_n, \\ d_{n+1} &= \frac{\sqrt{(a_n + b_n)(c_n + d_n)}}{2} \geq \frac{\sqrt{(d_n + d_n)(d_n + d_n)}}{2} = d_n, \end{aligned}$$

which imply  $a_n \geq a_{n+1} \geq b_{n+1} \geq c_{n+1} \geq d_{n+1} \geq d_n \geq 0$ . Since the sequences  $\{a_n\}$  and  $\{d_n\}$  are monotonous and bounded, they converge. Since

$$\begin{aligned} a_{n+1} - d_{n+1} &= \frac{1}{4}(\sqrt{a_n + b_n} - \sqrt{c_n + d_n})^2 \leq \frac{1}{4}(\sqrt{2a_n} - \sqrt{2d_n})^2 \\ &= \frac{1}{2}(a_n + d_n - 2\sqrt{a_n d_n}) \leq \frac{a_n - d_n}{2} \leq \frac{a - d}{2^{n+1}}, \end{aligned}$$

we have  $\lim_{n \rightarrow \infty} (a_n - d_n) = 0$ . Thus the quadruple sequence (1) has a common limit. □

*Remark 1.* 1. We have

$$\begin{aligned} (4a_{n+1})^2 - (4b_{n+1})^2 &= (a_n - b_n - c_n + d_n)^2, \\ (4a_{n+1})^2 - (4c_{n+1})^2 &= (a_n - b_n + c_n - d_n)^2, \\ (4a_{n+1})^2 - (4d_{n+1})^2 &= (a_n + b_n - c_n - d_n)^2. \end{aligned}$$

2. The quadruple sequence (1) quadratically converges, since

$$a_{n+1} - d_{n+1} \leq \frac{1}{4}(\sqrt{2a_n} - \sqrt{2d_n})^2 = \frac{1}{2} \frac{(a_n - d_n)^2}{(\sqrt{a_n} + \sqrt{d_n})^2}.$$

It is easy to see that

$$\begin{aligned} \mu(a, b, c, d) &= a\mu\left(1, \frac{b}{a}, \frac{c}{a}, \frac{d}{a}\right), \\ \mu(a, b, c, d) &= \mu\left(\frac{a+b+c+d}{4}, \frac{\sqrt{(a+d)(b+c)}}{2}, \frac{\sqrt{(a+c)(b+d)}}{2}, \frac{\sqrt{(a+b)(c+d)}}{2}\right). \end{aligned}$$

By putting  $x_1 = b/a$ ,  $x_2 = c/a$ ,  $x_3 = d/a$  for these equalities, we have the following lemma.

LEMMA 2. Let  $(y_1, y_2, y_3)$  be the image of  $(x_1, x_2, x_3)$  by the map  $\varphi$

$$\begin{aligned} \varphi(x_1, x_2, x_3) &= \left(\frac{2\sqrt{(1+x_3)(x_1+x_2)}}{1+x_1+x_2+x_3}, \frac{2\sqrt{(1+x_2)(x_1+x_3)}}{1+x_1+x_2+x_3}, \frac{2\sqrt{(1+x_1)(x_2+x_3)}}{1+x_1+x_2+x_3}\right). \end{aligned}$$

Then  $\mu$  satisfies the relation

$$(2) \quad \frac{4}{1+x_1+x_2+x_3} \mu(1, x_1, x_2, x_3) = \mu(1, y_1, y_2, y_3)$$

for  $0 < x_3 \leq x_2 \leq x_1 \leq 1$ .

**§3. Integrable Pfaffian systems**

In this section, we prepare some facts of integrable Pfaffian systems. We consider a system of first-order partial differential equations with  $r$  unknowns  $f_1, \dots, f_r$  and  $n$  variables  $x_1, \dots, x_n$  in the following form

$$(3) \quad df(x) = \Omega(x)f(x),$$

where  $x = (x_1, \dots, x_n)$  is in an open set  $U$ ,  $f(x) = {}^t(f_1(x), \dots, f_r(x))$  and  $\Omega(x)$  is an  $r \times r$  matrix whose entries are 1-forms on  $U$ . The system (3) is called a Pfaffian system on  $U$  and  $\Omega(x)$  is called the connection matrix of (3). If  $\Omega(x)$  satisfies the integrability condition

$$d\Omega(x) = \Omega(x) \wedge \Omega(x),$$

then the system (3) is integrable.

**FACT 1.** 1. *The system (3) has exactly  $r$  linearly independent vector valued solutions if and only if it is integrable.*

2. *If the system (3) is integrable, then there exists a unique solution  $f$  around  $u \in U$  such that  $f(u) = p$  for a given initial vector  $p \in \mathbb{C}^r$ .*

**FACT 2.** *For an integrable Pfaffian system (3) and an invertible  $r \times r$  functional matrix  $P(x)$  on  $U$ , the vector valued function  $g(x) = P(x)f(x)$  satisfies the Pfaffian system*

$$dg(x) = [P(x)\Omega(x)P(x)^{-1} + dP(x)P(x)^{-1}]g(x).$$

Let  $f_0$  be a function of  $n$ -variables  $(y_1, \dots, y_n)$  on an open set  $V$ . We assume that the vector valued function

$$f(y) = {}^t\left(f_0(y), \frac{\partial f_0}{\partial y_1}(y), \dots, \frac{\partial f_0}{\partial y_n}(y)\right)$$

satisfies an integrable Pfaffian system

$$df(y) = \Omega(y)f(y)$$

on  $V$ . Let  $\eta$  be a map from an open set  $U$  to  $V$  given as

$$\eta : U \ni x = (x_1, \dots, x_n) \longmapsto y = (\eta_1(x), \dots, \eta_n(x)) \in V,$$

and let  $J$  be the Jacobi matrix of  $\eta$ :

$$J = \left( \frac{\partial \eta_i}{\partial x_j} \right)_{ij} = \begin{pmatrix} \frac{\partial \eta_1}{\partial x_1} & \cdots & \frac{\partial \eta_1}{\partial x_n} \\ \vdots & \cdots & \vdots \\ \frac{\partial \eta_n}{\partial x_1} & \cdots & \frac{\partial \eta_n}{\partial x_n} \end{pmatrix}.$$

FACT 3. *If  $\det(J) \neq 0$  on  $U$  then the function  $h_0(x) = f_0(\eta(x))$  satisfies*

$$dh(x) = [\tilde{J}\Omega(x)\tilde{J}^{-1} + d\tilde{J}\tilde{J}^{-1}]h(x),$$

where

$$h(x) = {}^t \left( h_0(x), \frac{\partial h_0}{\partial x_1}(x), \dots, \frac{\partial h_0}{\partial x_n}(x) \right), \quad \tilde{J} = \begin{pmatrix} 1 & \\ & {}^t J \end{pmatrix},$$

and  $\Omega(x)$  is the pull-back of  $\Omega(y)$  under the map  $\eta$ .

**§4. Lauricella’s hypergeometric function  $F_D$**

Lauricella’s hypergeometric function  $F_D$  of  $m$ -variables  $z_1, \dots, z_m$  with parameters  $\alpha, \beta_1, \dots, \beta_m, \gamma$  is defined as

$$F_D(\alpha, \beta, \gamma; z) = \sum_{n_1, \dots, n_m \geq 0}^{\infty} \frac{(\alpha, \sum_{j=1}^m n_j) \prod_{j=1}^m (\beta_j, n_j)}{(\gamma, \sum_{j=1}^m n_j) \prod_{j=1}^m (1, n_j)} \prod_{j=1}^m z_j^{n_j},$$

where  $z = (z_1, \dots, z_m)$  satisfies  $|z_j| < 1$  ( $j = 1, \dots, m$ ),  $\beta = (\beta_1, \dots, \beta_m)$ ,  $\gamma \neq 0, -1, -2, \dots$  and  $(\alpha, n) = \alpha(\alpha + 1) \cdots (\alpha + n - 1) = \Gamma(\alpha + n)/\Gamma(\alpha)$ . This function admits the integral representation:

$$F_D(\alpha, \beta, \gamma; z) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma - \alpha)} \int_0^1 t^\alpha (1 - t)^{\gamma - \alpha} \prod_{j=1}^m (1 - z_j t)^{-\beta_j} \frac{dt}{t(1 - t)}.$$

When  $m = 1$ ,  $F_D(\alpha, \beta, \gamma; z)$  coincides with the Gauss hypergeometric function  $F(\alpha, \beta, \gamma; z)$ . We consider  $F_D(\alpha, \beta_1, \beta_2, \beta_3, \gamma; z_1, z_2, z_3)$  of three variables.

FACT 4. *The function  $F_D(\alpha, \beta, \gamma; z)$  of three variables satisfies the integrable Pfaffian system given as*

$$df = \sum_{1 \leq i < j \leq 5} A_{ij} d \log(z_i - z_j) f,$$

where  $f = {}^t(f_0, f_1, f_2, f_3)$ ,  $f_0 = F_D(\alpha, \beta, \gamma; z)$ ,  $f_i = z_i \frac{\partial f_0}{\partial z_i}$  ( $i = 1, 2, 3$ ),  $z_4 = 0$ ,  $z_5 = 1$  and

$$\begin{aligned}
 A_{12} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -\beta_2 & \beta_1 & 0 \\ 0 & \beta_2 & -\beta_1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & A_{14} &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 + \beta_2 + \beta_3 - \gamma & 0 & 0 \\ 0 & -\beta_2 & 0 & 0 \\ 0 & -\beta_3 & 0 & 0 \end{pmatrix}, \\
 A_{13} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -\beta_3 & 0 & \beta_1 \\ 0 & 0 & 0 & 0 \\ 0 & \beta_3 & 0 & -\beta_1 \end{pmatrix}, & A_{24} &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & -\beta_1 & 0 \\ 0 & 0 & 1 + \beta_1 + \beta_3 - \gamma & 0 \\ 0 & 0 & -\beta_3 & 0 \end{pmatrix}, \\
 A_{23} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -\beta_3 & \beta_2 \\ 0 & 0 & \beta_3 & -\beta_2 \end{pmatrix}, & A_{34} &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -\beta_1 \\ 0 & 0 & 0 & -\beta_2 \\ 0 & 0 & 0 & 1 + \beta_1 + \beta_2 - \gamma \end{pmatrix}, \\
 A_{15} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ -\alpha\beta_1 & \gamma - \alpha - \beta_1 - 1 & -\beta_1 & -\beta_1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 A_{25} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\alpha\beta_2 & -\beta_2 & \gamma - \alpha - \beta_2 - 1 & -\beta_2 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 A_{35} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\alpha\beta_3 & -\beta_3 & -\beta_3 & \gamma - \alpha - \beta_3 - 1 \end{pmatrix}.
 \end{aligned}$$

*Remark 2.* The  $A_{ij}$  and  $A_{i,n+1}$  in the proof of Proposition 9.1.4 in [IKSY] are wrong. Professor K. Ohara informed us of the correct Pfaffian system given by the system [O].

**§5. Main theorem**

**THEOREM 1.** *For any numbers  $x_1, x_2, x_3$  satisfying  $0 < x_3 \leq x_2 \leq x_1 \leq 1$ , we have*

$$\frac{1}{\mu(1, x_1, x_2, x_3)} = F_D\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, 1; 1 - x_1^2, 1 - x_2^2, 1 - x_3^2\right)^2,$$

where  $\mu(1, x_1, x_2, x_3)$  is the common limit of the quadruple sequence (1) with initial  $(1, x_1, x_2, x_3)$  and  $F_D$  is Lauricella's hypergeometric function.

We put

$$F(z_1, z_2, z_3) = F_D\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, 1; z_1, z_2, z_3\right).$$

PROPOSITION 1. *The function  $F$  satisfies*

$$\begin{aligned} & \frac{1+x_1+x_2+x_3}{4} F(1-x_1^2, 1-x_2^2, 1-x_3^2)^2 = F(1-y_1^2, 1-y_2^2, 1-y_3^2)^2 \\ & = F\left(\left(\frac{1-x_1-x_2+x_3}{1+x_1+x_2+x_3}\right)^2, \left(\frac{1-x_1+x_2-x_3}{1+x_1+x_2+x_3}\right)^2, \left(\frac{1+x_1-x_2-x_3}{1+x_1+x_2+x_3}\right)^2\right)^2, \end{aligned}$$

where  $(y_1, y_2, y_3) = \varphi(x_1, x_2, x_3)$  is defined in Lemma 2.

*Proof.* Put

$$(\xi_1, \xi_2, \xi_3) = \left(\frac{1-x_1-x_2+x_3}{1+x_1+x_2+x_3}, \frac{1-x_1+x_2-x_3}{1+x_1+x_2+x_3}, \frac{1+x_1-x_2-x_3}{1+x_1+x_2+x_3}\right).$$

Then we have

$$\begin{aligned} (x_1, x_2, x_3) &= \left(\frac{1-\xi_1-\xi_2+\xi_3}{1+\xi_1+\xi_2+\xi_3}, \frac{1-\xi_1+\xi_2-\xi_3}{1+\xi_1+\xi_2+\xi_3}, \frac{1+\xi_1-\xi_2-\xi_3}{1+\xi_1+\xi_2+\xi_3}\right), \\ \frac{1+x_1+x_2+x_3}{4} &= \frac{1}{1+\xi_1+\xi_2+\xi_3}, \\ (1-x_1^2, 1-x_2^2, 1-x_3^2) &= \left(\frac{4(1+\xi_3)(\xi_1+\xi_2)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_2)(\xi_1+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_1)(\xi_2+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}\right). \end{aligned}$$

Thus the equality in Proposition 1 is equivalent to

$$\begin{aligned} & \sqrt{1+\xi_1+\xi_2+\xi_3} F(\xi_1^2, \xi_2^2, \xi_3^2) \\ &= F\left(\frac{4(1+\xi_3)(\xi_1+\xi_2)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_2)(\xi_1+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_1)(\xi_2+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}\right) \end{aligned}$$

for  $0 \leq \xi_1 \leq \xi_2 \leq \xi_3 < 1$ . We show that the Pfaffian systems obtained by the functions in the both sides of the above equality coincide.

Let  $\Omega(x)$  be the connection 1-form in Fact 4 for  $\alpha = \beta_1 = \beta_2 = \beta_3 = 1/4$  and  $\gamma = 1$ . Fact 2 implies that the vector valued function

$$g(x) = {}^t\left(F, \frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial x_2}, \frac{\partial F}{\partial x_3}\right)$$

satisfies the Pfaffian system  $dg = \Omega_1(x)g$ , where

$$\Omega_1(x) = P\Omega(x)P^{-1} + dPP^{-1},$$

$$P = \text{diag}\left(1, \frac{1}{x_1}, \frac{1}{x_2}, \frac{1}{x_3}\right) = \begin{pmatrix} 1 & & & \\ & \frac{1}{x_1} & & \\ & & \frac{1}{x_2} & \\ & & & \frac{1}{x_3} \end{pmatrix}.$$

The vector valued function

$$h(\xi) = {}^t\left(h_0, \frac{\partial h_0}{\partial \xi_1}, \frac{\partial h_0}{\partial \xi_2}, \frac{\partial h_0}{\partial \xi_3}\right)$$

for

$$h_0(\xi_1, \xi_2, \xi_3) = \sqrt{1 + \xi_1 + \xi_2 + \xi_3}F(\xi_1^2, \xi_2^2, \xi_3^2)$$

satisfies  $h(0, 0, 0) = {}^t(1, 1/2, 1/2, 1/2)$  and the Pfaffian system  $dh = \Omega_2(\xi)h$ , where

$$\Omega_2(\xi) = Q[J_1\Omega_1(\xi)J_1^{-1} + dJ_1J_1^{-1}]Q^{-1} + dQQ^{-1}, \quad J_1 = \text{diag}(1, 2\xi_1, 2\xi_2, 2\xi_3),$$

$$Q = \begin{pmatrix} \zeta & & & \\ \frac{1}{2\zeta} & \zeta & & \\ \frac{1}{2\zeta} & & \zeta & \\ \frac{1}{2\zeta} & & & \zeta \end{pmatrix}, \quad \zeta = \sqrt{1 + \xi_1 + \xi_2 + \xi_3},$$

and  $\Omega_1(\xi)$  is the pull-back of  $\Omega_1(x)$  under the map

$$(\xi_1, \xi_2, \xi_3) \mapsto (x_1, x_2, x_3) = (\xi_1^2, \xi_2^2, \xi_3^2).$$

On the other hand, the vector valued function

$$h(x) = {}^t\left(h_0, \frac{\partial h_0}{\partial \xi_1}, \frac{\partial h_0}{\partial \xi_2}, \frac{\partial h_0}{\partial \xi_3}\right)$$

for

$$h_0(\xi_1, \xi_2, \xi_3) = F\left(\frac{4(1+\xi_3)(\xi_1+\xi_2)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_2)(\xi_1+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_1)(\xi_2+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}\right)$$

satisfies  $h(0, 0, 0) = {}^t(1, 1/2, 1/2, 1/2)$  and the Pfaffian system  $dh = \Omega_3(\xi)h$ , where

$$\Omega_3(\xi) = J_2\Omega'_1(\xi)J_2^{-1} + dJ_2J_2^{-1}, \quad J_2 = \begin{pmatrix} 1 & \\ & {}^tJ \end{pmatrix},$$

$\Omega'_1(\xi)$  is the pull-back of  $\Omega_1(x)$  under the map

$$\varphi' : (\xi_1, \xi_2, \xi_3) \mapsto \left( \frac{4(1+\xi_3)(\xi_1+\xi_2)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_2)(\xi_1+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2}, \frac{4(1+\xi_1)(\xi_2+\xi_3)}{(1+\xi_1+\xi_2+\xi_3)^2} \right),$$

and  $J$  is the Jacobi matrix of the map  $\varphi'$ . By a straight forward calculation, we can show that  $\Omega_2(\xi) = \Omega_3(\xi)$ . Thus we have the required equality around  $\xi = (0, 0, 0)$ . □

*Proof of Theorem 1.* Consider the quadruple sequence (1) with initial  $(a_0, b_0, c_0, d_0) = (1, x_1, x_2, x_3)$ . Lemma 2 and Proposition 1 imply that

$$\begin{aligned} & \mu(1, x_1, x_2, x_3)F(1 - x_1^2, 1 - x_2^2, 1 - x_3^2)^2 \\ &= \mu(1, y_1, y_2, y_3)F(1 - y_1^2, 1 - y_2^2, 1 - y_3^2)^2. \end{aligned}$$

Thus we have

$$\begin{aligned} & \mu(1, x_1, x_2, x_3)F(1 - x_1^2, 1 - x_2^2, 1 - x_3^2)^2 \\ &= \mu\left(1, \frac{b_n}{a_n}, \frac{c_n}{a_n}, \frac{d_n}{a_n}\right)F\left(1 - \left(\frac{b_n}{a_n}\right)^2, 1 - \left(\frac{c_n}{a_n}\right)^2, 1 - \left(\frac{d_n}{a_n}\right)^2\right)^2 \end{aligned}$$

for any  $n \in \mathbb{N}$ . Since

$$\lim_{n \rightarrow \infty} \frac{b_n}{a_n} = \lim_{n \rightarrow \infty} \frac{c_n}{a_n} = \lim_{n \rightarrow \infty} \frac{d_n}{a_n} = 1,$$

and  $\mu(1, 1, 1, 1) = F(0, 0, 0) = 1$ , we have

$$\begin{aligned} & \mu(1, x_1, x_2, x_3)F(1 - x_1^2, 1 - x_2^2, 1 - x_3^2)^2 \\ &= \lim_{n \rightarrow \infty} \mu\left(1, \frac{b_n}{a_n}, \frac{c_n}{a_n}, \frac{d_n}{a_n}\right)F\left(1 - \left(\frac{b_n}{a_n}\right)^2, 1 - \left(\frac{c_n}{a_n}\right)^2, 1 - \left(\frac{d_n}{a_n}\right)^2\right)^2 \\ &= \mu(1, 1, 1, 1)F(0, 0, 0)^2 = 1, \end{aligned}$$

which is the desired equality. □

**COROLLARY 1.** For  $1 > x_1 \geq x_2 \geq x_3 \geq 0$ , we have

$$F(1 - x_1^2, 1 - x_2^2, 1 - x_3^2) = \prod_{n=0}^{\infty} \sqrt{\frac{a_n}{a_{n+1}}},$$

where we set the initial of the quadruple sequence (1) as  $(a_0, b_0, c_0, d_0) = (1, x_1, x_2, x_3)$ .

*Proof.* By Proposition 1, we have

$$\begin{aligned} & F(1 - x_1^2, 1 - x_2^2, 1 - x_3^2) \\ &= \frac{2}{\sqrt{1+x_1+x_2+x_3}} F(1 - y_1^2, 1 - y_2^2, 1 - y_3^2) \\ &= \sqrt{\frac{4a_0}{a_0+b_0+c_0+d_0}} F\left(1 - \frac{b_1^2}{a_1^2}, 1 - \frac{c_1^2}{a_1^2}, 1 - \frac{d_1^2}{a_1^2}\right) \\ &= \sqrt{\frac{a_0}{a_1}} \sqrt{\frac{a_1}{a_2}} F\left(1 - \frac{b_2^2}{a_2^2}, 1 - \frac{c_2^2}{a_2^2}, 1 - \frac{d_2^2}{a_2^2}\right) \\ &= \left(\prod_{i=0}^{n-1} \sqrt{\frac{a_i}{a_{i+1}}}\right) F\left(1 - \frac{b_n^2}{a_n^2}, 1 - \frac{c_n^2}{a_n^2}, 1 - \frac{d_n^2}{a_n^2}\right), \end{aligned}$$

which implies this corollary. □

### §6. A specialization

For the case  $b = c = d$ , the quadruple sequence reduces to

$$a_{n+1} = \frac{a_n + 3b_n}{4}, \quad b_{n+1} = c_{n+1} = d_{n+1} = \sqrt{b_n \frac{a_n + b_n}{2}},$$

which is studied in [BB]. It is shown that the reciprocal of the common limit of the double sequences is  $F\left(\frac{1}{4}, \frac{3}{4}, 1; 1 - x^2\right)^2$ , where  $x = b/a$  and  $F(\alpha, \beta, \gamma; z)$  is the Gauss hypergeometric function. By our main theorem, we have

$$\begin{aligned} & F_D\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, 1; 1 - x^2, 1 - x^2, 1 - x^2\right)^2 \\ &= \frac{1}{\mu(1, x, x, x)} = \frac{1}{M_4(1, x)} = F\left(\frac{1}{4}, \frac{3}{4}, 1; 1 - x^2\right)^2. \end{aligned}$$

Note that the above reduction of  $F_D$  to  $F$  can be easily obtained by the integral representation of  $F_D$ .

### REFERENCES

- [B] C. W. Borchardt, *Über das arithmetisch-geometrische Mittel aus vier Elementen*, Berl. Monatsber, **53** (1876), 611–621.
- [BB] J. M. Borwein and P. B. Borwein, *A cubic counterpart of Jacobi’s identity and the AGM*, Trans. Amer. Math. Soc., **323** (2) (1991), 691–701.
- [IKSY] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida, *From Gauss to Painlevé*, Vieweg, Braunschweig, Wiesbaden, 1991.

- [KS1] K. Koike and H. Shiga, *Isogeny formulas for the Picard modular form and a three terms arithmetic geometric mean*, J. Number Theory, **124** (2007), 123–141.
- [KS2] K. Koike and H. Shiga, *Extended Gauss AGM and corresponding Picard modular forms*, J. Number Theory, **128** (2008), 2097–2126.
- [M] J. F. Mestre, *Moyenne de Borchardt et intégrales elliptiques*, C. R. Acad. Sci. Paris, **313** (1991), 273–276.
- [MM] D. V. Manna and V. H. Moll, *Landen survey*, *Probability, Geometry and Integrable Systems*, MSRI Publications, **55**, Cambridge University Press, Cambridge, 2008, pp. 287–319.
- [O] K. Ohara, *yang — a package for computation in the ring of differential-difference operators*, <http://www.openxm.org>, 2007.

Takayuki Kato

*MEC INC.*

*2-6 Futuka, Aoba-ku*

*Sendai 980-0802*

*Japan*

`katou-t@mec-inc.co.jp`

Keiji Matsumoto

*Department of Mathematics*

*Hokkaido University*

*Sapporo 060-0810*

*Japan*

`matsu@math.sci.hokudai.ac.jp`