Vol. XXXVII., No. 1

The Journal of Laryngology and Otology

EDITED BY

A. LOGAN TURNER AND J. S. FRASER

Contents

HEMORRHAGE DURING AND AFTER TONSILLECTOMY: SURGICAL PRINCIPLES AND METHODS FOR ITS CONTROL CORNELIUS G. COAKLEY

THE PATHOLOGICAL AND CLINICAL ASPECTS OF DEAF-MUTISM J. S. FRASER

CLINICAL RECORDS :--

A CASE OF FATAL MENINGITIS FOLLOWING SUB-

MUCOUS RESECTION OF THE NASAL SEPTUM. W. ELIAS POWELL FOREIGN BODY IN RIGHT BRONCHUS . T. ARTHUR MACGIBBON FACIAL PARALYSIS ASSOCIATED WITH ACUTE' MIDDLE-EAR SUPPURATION . . . ERNEST CULPIN

Societies' Proceedings :---

ROYAL SOCIETY OF MEDICINE-SECTION OF OTOLOGY.

ABSTRACTS :---

EAR.

NOSE AND ACCESSORY SINUSES.

PHARYNX.

REVIEWS OF BOOKS. GENERAL NOTES. LETTER TO EDITORS. EDITORIAL

Edinburgh: Oliver and Boyd London: 33 Paternoster Row

Monthly, 4/- net

Annual Subscription, 40/- net

"We have experimentally proved that Listerine is a powerful antiseptic, preventing the development of bacteria and decomposition of vegetable infusions."—British Medical fournal, May 3, 1890.

LISTERINE

A safe, non-poisonous, unirritating antiseptic solution

LISTERINE

is particularly useful in the treatment of abnormal conditions of the mucosa, and admirably suited for a wash, gargle, or douche, in catarrhal conditions of the nose and throat.

LISTERINE

in proper dilution may be freely and continuously used without prejudicial effect, either by injection or spray, in all the natural cavities of the body.

LISTERINE embodies a two-fold antiseptic effect, in that after the evaporation of its volatile constituents—thyme, eucalyptus, mentha, gaultheria, and ethyl alcohol—a film of boracic and benzoic acids remains upon the surface to which Listerine has been applied, affording more prolonged antiseptic protection.

LISTERINE is a trustworthy surgical dressing; it has no injurious effect upon the tissues in which the healing process is going on.

LISTERINE in teaspoonful doses will often afford relief in fermentative dyspepsia, and is largely prescribed, with excellent results, in the various forms of diarrhœa, occurring in infants and adults.

LISTERINE literature, including special pamphlets upon Respiratory Diseases, may be had, by physicians, upon application to

LAMBERT PHARMACAL COMPANY

2101 Locust Street

St Louis, Mo., U.S.A.

British Agents:

S. Maw, Son & Sons, 7 Aldersgate St., London, E.C.

The Journal of Laryngology and Otology

(Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

A. LOGAN TURNER AND J. S. FRASER

ASSISTED BY

DOUGLAS GUTHRIE AND IRWIN MOORE

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked and no additional matter should be added.

4. The order for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied, on which the price of the reprints is stated.

5. Authors of Original Communications on Oto-Laryngology in other Journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

6. Editorial Communications should be addressed to "EDITORS, *Journal of Laryngology*, c/o Messrs OLIVER AND BOYD, Tweeddale Court, Edinburgh."

7. The Annual Subscription is Forty Shillings or Ten Dollars, and is payable in advance.

8. Single copies of back numbers, both of the present and the previous series, are on sale at Four Shillings (One Dollar) each.

9. An Annual Subscription can be commenced with any number of the Journal, and the preceding numbers of the current year can be purchased at Four Shillings (One Dollar) each.

PHENATE OF SODA SOLUTION

Sol. Sodæ Phenatis (Woolley)

Invaluable as a Mouth and Throat Wash and for allaying tenderness of the Gums after Teeth Extraction ANTISEPTIC and ASTRINGENT

Price, Retail, of all Chemists-

¹/₄-Pint Bottles, 1/6; ¹/₂-Pint, 2/9; Pint, 4/6; Quart, 7/-; ¹/₂-Gallon, 12/6 Special Prices to the Medical and Dental Professions


A DENTIST writes :—"I have found it invaluable in dental practice for syringing the sockets after removal of teeth which have been subject to an acute attack of *periostitis*; the strength used is one part of the solution to 15 parts of hot water, the relief from intense pain being instantaneous."

JAMES WOOLLEY, SONS & CO., LTD.

Manufacturing Chemists

MANCHESTER

<section-header><section-header><section-header><section-header><section-header><section-header><text><text><text><text><text><text>



HOSPITAL FOR DI	SEASES OF QUARE, LONDON,	
HONORARY	MEDICAL	STAFF
Subgeons. Mr Jefferson Faulder Dr W. H. Kelson	OUT-PATIENTS. Wednesday, 2 P.M. Friday, 2 P.M.	Operations. Thursday, 10 A.M. Friday, 9.30 A.M.
Mr George W. Badgerow	{ Monday, 2 P.M. Tuesday, 6 P.M.	} Wednesday, 9 A.M.
Mr G. W. DAWSON Mr Gilbert Chubb	Friday, 6.30 P.M. Tuesday, 2 P.M. Thursday, 2 P.M. Saturday, 10 A.M.	Tuesday, 2 P.M. Thursday, 2 P.M. Tuesday, 10 A.M. Friday, 2 P.M.
	 Tuesday, 6 P.M. Saturday, 10 A.M. Tuesday, 2 P.M. Thursday, 2 P.M. 	} Monday, 10 A.M. } Wednesday, 10 A.M.

CLINICAL INSTRUCTION

Clinical Instruction in the Diagnosis and Treatment of Disease is given daily in the Out-Patient Department from 2 to 5 p.m.; on Taesdays and Fridays from 6.30 to 9 p.m. Major operations are per-formed at the times specified above, and minor operations by the Assistant Surgeons. The Hospital contains 60 beds for In-Patients. There is also an Out-Patient attendance of over 60,000 yearly. Practitioners and Medical Students are admitted to the Practice of the Hospital and from among them

appointed. For further information apply to the Dean-LIONEL COLLEDGE.



iv



CONTENTS

Editorial	I
HEMORRHAGE DURING AND AFTER TONSILLECTOMY: SURGICAL PRINCIPLES AND METHODS FOR ITS CONTROL. Cornelius G. Coakley, M.D., New York	
	3
THE PATHOLOGICAL AND CLINICAL ASPECTS OF DEAF-MUTISM. J. S. Fraser, M.B., F.R.C.S.Ed.	13
CLINICAL RECORDS : A Case of Fatal Meningitis following Submucous Resection of the Nasal	
Septum. W. Elias Powell, M.B.	39
Foreign Body in the Right Bronchus. T. Arthur MacGibbon, F.R.C.S.E., Christchurch, N.Z.	40
Facial Paralysis associated with Acute Middle-Ear Suppuration. Ernest Culpin, Brisbane	41
SOCIETIES' PROCEEDINGS :	42
Abstracts :	
Ear	45
Nose and Accessory Sinuses	47
Pharynx	49
REVIEWS OF BOOKS.	
	52
Letter to Editors	53
GENERAL NOTES	54

HEMOPLASTIN

(HÆMOSTATIC SERUM)

Remarkably efficacious in all forms of hæmorrhage

HEMOPLASTIN is a sterile fluid derived from normal serum of horses and cattle. It is composed principally of prothrombin, anti-antithrombin, and thrombokinase, in physiologically balanced solution ready for immediate use. It keeps well.

HEMOPLASTIN is indicated in the treatment of all types of hæmorrhage, particularly those depending upon imperfect coagulation of the blood. It has proved of great value in pulmonary hæmorrhage, purpura hæmorrhagica, and in operative hæmorrhage such as occurs in tonsillectomy, nasal and intracranial surgery. Its administration to hæmophiliac subjects about to undergo surgical operations is a valuable precaution against excessive post-operative hæmorrhage.

HEMOPLASTIN may be administered intravenously or subcutaneously. The dose, both for children and adults, is I to 2 c.c. every 4 to 6 hours until the hæmorrhage is controlled. Hemoplastin is also effective when applied on sterile gauze to bleeding surfaces.

Hemoplastin (P., D. & Co.) is supplied in bulbs containing 2 c.c.

PARKE, DAVIS & CO., ^{50 Beak St.,} LONDON, W.1