VOL. VIII.—1894.

THE

JOURNAL OF LARYNGOLOGY,

RHINOLOGY AND OTOLOGY;

AN ANALYTICAL RECORD OF CURRENT LITERATURE

RELATING TO

THE THROAT, NOSE AND EAR.

PUBLISHED MONTHLY.

Tondon:

F. J. REBMAN, 11, ADAM STREET, STRAND.

AMERICAN AGENTS:

THE F. A. DAVIS COMPANY, 1914-16, CHERRY STREET, PHILADELPHIA, PA.

ENTERED AT STATIONERS HALL

THE JOURNAL OF LARYNGOLOGY, RHINOLOGY, AND OTOLOGY.

Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN.

EDITED BY

R. NORRIS WOLFENDEN, M.D. Cantab.

JOHN MACINTYRE, M.B. Glasgow.

DUNDAS GRANT, M.A., M.D., F.R.C.S. Eng.

WITH THE ASSISTANCE OF

DR. SANDFORD (Cork), DR. MILLIGAN (Manchester),

AND WITH THE CO-OPERATION OF

Drs. Baron (Bristol), R. Botey (Barcelona),
Cartaz (Paris), Bryson Delavan (New York),
Draispul (St. Petersburg), Grazzi (Florence), Guye (Amsterdam),
Hicguet (Brussels), Middlemass Hunt (Liverpool),
Valerius Idelson (Berne), Joal (Paris), A. B. Kelly (Glasgow),
Karwowski (Warsaw), Lacoarret (Toulouse), Lake (London),
Lichtwitz (Bordeaux), John N. Mackenzie (Baltimore),
G. W. Major (Montreal), Prof. Massei (Naples),
Meyjes (Amsterdam), E. Meyer (Berlin), Michael (Hamburg),
Holger Mygind (Copenhagen), W. Porter (St. Louis),
Robertson (Newcastle), Sajous (Philadelphia), Sendziak (Warsaw),
Sota (Seville), and St. Clair Thomson (London).

CONTENTS.

Original Article:-

Nose and Naso-Pharynx, &c.—Microscopical Anatomy of Human Nasal Cavities, Suchannek, Hermann—Rare Case of Polypus of Septum, Lange, V.—Nasal Polypus, V.—William of Nasal Mucosa, Reth.—Epithelioma of Nasal Cavities, Preyfuss of Nasal Mucosa, Reth.—Polythelioma of Nasal Cavities, Preyfus of Nasal Indiana (Printed Markett, H. S.) Sis, Chiari—Nasal Tuberculosis Herzog—Empyema of Antrum of Highmore, Birkett, H. S.—Malignant Transformation of Antral Tunours, Finh.—Adenoids in Ear Disease, Leiderman, M. D.—Adenoids, Fraenkel — Enuresis Nocturna and Nasal Obstruction, Otto.

Pages 70 to 74.

Larynx.—Elongation of Epiglottis, Brown, Price—Falsetto Voice, Bennett, O. P.—
Hysterical Dumbness, Kayser—Spontaneous Cure of Laryngeal Tuberculosis, Grayson—
Crico-Arytenoid Anchylosis, Bennett, O. P.—Bony Occlusion of Posterior Nares, Bennett,
O. P.—Rheumatism of Crico-Arytenoid, Newcomb, J. E.—Lipomata of Upper Air-Pa-sages,
Künne—Foreign Body in Larynx, Hopkins, F. E.—Instruments for Operations through
Fenestrated Intubation Tubes, Lichtwitz—Subcordal Sarcoma, Birkett—Extirpation of
Half the Larynx, Gluck—Laryngo-Fissure, Pinaczek.

Pages 74 to 78.

Thyroid Gland, Neck, &c.—Goitre. Dupuis, T. R.—Graves' Disease, Troje— Electrolysis in Goitre, Dickson, C. R.—Some Diseases of Thyroid, Greenfield—Myxcedema, Moster—Treatment of Myxcedema, Wiehmann—Thyroid Treatment in Health and Myxcedema, Vermehren—Abscesses of Neck, Poulsen—Enlarged Cervical Glands, Allbutt, T. C. Pages 78 to 80.

Ears.—Anthropology of the Ear in Criminals, Daac, II.—Removal of Stapes, Becold, Blake, C. J.—Removal of Ossicles, Sheppard.—Schwartze's Modification of Stacke's Operation, Holmes, C. R.—Bezold's Mastoiditis, Vulpius, II.—Septic Thrombosis of Lateral Sinus, Vickery.—Bilateral Fracture of Temporal Bone, Vulpius, IV.—Deafness in Cerebellar Disease, Dercum, F. K.—Reciprocal Interaction of the Ears, Urbantschitsch.

Pages 81 to 81.

Association Meetings:-

The British Laryngological and Rhinological Association. Laryngological Society of Berlin.

Page 87. Page 110.

PAGE

Reviews:—Late Forms of Hereditary Syphilis in the Upper Air-Passages, Gerber—Normal and Pathological Anatomy of the Nasal Cavity, and its Accessory Cavities, Zuckerkandl—Text-Book of Diseases of the Ear, Gruber (Law and Jewell)—Manuel pratique des Maladies des Fosses Nasales, etc., Moure, E. J.

Pages 111 to 114.

GUIDE TO ADVERTISEMENTS.

Allen & Hanburys vii	., ix.
Bullock & Co.	XIV.
Burroughs, Wellcome & Co iv., vi.,	XIII.
Central London Throat Hospital	xvı.
Godfrey & Cooke	viii.
Houdé	XV.
Ingram & Royle	х.
Lambert Pharmaca Co	11.
Parks Davis & Co	VIII.
Rebman viii., xi., xii.	,
Watt & Son	xiv.

KEPLER PREPARATIONS.

THE KEPLER EXTRACT OF MALT.

The many and varied conditions of disease in which the Kepler Extract of Mala as a food is indicated, and the benefits which are to be obtained from its use sufficiently explain the wide popularity it has attained. It is manufactured from barley and barley only, and our perfected machinery and long experience have enabled us to obtain a product containing all the nutritive principles of the grain intact. The mair advantages of a good Malt Extract are that it is highly nutritious, readily assimilated and rich in that important factor diastase, which converts starchy food into maltose and other saccharine and easily digested substances. It is consequently an effective aid to digestion, and can be tolerated by the stomach when cod liver oil alone would be out of the question. Freedom from coarse and irritating elements makes it the safes aliment in inflammatory conditions of disease; it is an admirable demulcent in irritable states of the throat, and in debilitated conditions generally it is unrivalled.

Supplied to the Medical Profession in alb. and 18lb. bottles at 1s. 8d. and 3s. each.

THE KEPLER ESSENCE OF MALT.

In dietetics for the invalid, the Kepler Essence of Malt has already won its way to an enviable position, and has distinctly shown its superiority over all so-called nourishing stouts and ales as a food beverage. Its superior diastasic property, it palatable character, and its freedom from intoxicating qualities raise it at once fa above the ordinary level of the very best and most carefully-prepared fermented mal liquors. We are assured by competent judges who have tried it in their clinics that two fluid ounces of the Kepler Malt Essence are equal in therapeutic value to a pint the best stout or ale. This is a most important fact, to be kept well in mind wherecommending or prescribing suitable diet for the dyspeptic, the ill-nourished, the sick or the convalescent. The Kepler Extract of Malt makes a pleasant beverage whe mixed with aerated water and milk.

Supplied to the Medical Profession in champagne pints and quarts at 1s. 2d. and 2s. per bottle.

THE KEPLER SOLUTION OF COD LIVER OIL IN EXTRACT OF MALT.

The Kepler Solution was the first preparation which represented a rational metho of administering cod liver oil. When an oil is given alone, a considerable an prolonged digestive effort is required before assimilation is possible, and in not a fe instances it is voided unchanged. Moreover, the digestion is generally soon derange. The two problems which remained unsolved for so many years were: (1) How to cove the unpleasant taste of the oil; (2) how to increase its assimilability. Pharmacis tried every possible means to attain these ends, but since they lost sight of the fathat cod liver oil is a food, and treated it as a drug, they met with only partial success In emulsions the taste was somewhat masked, but it was found that in the stomach toil globules coalesced, and unpleasant and nauseating eructations resulted. In the Kepler Solution the oil is actually dissolved in Extract of Malt. It is not emulsified, ar no oil globules are visible under the microscope. No unpleasant effects are produce when it is taken into the stomach, but it passes through the pylorus unaffected, ar is readily absorbed by the lacteals.

Supplied to the Medical Profession in $\frac{3}{4}$ lb. and $1\frac{1}{2}$ lb. bottles at 1s. 8d. and 3s. each.

BURROUGHS, WELLCOME & CO., Manufacturing Chemist SNOW HILL BUILDINGS, LONDON.

Cable and Telegraphic Address: "BURCOME, LONDON."