Index of Subjects

Airborne non-sporeforming anaerobic bacteria, 181

Airborne operating room bacteria, evaluation of with a Biap slit sampler, 37

Age-sex distribution of Toxoplasma antibody in South Australia, 315

Anaerobic non-sporeforming airborne bacteria, 181

Anaerobic bacilli, gram-negative, isolated from clinical infections, 301

Antibiotic-containing diets: their effect on colonization of the alimentary tract of poultry by salmonella organisms, 137

Antibody responses and virus shedding following trivalent oral poliomyelitis vaccines prepared in monkey kidney or in human diploid cell substrates, 17

Anticoagulant rodenticides, the susceptibility of Bandicota bengalensis from Rangoon to, 127

Bacillus cereus spores, heat resistance of, and growth of the organism in boiled rice, 77 Bacteriological survey of sanitary dressings, and development of an effective means for their disposal, 41

Bandicota bengalensis from Rangoon; its susceptibility to several anticoagulant rodenticides, 127

Biap slit sampler for evaluation of airborne operating room bacteria, 37

Boiled compared with formalized leptospiral vaccines in rabbits, hamsters and humans, 1

Bovine udder experimentally infected with Campylobacter colificiuni, 421

Bromadialone as a rodenticide for use against warfarin-resistant and non-resistant rats and mice, 263

Brucella culture supernatants, an RNA fraction present in, 223

Brucellosis, bovine, the serodiagnosis of by means of non-specific immunological factors, 405

Campylobacter colificiuni, experimental infection of the bovine udder with, 421

Campylobacter infection, a milk-borne outbreak, 415

Chicken factory, an outbreak of streptococcal infection in, 71

Cleaning effect of flushing disinfectors, measurement of, 321

Clostridium perfringens, methods for counting in poultry meat and other products, 151

Clostridium perfringens food poisoning, serological typing in outbreaks of, 443

Coliphage f2, inactivation of in municipal effluent by various disinfectants, 247

Conglutinin and other non-specific immunological factors in the serodiagnosis of bovine brucellosis, 405

Contaminated medicaments in use in a hospital for diseases of the skin, 103

Down's syndrome subjects in institutions, their response to enterovirus infections, 433

Echinococcus granulosus, Taenia hydatigena and T. ovis, simulating strategies for control of, 389

Encephalitozoon cuniculi in the wild rabbit in Victoria, Australia, 295

Enterotoxigenicity of environmental Escherichia coli, 411

Enterovirus infections, the response of institutionalized Down's syndrome subjects to, 433

Escherichia coli in the environment, belonging to serotypes normally considered enterotoxigenic, 411

Escherichia coli in market pigs, 1956–1979: emergence of organisms with plasmid-borne trimethoprim resistance, 467

Faeces of babies born in hospital and at home, gram-negative aerobic bacilli in, 91

Fleas, European rabbit; the emergence of males and females from laboratory cultures, 109

Flotation procedures for recovering ova of Toxocara spp. from soil, 83

Flushing disinfectors, a method for measuring the cleaning effect of, 321

Food poisoning by Clostridium perfringens; application of serological typing to the investigation of outbreaks, 443

Foot-and-mouth disease virus, heat inactivation of in milk from intramammarily infected susceptible cows, 159

Gram-negative aerobic bacilli in the faeces of babies born in hospital compared with those born at home, 91

Gram-negative anaerobic bacilli isolated from clinical infections, identification of, 301 Growth medium composition, the effect of on the antigenicity of *Mycoplasma bovis*, 29

Heat inactivation of foot-and-mouth disease virus in milk from intramammarily infected susceptible cows, 159

Heat resistance of *Bacillus cereus* spores, and growth of the organism in boiled rice, 77 Hepatitis B serology in Greek prostitutes: significance of the different serum markers, 257 Hospital food and kitchens, *Klebsiella* species in, 97

Immunogenicity of boiled compared with formalized leptospiral vaccines in rabbits, hamsters and humans, 1

Infection by viruses in small British rodents, 285

Infection prevention in patients with cancer: microbiological evaluation of laminar air flow isolation, topical chlorhexidine, and oral non-absorbable antibiotics, 457

Influenza vaccine, polyvalent; protection by, and persistence of homologous and heterologous HI antibodies during two epidemic seasons, 237

Intradermal versus subcutaneous immunization with typhoid vaccine, 11

Iodine, the effects of on biological activities of myxoviruses, 381

Isolation and classification of sixteen strains of saprophytic leptospires, 173

Klebsiella species in hospital food and kitchens: a source of organisms in the bowel of patients,

Laminar air flow isolation, topical chlorhexidine and oral non-absorbable antibiotics compared for infection prevention in patients with cancer, 457

Leptospiral vaccines, immunogenicity of boiled compared with formalized in rabbits, hamsters and humans, 1

Leptospires, isolation and classification of sixteen saprophytic strains of, 173

Macba plate, technique for cultural examination of urine using a single plate, 429

Media, and methods for counting Clostridium perfringens in poultry meat and furtherprocessed products, 151

Milk-borne outbreak of Campylobacter infection, 415

Milk products, commercial, and indigenous weaning foods in a rural West African environment: a bacteriological perspective, 191

Municipal effluent, inactivation of f2 coliphage in by various disinfectants, 247

Municipal effluent, inactivation of viruses in by chlorine, 63

Mycoplasma bovis, the effect of growth medium composition on the antigenicity of, 29

Myxomatosis; the emergence of male and female European rabbit fleas $\bar{S}pilopsyllus$ cuniculi (Dale) from laboratory cultures, 109

Myxoviruses, the effects of iodine on the biological activities of, 381

Neomycin and polymyxin B, the doses which can be applied for selective decontamination of the digestive tract in mice, 331

Plesiomonas shigelloides from aquatic environments, domestic animals, pets and humans, 203 Poliomyelitis trivalent oral vaccines, prepared in either monkey kidney or human diploid cell substrates; comparison of antibody responses and virus shedding following administration of, 17

Prostitutes, Greek; hepatitis B serology in, 257

Pyriminyl as a poison for Rattus norvegicus, 347

Radial haemolysis for measurement of rubella antibody: a simple screening method, 213

Rattus norvegicus, laboratory and field evaluation of pyriminyl as a poison for, 347 RNA fraction present in Brucella culture supernatants; isolation and properties of, 223

Rodenticides, acute; statistical analysis of comparative field trials of, 341

R plasmids derived from Salmonella and Escherichia coli obtained from a pig farm, genetical relationship between, 365

Rubella antibody measured by radial haemolysis. A simple screening method for diagnostic laboratories, 213

Salmonella and Escherichia coli obtained from a pig farm; genetical relationship between R plasmids derived from, 365

Salmonella organisms in the alimentary tract of poultry; the effect of antibiotic-containing diets on, 137

Salmonella typhimurium, phage type/biotype groups of in Scotland, 1974-6, 115

Salmonella strains, virulence for and excretion by chickens, 479

Salmonellas in sewage and sewage sludges; the occurrence, and significance to animal health of, 47

Sanitary dressings, a bacteriological survey of, and development of an effective means for their disposal, 41

Selective decontamination of the digestive tract in mice; the dose at which neomycin and polymyxin B can be applied, 331

Serodiagnosis of bovine brucellosis; the use of non-specific immunological factors in, 405

Serological typing applied to the investigation of outbreaks of Clostridium perfringens food poisoning, 1970–1978, 443

 $Serratia\ marcescens$ from outbreaks of infection in hospitals; epidemiological type identification of, 269

Sewage and sewage sludges, the occurrence and significance to animal health of salmonellas in,
47

Simulating strategies for control of *Echniococcus granulosus*, *Taenia hydatigena* and *T. ovis*, 389

Skin diseases, contaminated medicaments in use in a hospital for, 103

Statistical analysis of comparative field trials of acute rodenticides, 341

Streptococcal infection, an outbreak in a chicken factory, 71

Swine vesicular disease: pathways of infection, 355

Toxocara spp., a comparison of flotation procedures for recovering ova from soil, 83 Toxoplasma antibody in the South Australian population, age-sex distribution of, 315 Trimethoprim resistance, plasmid-borne, in antibiotic-resistant Escherichia coli in market pigs, 1956-1979, 467

Type identification of Serratia marcescens from outbreaks of infection in hospitals, 269 Typhoid vaccine; intradermal versus subcutaneous immunization with, 11

Urine, cultural examination of using a single plate (the Macba plate), 429

Vaccine, polyvalent influenza; protection by and persistence of homologous and heterologous HI antibodies during two epidemic seasons, 237

Virulence of salmonella strains for chickens; their exerction by infected chickens, 479 Viruses in municipal effluent, inactivation by chlorine, 63

Viruses in small British field rodents, evidence of infection by, 285

Warfarin-resistant and non-resistant rats and mice: evaluation of bromadiolone as a rodenticide for use against, 263

West African rural environment: commercial milk products and indigenous weaning foods in, 191