



# All Irrational Extended Rotation Algebras are AF Algebras

George A. Elliott and Zhuang Niu

*Abstract.* Let  $\theta \in [0, 1]$  be any irrational number. It is shown that the extended rotation algebra  $\mathcal{B}_\theta$  introduced by the authors in *J. Reine Angew. Math.* **665**(2012), pp. 1–71, is always an AF algebra.

## 1 Introduction

In [8], a natural embedding of the irrational rotation  $C^*$ -algebra in a simple nuclear  $C^*$ -algebra with trivial  $K_1$ -group was constructed, which gives rise to an isomorphism of ordered  $K_0$ -groups. For a dense  $G_\delta$  of irrational numbers, the  $C^*$ -algebra constructed (by adjoining natural spectral projections of the canonical unitary generators of the rotation algebra) was shown to be AF.

In the present paper, using the remarkable recent work of Matui and Sato ([18], [14], [13]), together with what might be called the Winter–Lin–Niu deformation technique ([20], [11], [12]), the new  $C^*$ -algebra is shown to be AF for every irrational number. (See Corollary 4.7.)

By Remark 2.8 of [8], the flip of the rotation algebra extends to the larger algebra, and it is easily seen that the automorphism of order four known as the Fourier transform does also. It is an interesting question whether the whole of the natural  $SL(2, \mathbb{Z})$ -action also extends. Also of interest is whether the (unique) extendibility of, say, the flip automorphism, determines the embedding up to an automorphism. (Note that, as pointed out in [9] and [7], all embeddings are approximately unitarily equivalent, but the question of when two differ by an automorphism—in particular, ours and the very concrete (if not absolutely unique) one of Pimsner and Voiculescu constructed in [16]—is clearly an important question—analogue perhaps to the basic question in subfactor theory.)

## 2 Irrational Extended Rotation Algebras

Consider the  $C^*$ -algebra  $C(\mathbb{T})$  as the canonical sub- $C^*$ -algebra of  $L^\infty(\mathbb{T})$ , and denote by  $\sigma$  the automorphism of  $L^\infty(\mathbb{T})$  induced by translation by  $e^{2\pi i\theta}$ :

$$f(z) \mapsto f(e^{2\pi i\theta} z).$$

---

Received by the editors February 5, 2014.

Published electronically September 15, 2014.

This research was supported by the Natural Sciences and Engineering Research Council of Canada and the Fields Institute for Research in Mathematical Sciences.

AMS subject classification: 46L05, 46L35, 46L55, 46L80.

Keywords: irrational rotation algebra, extended irrational rotation algebra, AF-embedding.

Note that  $C(\mathbb{T})$  is invariant under the action of  $\sigma$ .

Consider two collections of (closed, open, or half-open) subintervals  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  of  $\mathbb{T}$ , and still denote again by  $f_i$  and  $g_j$  the spectral projections of the canonical unitary  $f(z) = z$  in  $L^\infty(\mathbb{T})$  corresponding to the subintervals  $f_i$  and  $g_j$ .

Consider the following two commutative  $C^*$ -algebras:

$$C(\Omega_u) := C^*(C(\mathbb{T}) \cup \{\sigma^{-k}(f_i) ; i \in \Lambda_1, k \in \mathbb{Z}\}) \subseteq L^\infty(\mathbb{T})$$

and

$$C(\Omega_v) := C^*(C(\mathbb{T}) \cup \{\sigma^k(g_j) ; j \in \Lambda_2, k \in \mathbb{Z}\}) \subseteq L^\infty(\mathbb{T}),$$

where  $\Omega_u$  and  $\Omega_v$  denote the spectra of these algebras. Note that the rotation  $\sigma$  can be extended to an automorphism of  $C(\Omega_u)$  (or  $C(\Omega_v)$ ). Denote also by  $u$  and  $v$  the canonical generators of  $C(\mathbb{T})$  inside  $C(\Omega_u)$  and  $C(\Omega_v)$ .

**Definition 2.1** For an irrational number  $\theta$ , and two collections of subintervals  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  of the unit circle  $\mathbb{T}$ , let us (as in [8]) refer to the universal  $C^*$ -algebra generated by  $C(\Omega_u)$  and  $C(\Omega_v)$  with respect to the relations

- (i)  $uv = e^{2\pi i \theta} v u$ ,
- (ii)  $u\sigma^k(g_j)u^* = \sigma^{k+1}(g_j)$  for any  $j \in \Lambda_2$  and  $k \in \mathbb{Z}$ , and
- (iii)  $v\sigma^{-k}(f_i)v^* = \sigma^{-k-1}(f_i)$  for any  $i \in \Lambda_1$  and  $k \in \mathbb{Z}$ ,

as the (irrational) extended rotation algebra, and denote it by  $\mathcal{B}_\theta (= \mathcal{B}_\theta(\{f_i\}, \{g_j\}))$ .

**Remark 2.2** If the intervals in the collections  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are arbitrary mixtures of open, closed, or half-open, then by [8, 5.14], there is a short exact sequence

$$0 \longrightarrow \bigoplus \mathcal{K} \longrightarrow \mathcal{B}_\theta \longrightarrow \mathcal{B}'_\theta \longrightarrow 0,$$

where  $\mathcal{B}'_\theta$  is an extended rotation algebra which can be generated by half-open intervals with the same orientation, and  $\mathcal{K}$  is the algebra of compact operators. In this paper, we will show that the  $C^*$ -algebra  $\mathcal{B}'_\theta$  is an AF algebra, and hence (by [3] and [5]) the  $C^*$ -algebra  $\mathcal{B}_\theta$  is AF as well.

If  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation, then there is another set of generators and relations for  $\mathcal{B}_\theta$ , as we shall now describe.

By Lemma 2.3 of [8], there is a  $\theta$ -independent set of real numbers  $\{a_k ; k \in \Lambda_u\}$  for some countable index set  $\Lambda_u$  (finite or infinite) such that the  $C^*$ -algebra  $C(\Omega_u)$  is generated by

$$\{\sigma^n(p_k), \sigma^n(e_k) ; n \in \mathbb{Z}, k \in \Lambda_u\},$$

where  $p_k$  is the spectral projection corresponding to  $[a_k, a_k + \theta)$  or  $(a_k, a_k + \theta]$ , and  $e_k$  is the minimal projection corresponding to  $\{a_k\}$  or zero. Since the half-open intervals  $f_i$ ,  $i \in \Lambda_1$ , are chosen to have the same orientation, the projection  $e_k$  is always zero. Let us refer to the points  $\{a_k ; k \in \Lambda_u\}$  as the cutting points of the canonical unitary  $u$ . A similar argument also works for  $C(\Omega_v)$ ; let  $\{b_l ; l \in \Lambda_v\}$  denote the corresponding cutting points of  $v$ , where  $\Lambda_v$  is a countable index set.

**Theorem 2.3** ([8, Corollary 2.10]) *Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. The  $C^*$ -algebra  $\mathcal{B}_\theta$  is the universal  $C^*$ -algebra generated by unitaries  $u$  and  $v$  and positive elements  $\{h_{u,k} ; k \in \Lambda_u\}$  and  $\{h_{v,l} ; l \in \Lambda_v\}$  with respect to the relations*

- (i)  $uv = e^{2\pi i\theta}vu$ ,
- (ii)  $\|h_{u,k}\| = \|h_{v,l}\| = 1$ ,
- (iii)  $u = e^{2\pi i(h_{u,k} + a_k)}$ , and
- (iv)  $v = e^{2\pi i(h_{v,l} + b_l)}$ ,

where  $\{a_k\}$  and  $\{b_l\}$  are as above.

For the extended rotation algebra  $\mathcal{B}_\theta$  in the case considered in Theorem 2.3 (which we shall usually consider now, unless otherwise specified—see Proposition 2.5 and Corollary 4.7), one has the following.

**Theorem 2.4** ([8, Theorems 3.6 and 5.1]) *Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. The  $C^*$ -algebra  $\mathcal{B}_\theta$  is simple and nuclear, and has a unique tracial state  $\tau$ .*

Consider the two  $C^*$ -algebras

$$B_u := C^*\{f, u ; f \in C(\Omega_v)\} \quad \text{and} \quad B_v := C^*\{f, v ; f \in C(\Omega_u)\}.$$

Then one has

$$B_u = C(\Omega_v) \rtimes_\sigma \mathbb{Z} \quad \text{and} \quad B_v = C(\Omega_u) \rtimes_\sigma \mathbb{Z}.$$

Both  $B_u$  and  $B_v$  contain the rotation algebra  $A_\theta$ , and one also has that  $\mathcal{B}_\theta = B_u *_{A_\theta} B_v$ .

**Proposition 2.5** (Propositions 3.4 and 5.3 of [8]) *There exist conditional expectations  $\mathbb{E}_u : \mathcal{B}_\theta \rightarrow C(\Omega_u)$  and  $\mathbb{E}_v : \mathcal{B}_\theta \rightarrow C(\Omega_v)$ . Moreover, if  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are half-open subintervals with the same orientation, then  $\mathbb{E}_u$  and  $\mathbb{E}_v$  are faithful.*

### 3 Strict Comparison of Positive Elements

In this section, we shall show that any irrational extended rotation algebra has strict comparison for positive elements (Theorem 3.8). The technique we are going to use is that of the (one-sided) large sub- $C^*$ -algebra due to N. C. Phillips (based on the method of Putnam in [17]; see [15]). Since (as shown in [8]) irrational extended rotation algebras are simple and nuclear, and have a unique tracial state, by a result of Matui and Sato these  $C^*$ -algebras are  $\mathcal{Z}$ -stable, *i.e.*,  $\mathcal{B}_\theta \otimes \mathcal{Z} \cong \mathcal{B}_\theta$ .

**Lemma 3.1** *Let  $q$  be a spectral projection of  $v$ , and let  $\varepsilon > 0$ . Then there is  $\delta > 0$  such that if  $e$  is a spectral projection of  $u$  with support in an interval with length at most  $\delta$ , one has*

$$\|eqe - \tau(q)e\| < \varepsilon,$$

where  $\tau$  is the canonical tracial state.

**Proof** The proof is almost the same as the proof of Lemma 5.7 of [8]. Let  $f, g \in C^*(v)$  be such that  $f \geq q \geq g$  and

$$\tau(f - g) < \varepsilon/2.$$

Choose polynomials  $F(v)$  and  $G(v)$  such that

$$\|f - F\| < \varepsilon/2 \quad \text{and} \quad \|g - G\| < \varepsilon/2.$$

Denote by  $n$  the larger of the degrees of  $F$  and  $G$ . Since  $\theta$  is irrational, there is  $\delta > 0$  such that if  $e$  is a spectral projection of  $u$  with support in an interval with length at most  $\delta$ , one has  $ev^i e = 0$  for all  $1 \leq i \leq n$ . In particular, this implies that  $eFe = F(0)e$  and  $eGe = G(0)e$ , where  $F(0)$  and  $G(0)$  are the constant terms of  $F$  and  $G$ , respectively. Note that

$$|F(0) - \tau(q)| < \varepsilon/2 \quad \text{and} \quad |G(0) - \tau(q)| < \varepsilon/2,$$

and also note that

$$F(0)e = eFe \approx_{\varepsilon/2} efe \geq eqe \geq ege \approx_{\varepsilon/2} eGe = G(0)e.$$

Then

$$(\tau(q) + \varepsilon)e \geq eqe \geq (\tau(q) - \varepsilon)e,$$

which is the conclusion of the lemma. ■

**Lemma 3.2** *Let  $a \in \mathcal{B}_\theta$ . For any  $\varepsilon > 0$ , there is  $\delta$  such that if  $e$  is a spectral projection of  $u$  with support in an interval with length at most  $\delta$ , then*

$$\|eae - \mathbb{E}_u(a)e\| < \varepsilon$$

(where  $\mathbb{E}_u$  is the canonical conditional expectation from  $\mathcal{B}_\theta$  to  $C(\Omega_u)$ ).

**Proof** Choose  $\sum_{j=1}^n c_j p_j q_j$  such that

$$\left\| a - \sum_{j=1}^n c_j p_j q_j \right\| < \varepsilon/3.$$

Then, by Lemma 3.1, there is  $\delta > 0$  such that if  $e$  is a spectral projection of  $u$  with support in an interval with length at most  $\delta$ , then

$$\left\| e \left( \sum_{j=1}^n c_j p_j q_j \right) e - \left( \sum_{j=1}^n c_j p_j \tau(q_j) \right) e \right\| < \varepsilon/3.$$

Note that

$$\mathbb{E}_u \left( \sum_{j=1}^n c_j p_j q_j \right) = \sum_{j=1}^n c_j p_j \tau(q_j)$$

and, therefore,

$$\begin{aligned} \|eae - \mathbb{E}_u(a)e\| &\leq \left\| e \left( \sum_{j=1}^n c_j p_j q_j \right) e - \mathbb{E}_u \left( \sum_{j=1}^n c_j p_j q_j \right) e \right\| + 2\varepsilon/3 \\ &\leq \varepsilon, \end{aligned}$$

as desired. ■

**Lemma 3.3** Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  each with the same orientation. Let  $b$  be a non-zero positive element of  $\mathcal{B}_\theta$ . Then there is a non-zero element  $a \in C(\Omega_u)$  such that  $a \preceq b$ . In particular, the  $C^*$ -algebra  $\mathcal{B}_\theta$  has the property (SP).

**Proof** By Proposition 2.5, the conditional expectation  $\mathbb{E}_u$  is faithful. Choose  $\varepsilon$  with

$$0 < \varepsilon < \|\mathbb{E}_u(b)\|/2.$$

By Lemma 3.2, there is  $e \in C(\Omega_u)$  such that

$$\|ebe - \mathbb{E}_u(b)e\| < \varepsilon.$$

Moreover, again since  $\mathbb{E}_u(b) \neq 0$ , the spectral projection  $e$  can be chosen so that

$$|\|\mathbb{E}_u(b)e\| - \|\mathbb{E}_u(b)\|| < \frac{\varepsilon}{2}.$$

Put

$$a' := \mathbb{E}_u(b)e \in C(\Omega_u)$$

and set  $a = (a' - \varepsilon)_+$ . Then

$$a \preceq ebe \preceq b.$$

Note that

$$\|a'\| = \|\mathbb{E}_u(b)e\| > \|\mathbb{E}_u(b)\| - \frac{\varepsilon}{2} > \frac{3\varepsilon}{2},$$

and hence  $a \neq 0$ , as desired. ■

**Lemma 3.4** Let  $A$  be a  $C^*$ -algebra and let  $a_1, a_2, \dots, a_n \in A$ . Then

$$(a_1 + \dots + a_n)^*(a_1 + \dots + a_n) \leq na_1^*a_1 + \dots + na_n^*a_n.$$

In particular,

$$(a_1 + \dots + a_n)^*(a_1 + \dots + a_n) \preceq a_1^*a_1 \oplus \dots \oplus a_n^*a_n.$$

**Proof** For any  $a, b \in A$ , since  $(a - b)^*(a - b) \geq 0$ , one has

$$a^*b + b^*a \leq a^*a + b^*b.$$

Then

$$\begin{aligned} (a_1 + \dots + a_n)^*(a_1 + \dots + a_n) &= \sum_{i=1}^n a_i^*a_i + \sum_{i < j} (a_i^*a_j + a_j^*a_i) \\ &\leq \sum_{i=1}^n a_i^*a_i + \sum_{i < j} (a_i^*a_i + a_j^*a_j) \\ &= n \sum_{i=1}^n a_i^*a_i, \end{aligned}$$

as desired. ■

**Lemma 3.5** Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. For any  $a \in \mathcal{B}_\theta$ , any  $\varepsilon > 0$ , and any  $b \in \mathcal{B}_\theta \setminus \{0\}$ , there are  $c \in A_\theta$  and  $g \in \mathcal{B}_\theta$  such that

- (i)  $\|a - (c + g)\| < \varepsilon$ ,
- (ii)  $g^*g \preceq b$  in  $\mathcal{B}_\theta$ .

**Proof** By Lemma 3.3, one may assume that  $b \in C(\Omega_u)$ , and hence one may assume that  $b$  is a projection in  $A_\theta$ .

By Theorem 1 of [9], one can choose  $\sum_{i=1}^n c_i p_i q_i$  such that

$$\left\| a - \sum_{i=1}^n c_i p_i q_i \right\| < \varepsilon/2,$$

where  $p_i$  and  $q_i$  are spectral projections in  $C(\Omega_u)$  and  $C(\Omega_v)$ , respectively. Then choose a projection  $e \in A_\theta$  with  $e < b$  and  $4n[e] < [b]$ .

Since (by [17]) the  $C^*$ -algebra  $C(\Omega_u) \rtimes \mathbb{Z}$  is an AT algebra, it has strict comparison of positive elements. Then, for each spectral projection  $p_i$ , we can choose  $f_i^-$ ,  $f_i$  and  $f_i^+$  such that

$$p_i = f_i^- + f_i + f_i^+,$$

with  $f_i \in C^*(u)$  and

$$f_i^-, f_i^+ \preceq e, \quad i = 1, \dots, n.$$

Similarly, one also chooses  $g_i^-, g_i$  and  $g_i^+$  such that

$$q_i = g_i^- + g_i + g_i^+,$$

with  $g_i \in C^*(v)$  and

$$g_i^-, g_i^+ \preceq e, \quad i = 1, \dots, n.$$

Then,

$$\begin{aligned} \sum_{i=1}^n c_i p_i q_i &= \sum_{i=1}^n c_i (f_i^- + f_i + f_i^+) (g_i^- + g_i + g_i^+) \\ &= \sum_{i=1}^n c_i f_i g_i + \sum_{i=1}^n c_i f_i^- q_i + \sum_{i=1}^n c_i f_i^+ q_i + \sum_{i=1}^n c_i f_i q g_i^- + \sum_{i=1}^n c_i f_i q g_i^+. \end{aligned}$$

Put

$$c = \sum_{i=1}^n c_i f_i g_i$$

and

$$g = \sum_{i=1}^n c_i f_i^- q_i + \sum_{i=1}^n c_i f_i^+ q_i + \sum_{i=1}^n c_i f_i q g_i^- + \sum_{i=1}^n c_i f_i q g_i^+.$$

By Lemma 3.4, one has

$$\begin{aligned} g^*g &\preceq \bigoplus_{i=1}^n f_i^- \oplus \bigoplus_{i=1}^n f_i^+ \oplus \bigoplus_{i=1}^n g_i^- \oplus \bigoplus_{i=1}^n g_i^+ \\ &\preceq \bigoplus_{i=1}^{4n} e \preceq b, \end{aligned}$$

as desired. ■

We thank N. C. Phillips for communicating to us the following two lemmas ([15, Lemmas 1.13 and 1.15]).

**Lemma 3.6** (1.13 of [15]) *Let  $A$  be a  $C^*$ -algebra, let  $a, b \in A$  be positive, and let  $\alpha, \beta \geq 0$ . Then*

$$(a + b - (\alpha + \beta))_+ \preceq (a - \alpha)_+ + (b - \beta)_+ \preceq (a - \alpha)_+ \oplus (b - \beta)_+.$$

**Lemma 3.7** (1.15 of [15]) *Let  $A$  be a  $C^*$ -algebra, and let  $a, b \in A$  be such that  $0 \leq a \leq b$ . Let  $\varepsilon > 0$ . Then  $(a - \varepsilon)_+ \preceq (b - \varepsilon)_+$ .*

**Theorem 3.8** *Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. The irrational extended rotation algebra  $\mathcal{B}_\theta$  has strict comparison of positive elements.*

**Proof** Let  $a, b$  be positive elements of  $\mathcal{B}_\theta$  (or of a matrix algebra over  $\mathcal{B}_\theta$ ) such that  $d_\tau(a) < d_\tau(b) - \delta$  for some  $\delta > 0$ , where  $\tau$  is the canonical trace.

Suppose that 0 is not an isolated point of  $\text{sp}(b)$ . Choose  $\delta_1 > 0$  such that

$$d_\tau((b - \eta)_+) > d_\tau(b) - \delta/4, \quad \text{for all } \eta \in (0, \delta_1).$$

Fix  $\varepsilon > 0$  with  $\varepsilon < \delta_1/9$ . Since 0 is not an isolated point of  $\text{sp}(b)$ , we may also assume that  $h_{(0, \varepsilon/2)}(b) \neq 0$  and  $h_{(\varepsilon/2, \varepsilon)}(b) \neq 0$  for continuous positive functions  $h_{(0, \varepsilon/2)}$  and  $h_{(\varepsilon/2, \varepsilon)}$  with support in  $(0, \varepsilon/2)$  and  $(\varepsilon/2, \varepsilon)$ , respectively.

By Lemma 3.5, there are  $b_0 \in A_\theta$  and  $b_1 \in \mathcal{B}_\theta$  such that  $\|(b - \varepsilon)^{1/2} - (b_0 + b_1)\|$  is sufficiently small that

$$(3.1) \quad \|(b - \varepsilon)_+ - (b_0 + b_1)^*(b_0 + b_1)\| < \varepsilon$$

and also

$$(3.2) \quad b_1^* b_1 \preceq h_{(0, \varepsilon/2)}(b).$$

Moreover, we may assume that

$$(3.3) \quad \|(b - 8\varepsilon)_+ - ((b_0 + b_1)^*(b_0 + b_1) - 7\varepsilon)_+\| < \varepsilon.$$

Then, by Lemma 3.4,

$$b_0^* b_0 = (b_0 + b_1 - b_1)^*(b_0 + b_1 - b_1) \leq 2(b_0 + b_1)^*(b_0 + b_1) + 2b_1^* b_1,$$

and then by Lemma 3.7 and Lemma 3.6,

$$(3.4) \quad \begin{aligned} (b_0^* b_0 - 3\varepsilon)_+ &\preceq (2(b_0 + b_1)^*(b_0 + b_1) + 2b_1^* b_1 - 3\varepsilon)_+ \\ &\preceq (2(b_0 + b_1)^*(b_0 + b_1) - 2\varepsilon)_+ + (2b_1^* b_1 - \varepsilon)_+ \\ &\preceq (b - \varepsilon)_+ + h_{(0, \varepsilon/2)}(b) \quad (\text{by (3.1) and (3.2)}). \end{aligned}$$

In particular,

$$d_\tau((b_0^* b_0 - 3\varepsilon)_+) \leq d_\tau(b).$$

On the other hand,

$$\begin{aligned}
 (3.5) \quad (b - 9\varepsilon)_+ &= ((b - 8\varepsilon)_+ - \varepsilon)_+ \\
 &\preceq ((b_0 + b_1)^*(b_0 + b_1) - 7\varepsilon)_+ \quad (\text{by (3.3)}) \\
 &\preceq (2b_0^*b_0 + 2b_1^*b_1 - 7\varepsilon)_+ \\
 &\preceq (2b_0^*b_0 - 6\varepsilon)_+ + (2b_1^*b_1 - \varepsilon)_+ \\
 &\preceq (b_0^*b_0 - 3\varepsilon)_+ \oplus b_1^*b_1.
 \end{aligned}$$

In particular,

$$d_\tau((b - 9\varepsilon)_+) \leq d_\tau((b_0^*b_0 - 3\varepsilon)_+) + d_\tau(b_1^*b_1),$$

and hence

$$\begin{aligned}
 (3.6) \quad d_\tau((b_0^*b_0 - 3\varepsilon)_+) &\geq d_\tau((b - 9\varepsilon)_+) - d_\tau(b_1^*b_1) \\
 &> d_\tau(b) - \delta/4 - \delta/4 = d_\tau(b) - \delta/2.
 \end{aligned}$$

Applying Lemma 3.5 to  $a^{1/2}$ , we define  $a_0 \in A_\theta$  and  $a_1 \in \mathcal{B}_\theta$  such that

$$\begin{aligned}
 \|a - (a_0 + a_1)^*(a_0 + a_1)\| &< \varepsilon, \\
 a_1^*a_1 &\preceq h_{(\varepsilon/2, \varepsilon)}(b),
 \end{aligned}$$

and

$$\|(a - 7\varepsilon)_+ - ((a_0 + a_1)^*(a_0 + a_1) - 7\varepsilon)_+\| < \varepsilon.$$

Then the same argument as for (3.5) shows that

$$(3.7) \quad (a - 8\varepsilon)_+ \preceq (a_0^*a_0 - 3\varepsilon)_+ + a_1^*a_1,$$

and since

$$a_0^*a_0 \leq 2(a_0 + a_1)^*(a_0 + a_1) + 2a_1^*a_1,$$

the same argument as for (3.4) shows that

$$\begin{aligned}
 (a_0^*a_0 - 3\varepsilon)_+ &\preceq ((a_0 + a_1)^*(a_0 + a_1) - \varepsilon) + (a_1^*a_1 - \varepsilon/2)_+ \\
 &\preceq a \oplus a_1^*a_1.
 \end{aligned}$$

Therefore, by (3.6),

$$\begin{aligned}
 d_\tau((a_0^*a_0 - 3\varepsilon)_+) &\leq d_\tau(a) + d_\tau(a_1^*a_1) < d_\tau(a) + \delta/2 \\
 &< d_\tau(b) - \delta/2 < d_\tau((b_0^*b_0 - 3\varepsilon)_+).
 \end{aligned}$$

Note that  $(a_0^*a_0 - 3\varepsilon)_+ \in A_\theta$  and  $(b_0^*b_0 - 3\varepsilon)_+ \in A_\theta$ . Since (by [6] or [2])  $A_\theta$  has strict comparison for positive elements, one has

$$(a_0^*a_0 - 3\varepsilon)_+ \preceq (b_0^*b_0 - 3\varepsilon)_+,$$

and hence

$$\begin{aligned}
 (a - 8\varepsilon)_+ &\preceq (a_0^*a_0 - 3\varepsilon)_+ + a_1^*a_1 \quad (\text{by (3.7)}) \\
 &\preceq (b_0^*b_0 - 3\varepsilon)_+ + a_1^*a_1 \\
 &\preceq (b - \varepsilon)_+ + h_{(0, \varepsilon/2)}(b) + h_{(\varepsilon/2, \varepsilon)}(b) \quad (\text{by (3.4)}) \\
 &\preceq b.
 \end{aligned}$$

Since (by Lemma 3.3)  $\varepsilon$  is arbitrary, and the left side converges to  $a$  as  $\varepsilon$  converges to zero, by inspection of the definition of Cuntz comparison, one has

$$a \preceq b.$$

Suppose now that 0 is an isolated point of  $\text{sp}(b)$ . Then the range projection of  $b$  in the bidual of  $\mathcal{B}_\theta$  belongs to  $\mathcal{B}_\theta$ , and is Cuntz equivalent to  $b$ , and so we may assume that  $b$  is a projection. Since (by Lemma 3.3)  $\mathcal{B}_\theta$  has property (SP), if  $b \neq 0$  (as we may suppose), there is a non-zero projection  $p < b$  such that  $\tau(p) < \tau(b) - d_\tau(a)$ . Pick a positive element  $c \in p\mathcal{B}_\theta p$  with  $\text{sp}(c) = [0, 1]$ , and consider the positive element

$$b' := (b - p) + c.$$

Then

$$d_\tau(b') \geq \tau(b - p) > d_\tau(a),$$

$\text{sp}(b') = [0, 1]$ , and  $b' < b$ . By the first part of the proof, one has that

$$a \preceq b'.$$

Since  $b' < b$ , we again have  $a \preceq b$ . ■

**Corollary 3.9** *Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. Then the irrational extended rotation algebra  $\mathcal{B}_\theta$  is  $\mathcal{Z}$ -stable.*

**Proof** By Theorem 5.1 and Theorem 3.6 of [8],  $\mathcal{B}_\theta$  is simple and has a unique tracial state. By Corollary 7.5 of [8],  $\mathcal{B}_\theta$  is nuclear. Hence, by Theorem 1.1 of [13],  $\mathcal{B}_\theta$  is  $\mathcal{Z}$ -stable. ■

### 4 Quasidiagonality and the UCT

In this section, let us show that any  $\mathcal{B}_\theta$  is quasidiagonal and satisfies the UCT. Then, by a result of Matui and Sato in [14] and a recent classification theorem, it will follow that the  $C^*$ -algebra  $\mathcal{B}_\theta$  is an AF algebra.

**Theorem 4.1** *Assume that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open subintervals of  $\mathbb{T}$  with the same orientation. Then, for any irrational  $\theta$ , the extended rotation algebra  $\mathcal{B}_\theta$  is quasidiagonal.*

**Proof** Since  $\mathcal{B}_\theta$  is nuclear ([8, 7.5]), it is enough to show that  $\mathcal{B}_\theta$  can be (unitaly) embedded into  $\prod_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C}) / \bigoplus_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C})$  for suitable natural numbers  $n_\lambda$ .

Let  $m_\lambda, n_\lambda$  be natural numbers such that  $m_\lambda/n_\lambda \rightarrow \theta$  as  $\lambda \rightarrow \infty$ . Set  $\omega_\lambda = e^{2\pi i m_\lambda/n_\lambda}$ ,

$$u_\lambda := \begin{pmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \ddots & \ddots & \\ 0 & 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \in M_{n_\lambda}(\mathbb{C}),$$

and

$$v_\lambda := \text{diag}\{\omega_\lambda, \omega_\lambda^2, \dots, \omega_\lambda^{n_\lambda}\} \in M_{n_\lambda}(\mathbb{C}).$$

For each  $k \in \Lambda_u$ , pick

$$h_{u,k,\lambda} = \frac{1}{2\pi i} \log(u_\lambda e^{-ia_k})$$

with  $0 \leq h_{u,k,\lambda} \leq 1$ , and for each  $l \in \Lambda_v$ , pick

$$h_{v,l,\lambda} = \frac{1}{2\pi i} \log(v_\lambda e^{-ib_l})$$

with  $0 \leq h_{v,l,\lambda} \leq 1$ .

Consider the elements

$$u := (\widetilde{u_\lambda}), \quad v := (\widetilde{v_\lambda}), \quad h_{u,k} := (\widetilde{h_{u,k,\lambda}}), \quad \text{and} \quad h_{v,l} := (\widetilde{h_{v,l,\lambda}})$$

in  $\prod_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C}) / \bigoplus_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C})$ . We shall show that that the  $C^*$ -algebra generated by these elements is isomorphic to  $\mathcal{B}_\theta$ , giving the desired embedding.

By Theorem 2.3, it is enough to show that  $u, v, h_{u,k}, h_{v,l}$  satisfy the relations

- (1)  $uv = e^{2\pi i\theta}vu$ ,
- (2)  $\|h_{u,k}\| = \|h_{v,l}\| = 1$ ,
- (3)  $u = e^{2\pi i(h_{u,k}+a_k)}$ , and
- (4)  $v = e^{2\pi i(h_{v,l}+b_l)}$ .

One only has to verify condition (1) since the other conditions are satisfied straightforwardly. A calculation shows that

$$u_\lambda v_\lambda u_\lambda^* v_\lambda^* = e^{2\pi i m_\lambda / n_\lambda},$$

and hence

$$\lim_{\lambda \rightarrow \infty} u_\lambda v_\lambda u_\lambda^* v_\lambda^* = \lim_{\lambda \rightarrow \infty} e^{2\pi i m_\lambda / n_\lambda} = e^{2\pi i\theta},$$

which implies

$$uv = e^{2\pi i\theta}vu$$

in  $\prod_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C}) / \bigoplus_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C})$ . Therefore, the elements  $u, v, h_{u,k}, h_{v,l}$  generate a copy of  $\mathcal{B}_\theta$  in  $\prod_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C}) / \bigoplus_{\lambda=1}^\infty M_{n_\lambda}(\mathbb{C})$ , as desired. ■

Let us now show that the  $C^*$ -algebra  $\mathcal{B}_\theta$  satisfies the UCT. It will be convenient to show at the same time, for use in the final classification, that  $K_1(\mathcal{B}_\theta) = \{0\}$  and that  $K_0(\mathcal{B}_\theta)$  is torsion free.

Note that  $\mathcal{B}_\theta = B_u *_A B_v$ . (In the case that there is only one cutting point for each of  $u$  and  $v$ , it follows directly from the Cuntz–Germain–Thomsen exact sequence that  $K_0(\mathcal{B}_\theta) = \mathbb{Z} + \theta\mathbb{Z}$  and  $K_1(\mathcal{B}_\theta) = \{0\}$ .) Denote by  $i_u$  and  $i_v$  the embeddings of  $A_\theta$  into  $B_u$  and  $B_v$ , respectively, and denote by  $j_u$  and  $j_v$  the embeddings of  $B_u$  and  $B_v$  into  $\mathcal{B}_\theta$ , respectively.

Before looking at  $K_*(\mathcal{B}_\theta)$ , let us consider the  $C^*$ -algebras  $B_u$  and  $B_v$ , and rewrite them as certain amalgamated free products.

Recall that  $B_u = C(\Omega_v) \rtimes_\sigma \mathbb{Z}$ , and suppose that there are only finitely many cutting points  $\{b_l; l \in \Lambda_v\}$  for the unitary  $v$ . Put  $0 = b_1 < b_2 < \dots < b_{|\Lambda_v|} < 1$ .

For each  $l \in \Lambda_\nu$ , denote by  $I_l$  the closed interval  $[b_l, b_{l+1}]$  (assume  $b_{|\Lambda_\nu|+1} = 1$ ), and consider the  $C^*$ -algebra  $\bigoplus_{l \in \Lambda_\nu} C(I_l)$ . For each  $l \in \Lambda_\nu$ , define a function  $h_l: [0, 1] \rightarrow [0, 1]$  by

$$h_l: t \mapsto \begin{cases} t - b_l, & \text{if } t \in \bigcup_{s \geq l} I_s, \\ t + (1 - b_l), & \text{otherwise.} \end{cases}$$

Then  $\{1, h_l; l \in \Lambda_\nu\}$  is a set of generators for the  $C^*$ -algebra  $\bigoplus_{l \in \Lambda_\nu} C(I_l)$ . Regard the unitary  $\nu$  as the function

$$t \mapsto e^{2\pi i t}, \quad t \in \bigcup_l I_l,$$

in  $\bigoplus_{l \in \Lambda_\nu} C(I_l)$ . Then a direct verification shows that

$$\nu = e^{2\pi i(h_l + b_l)}, \quad l \in \Lambda_\nu.$$

On the other hand, in the concrete  $C^*$ -algebra  $B_u$ , the commutative  $C^*$ -algebra  $C(\Omega_\nu)$  contains a copy of  $\bigoplus_{l \in \Lambda_\nu} C(I_l)$ . Let  $\bar{h}_l$  denote the generator in  $C(\Omega_\nu)$  corresponding to the element  $h_l$ . Then, there is a homomorphism  $\phi$  from the amalgamated free product  $A_\theta *_{C(\mathbb{T})} (\bigoplus_{l \in \Lambda_\nu} C(I_l))$  to  $B_u$  induced by

$$(4.1) \quad \phi(u) = u, \quad \phi(h_l) = \bar{h}_l, \quad l \in \Lambda_\nu.$$

Since the image contains  $\{u^{-n} \bar{h}_l u^n; n \in \mathbb{Z}, l \in \Lambda_\nu\}$ , it contains all the elements of  $C(\Omega_\nu)$ , and therefore  $\phi$  is surjective. Let us show that the map  $\phi$  is also injective.

**Lemma 4.2** *Under the assumption that  $|\Lambda_\nu| < \infty$ , the map  $\phi$  defined in (4.1) is injective. In particular, the  $C^*$ -algebra  $B_u$  is isomorphic to  $A_\theta *_{C(\mathbb{T})} (\bigoplus_{l \in \Lambda_\nu} C(I_l))$ .*

**Proof** The argument is similar to that of Theorem 2.9 of [8]. Set

$$B'_u = A_\theta *_{C(\mathbb{T})} \left( \bigoplus_{l \in \Lambda_\nu} C(I_l) \right).$$

Choose a faithful representation  $\pi$  of  $B'_u$  on some Hilbert space  $\mathcal{H}$ , and let us still use the same notation for the images of the elements of  $B'_u$  as for the elements themselves.

Since  $\nu = e^{2\pi i(h_l + b_l)}$ , one has

$$(4.2) \quad h_l = \frac{1}{2\pi i} \log(e^{-2\pi i b_l} \nu) + e_l,$$

where  $e_l$  is a subprojection of the spectral projection  $E_\nu(\{e^{2\pi i b_l}\})$ .

Consider the positive elements  $g_1 := f_1(h_l)$  and  $g_2 := f_2(h_l)$ , where

$$f_1(x) = \begin{cases} 0 & \text{if } 0 \leq x \leq 1/2, \\ \text{linear} & \text{if } 1/2 \leq x \leq 1 - \theta, \\ 1 & \text{otherwise,} \end{cases}$$

and

$$f_2(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq \theta, \\ \text{linear} & \text{if } \theta \leq x \leq 1/2, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$g_1 = f_1\left(\frac{1}{2\pi i} \log(v)\right) + e_l,$$

$$g_2 = f_2\left(\frac{1}{2\pi i} \log(v)\right) + (E_v(\{e^{2\pi i b_l}\}) - e_l),$$

and hence

(4.3)

$$g_1(ug_2u^*)$$

$$= \left( f_1\left(\frac{1}{2\pi i} \log(v)\right) + e_l \right) \left( u f_2\left(\frac{1}{2\pi i} \log(v)\right) u^* + u(E_v(\{e^{2\pi i b_l}\}) - e_l) u^* \right)$$

$$= f_1\left(\frac{1}{2\pi i} \log(v)\right) \cdot u f_2\left(\frac{1}{2\pi i} \log(v)\right) u^* + e_l + u(E_v(\{e^{2\pi i b_l}\}) - e_l) u^*$$

$$= E_v((b_l - \theta, b_l)) + e_l + u(E_v(\{e^{2\pi i b_l}\}) - e_l) u^*.$$

Therefore, the element  $g_1ug_2u^*$  is a projection. Let us define

$$d_{l,n} := u^{-n-1}(g_1vg_2u^*)u^{n+1}.$$

Then the elements

$$\{v, d_{l,n} ; n \in \mathbb{Z}, k \in \Lambda_v\}$$

satisfy the set of relations  $\mathcal{R}'$  of [8], and by Lemma 2.6 of [8], the  $C^*$ -algebra generated by  $\{v, d_{l,n}; n \in \mathbb{Z}, k \in \Lambda_v\}$  is isomorphic to  $C(\Omega_v)$  under the map

$$v \mapsto z, \quad d_{l,n} \mapsto \sigma^{-n}(\chi_{[b_l, b_l + \theta]}).$$

Therefore, there is a homomorphism

$$\psi: B_u \cong C(\Omega_v) \rtimes \mathbb{Z} \rightarrow B'_u$$

with

$$\psi(u) = u \quad \text{and} \quad \psi(\sigma^{-n}(\chi_{[b_l, b_l + \theta]})) = d_{l,n}.$$

In particular, by (4.2) and (4.3), one has

$$\psi(\bar{h}_l) = h_l, \quad l \in \Lambda_v,$$

and hence  $\psi \circ \phi = \text{id}_{B'_u}$ , which implies that the map  $\phi$  is injective. ■

**Lemma 4.3** Consider the  $C^*$ -algebra  $B_u$  (or  $B_v$ ). Let  $a \in K_0(B_u)$  such that  $na \in (i_u)_0(K_0(A_\theta))$  for some non-zero  $n \in \mathbb{N}$ . Then  $a \in (i_u)_0(K_0(A_\theta))$ .

**Proof** Assume that  $|\Lambda_v| < \infty$ . By Lemma 4.2 and Theorem 6.4 of [19], a straightforward calculation shows that the sequence

$$0 \longrightarrow K_0(C(\mathbb{T})) \xrightarrow{\iota} K_0(A_\theta) \oplus \left( \bigoplus K_0(C(I_l)) \right) \xrightarrow{(i_u)_0 - (\eta)_0} K_0(B_u)$$

is exact, where

$$\iota(1) = (0, 1) \oplus (1, \dots, 1),$$

and  $\eta$  is the embedding of  $\bigoplus_l C(I_l)$  into  $B_u$ .

Let  $(a, b) \oplus (c_1, \dots, c_{|\Lambda_v|-1}) \in K_0(A_\theta) \oplus (\bigoplus K_0(C(I_l)))$  be a representative of  $a$ . One then has that

$$((na, nb) \oplus (nc_1, \dots, nc_{|\Lambda_v|-1})) - ((a', b') \oplus (0, \dots, 0)) = (0, m) \oplus (m, \dots, m)$$

for some  $a', b', m \in \mathbb{Z}$ . In particular, this implies that  $m$  is divisible by  $n$ , and

$$c_1 = \dots = c_{|\Lambda_v|-1} = m/n.$$

Then the element

$$(a, b) \oplus (c_1, \dots, c_{|\Lambda_v|-1}) - (0, m/n) \oplus (m/n, \dots, m/n) = (a, b - m/n) \oplus (0, \dots, 0)$$

is still a representative of  $a$ , and it is in the image of  $K_0(A_\theta)$ , as desired.

If  $|\Lambda_v| = |\{b_1, b_2, \dots, b_i, \dots\}| = \infty$ , then for each  $n = 1, 2, \dots$ , denote by  $\Omega_{v,n}$  the commutative  $C^*$ -algebra generated by the spectral projections

$$\{\chi_{[b_i+k\theta, b_i+(k+1)\theta)} ; i = 1, \dots, n, k \in \mathbb{Z}\},$$

and consider the  $C^*$ -algebra crossed product

$$B_{u,n} := C(\Omega_{v,n}) \rtimes_\sigma \mathbb{Z}.$$

Then, as a sub- $C^*$ -algebra of  $B_u$ , each  $B_{u,n}$  contains  $A_\theta$ , and  $B_u = \overline{\bigcup_{n=1}^\infty B_{u,n}}$ . The conclusion follows from the preceding case, that there are only finitely many cutting points. ■

**Lemma 4.4** *With the setting as above, the following facts obtain:*

- (i)  $K_1(\mathcal{B}_\theta) = \{0\}$ , and  $K_0(\mathcal{B}_\theta)$  is torsion free.
- (ii) The map  $(i_u, -i_v): A_\theta \rightarrow B_u \oplus B_v$  induces an injective map on the  $K$ -groups.

**Proof** Since  $B_u = C(\Omega_v) \rtimes \mathbb{Z}$  and the action of  $\sigma$  on  $\Omega_v$  has no nontrivial clopen subset, a direct calculation using the Pimsner–Voiculescu six-term exact sequence shows that  $K_1(B_u)$  is isomorphic to  $\mathbb{Z}$ , and is generated by the canonical unitary  $u$ . The same argument also works for  $B_v$ . In particular, this implies that  $(i_u, -i_v)$  (or  $(i_u, i_v)$ ) induces an isomorphism between  $K_1(A_\theta)$  and  $K_1(B_u) \oplus K_1(B_v)$ .

For the injectivity on  $K_0$ -groups, by applying the standard trace on  $B_u$  (or  $B_v$ ), one has that the map  $\iota_u$  (or  $\iota_v$ ) also induces an embedding of  $K_0(A_\theta)$  into  $K_0(B_u)$  (or  $K_0(B_v)$ ). In particular, the map  $(i_u, -i_v)$  (or  $(i_u, i_v)$ ) induces an injective map from  $K_1(A_\theta)$  to  $K_1(B_u) \oplus K_1(B_v)$ .

By Theorem 6.4 of [19], one has the exact sequence

$$\begin{array}{ccccc} K_0(A_\theta) & \xrightarrow{(i_{u_0}, i_{v_0})} & K_0(B_u) \oplus K_0(B_v) & \xrightarrow{j_{u_0} - j_{v_0}} & K_0(\mathcal{B}_\theta) \\ \uparrow & & & & \downarrow \\ K_1(\mathcal{B}_\theta) & \xleftarrow{j_{u_1} - j_{v_1}} & K_1(B_u) \oplus K_1(B_v) & \xleftarrow{(i_{u_1}, i_{v_1})} & K_1(A_\theta). \end{array}$$

Since  $(i_{u_1}, i_{v_1}): K_1(A_\theta) \rightarrow K_1(B_u) \oplus K_1(B_v)$  is an isomorphism, one has that  $K_1(\mathcal{B}_\theta)$  embeds into  $K_0(A_\theta)$  with image the kernel of the map  $(i_{u_0}, i_{v_0}): K_0(A_\theta) \rightarrow K_0(B_u) \oplus K_0(B_v)$ . But the map  $(i_{u_0}, i_{v_0})$  is injective, as shown above. Therefore,  $K_1(\mathcal{B}_\theta) = \{0\}$ .

Let us show that  $K_0(\mathcal{B}_\theta)$  is torsion free. As shown above, one has that

$$K_0(\mathcal{B}_\theta) = (K_0(B_u) \oplus K_0(B_v)) / (i_{u0}, i_{v0})(K_0(A_\theta)).$$

Let  $(a, b) \in K_0(B_u) \oplus K_0(B_v)$  with  $\overline{n(a, b)} = 0$  for some nonzero  $n \in \mathbb{Z}$ ; that is

$$n(a, b) = ((i_u)_0(c), (i_v)_0(c))$$

for some  $c \in K_0(A_\theta)$ , and hence

$$na = (i_u)_0(c) \quad \text{and} \quad nb = (i_v)_0(c).$$

By Lemma 4.3, one has that

$$a \in (i_u)_0(K_0(A_\theta)) \quad \text{and} \quad b \in (i_v)_0(K_0(A_\theta)).$$

Denote by  $a', b' \in K_0(A_\theta)$  the preimages of  $a, b$ , respectively. Since the maps  $(i_u)_0$  and  $(i_v)_0$  are injective, one has

$$na' = c = nb'.$$

Since  $K_0(A_\theta)$  is torsion free, one has  $a' = b'$ , and therefore

$$(a, b) = ((i_u)_0(a'), (i_v)_0(b')) \in ((i_u)_0, (i_v)_0)(K_0(A_\theta)),$$

which implies

$$\overline{(a, b)} = 0 \in K_0(\mathcal{B}_\theta).$$

This shows that the group  $K_0(\mathcal{B}_\theta)$  is torsion free. ■

Let  $A, B$  be  $C^*$ -algebras. In what follows, let

$$\gamma(A, B): KK(A, B) \rightarrow \text{Hom}(K_*(A), K_*(B))$$

denote the canonical homomorphism. Let us also use the same notation for the analogous homomorphism with domain  $E(A, B)$ .

**Proposition 4.5** (23.8.1 of [1]) *Let  $A$  be a separable  $C^*$ -algebra. Suppose that for every separable  $C^*$ -algebra  $B$  with divisible  $K$ -groups,  $\gamma(A, B)$  is an isomorphism. Then for every separable  $C^*$ -algebra  $B$ , the exact sequence of the UCT holds for  $A$  and  $B$ .*

**Theorem 4.6** *For any irrational  $\theta$ , the extended rotation algebra  $\mathcal{B}_\theta$  satisfies the UCT.*

**Proof** Since  $\mathcal{B}_\theta$  is nuclear, the group  $E(\mathcal{B}_\theta, D)$  and  $KK(\mathcal{B}_\theta, D)$  are canonically isomorphic for any separable  $D$ . Therefore by Proposition 4.5, it is enough to show that  $\gamma(\mathcal{B}_\theta, D)$  is an isomorphism between  $E(\mathcal{B}_\theta, D)$  and  $\text{Hom}(K_*(\mathcal{B}_\theta), K_*(D))$  for any separable  $D$  with divisible  $K$ -groups.

By Theorem 6.3 of [19], one has the exact sequence

$$(4.4) \quad \begin{array}{ccccc} E(A_\theta, D) & \xleftarrow{i_u^* - i_v^*} & E(B_u, D) \oplus E(B_v, D) & \xleftarrow{(j_u^*, j_v^*)} & E(\mathcal{B}_\theta, D) \\ \downarrow & & & & \uparrow \\ E(\mathcal{B}_\theta, SD) & \xrightarrow{(j_u^*, j_v^*)} & E(B_u, SD) \oplus E(B_v, SD) & \xrightarrow{i_u^* - i_v^*} & E(A_\theta, SD). \end{array}$$

Applying the functor  $\gamma$  to the lower-right corner of (4.4), one has the commutative diagram

$$\begin{CD} E(B_u, SD) \oplus E(B_v, SD) @>{i_u^* - i_v^*}>> E(A_\theta, SD) \\ @V{\gamma(B_u, SD) \oplus \gamma(B_v, SD)}VV @VV{\gamma(A_\theta, SD)}V \\ \text{Hom}(K_*(B_u), K_*(SD)) \oplus \text{Hom}(K_*(B_v), K_*(SD)) @>{i_u^* - i_v^*}>> \text{Hom}(K_*(A_\theta), K_*(SD)). \end{CD}$$

By Lemma 4.4 (ii), the map  $(i_u, -i_v): A_\theta \rightarrow B_u \oplus B_v$  induces an embedding of  $K$ -groups. Then, since  $K_*(D)$  is divisible, the map  $i_u^* - i_v^*$  in the bottom row is a surjective homomorphism. Since the  $C^*$ -algebras  $B_u, B_v$ , and  $A_\theta$  are nuclear and satisfy the UCT, the vertical maps induced by the functor  $\gamma$  are isomorphisms, and therefore the map

$$i_u^* - i_v^*: E(B_u, SD) \oplus E(B_v, SD) \rightarrow E(A_\theta, SD)$$

must be surjective.

Then, by exactness of the sequence (4.4), the map  $E(A_\theta, SD) \rightarrow E(\mathcal{B}_\theta, D)$  is zero, and therefore the map

$$(j_u^*, j_v^*): E(\mathcal{B}_\theta, D) \rightarrow E(B_u, D) \oplus E(B_v, D)$$

is injective.

Let us consider the map  $\gamma(\mathcal{B}_\theta, D)$  and show that it is an isomorphism. Applying the functor  $\gamma$  to the top part of (4.4), one has the commutative diagram (4.5)

$$\begin{CD} \text{Hom}(K_*(A_\theta), K_*(D)) @<{i_u^* - i_v^*}<< \bigoplus_{\bullet=u,v} \text{Hom}(K_*(B_\bullet), K_*(D)) @<{(j_u^*, j_v^*)}<< \text{Hom}(K_*(\mathcal{B}_\theta), K_*(D)) \\ @V{\gamma(A_\theta, D)}VV @V{\gamma(B_u, D) \oplus \gamma(B_v, D)}VV @V{\gamma(\mathcal{B}_\theta, D)}VV \\ E(A_\theta, D) @<{i_u^* - i_v^*}<< \bigoplus_{\bullet=u,v} E(B_\bullet, D) @<{(j_u^*, j_v^*)}<< E(\mathcal{B}_\theta, D). \end{CD}$$

Since the map  $(j_u^*, j_v^*)$  in the bottom row is injective, and as before, the first two vertical maps (actually, we only need the middle one here) are isomorphisms, the map  $\gamma(\mathcal{B}_\theta, D)$  must be injective.

Let us show that  $\gamma(\mathcal{B}_\theta, D)$  is also surjective. Note that the sequence

$$0 \longrightarrow K_0(A_\theta) \xrightarrow{(i_u^*, -i_v^*)} K_0(B_u) \oplus K_0(B_v) \xrightarrow{j_u^* + j_v^*} K_0(\mathcal{B}_\theta) \longrightarrow 0$$

is exact, and  $K_1(\mathcal{B}_\theta) = \{0\}$ . Then a direct calculation shows that the top sequence of (4.5) is exact in the middle, and the map  $(j_u^*, j_v^*)$  (in the top row) is injective. Since the  $C^*$ -algebras  $A_\theta, B_u$ , and  $B_v$  satisfy the UCT, the maps  $\gamma(A_\theta, D), \gamma(B_u, D)$ , and  $\gamma(B_v, D)$  are isomorphisms. Let  $a \in \text{Hom}(K_0(\mathcal{B}_\theta), K_0(D))$ , and denote the image of  $a$  in  $E(B_u, D) \oplus E(B_v, D)$  by  $a'$ . Then, by the exactness of the top sequence, the element  $a'$  must be sent to 0 in  $E(A_\theta, D)$ , whence, by the exactness of the lower sequence, there is an element  $a'' \in E(\mathcal{B}_\theta, D)$  which is sent to  $a'$ . Since  $(j_u^*, j_v^*)$  is injective also at the level of  $\text{Hom}$  (in the top row), the element  $a''$  must be sent to  $a$  under the map  $\gamma(\mathcal{B}_\theta, D)$ . This shows that the map  $\gamma(\mathcal{B}_\theta, D)$  is surjective, as desired. ■

**Corollary 4.7** For arbitrary collections of sub-intervals  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$ , the irrational extended rotation algebra  $\mathcal{B}_\theta = \mathcal{B}_\theta(\{f_i\}, \{g_j\})$  is an AF algebra.

**Proof** Suppose that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of half-open intervals with the same orientation; then  $\mathcal{B}_\theta$  is simple, unital, nuclear, and has a unique tracial state. It is quasidiagonal by Theorem 4.1. Hence by [14, Theorem 6.1],  $\mathcal{B}_\theta \otimes \mathcal{Q}$  is TAF for the universal UHF algebra  $\mathcal{Q}$ . In other words,  $\mathcal{B}_\theta$  is rationally TAF.

By Theorem 4.6,  $\mathcal{B}_\theta$  satisfies the UCT; and by Corollary 3.9, it is  $\mathcal{Z}$ -stable. Therefore, it is covered by the classification theorem of [20], [11], and [12]. By Lemma 4.4 (i), the group  $K_1(\mathcal{B}_\theta)$  is zero and  $K_0(\mathcal{B}_\theta)$  is torsion free. Also, as  $\mathcal{B}_\theta$  is  $\mathcal{Z}$ -stable, by [10], the ordered group  $K_0(\mathcal{B}_\theta)$  is unperforated. Since  $\mathcal{B}_\theta$  has a unique tracial state, the ordered group  $K_0(\mathcal{B}_\theta)$  has a unique state. (It is the same to show that  $K_0(\mathcal{B}_\theta \otimes \mathcal{Q})$  has a unique state, but this holds as  $\mathcal{B}_\theta \otimes \mathcal{Q}$  has a unique tracial state and is TAF—see above.) Furthermore, the image of  $K_0(\mathcal{B}_\theta)$  is dense in  $\mathbb{R}$  (it contains the subgroup  $\mathbb{Z} + \mathbb{Z}\theta$ ). Therefore,  $K_0(\mathcal{B}_\theta)$  is a Riesz group.

It follows by [4] that there is an AF algebra with the same invariant, which is also covered by this classification theorem, and so the  $C^*$ -algebra  $\mathcal{B}_\theta$  is isomorphic to that AF algebra.

For the general case, that  $\{f_i\}_{i \in \Lambda_1}$  and  $\{g_j\}_{j \in \Lambda_2}$  are two collections of arbitrary intervals, by 5.14 of [8], there is a short exact sequence

$$0 \longrightarrow \bigoplus \mathcal{K} \longrightarrow \mathcal{B}_\theta \longrightarrow \mathcal{B}'_\theta \longrightarrow 0,$$

where  $\mathcal{B}'_\theta$  is an extended rotation algebra which can be generated by half-open intervals with the same orientation, and  $\mathcal{K}$  is the algebra of compact operators. Since the previous argument shows that  $\mathcal{B}'_\theta$  is an AF algebra, the  $C^*$ -algebra  $\mathcal{B}_\theta$  is an extension of AF algebras, and therefore (by [3] and [5]) it is an AF algebra as well. ■

## References

- [1] B. Blackadar, *K-theory for Operator Algebras*. Second edition, Math. Sci. Res. Inst. Publ. 5, Cambridge University Press, Cambridge, 1998.
- [2] B. Blackadar, A. Kumjian, and M. Rørdam, *Approximately central matrix units and the structure of noncommutative tori*. *K-Theory* **6**(1992), 267–284. <http://dx.doi.org/10.1007/BF00961466>
- [3] L. G. Brown, *Extensions of AF-algebras: the projection lifting problem*. In: *Operator algebras and applications Symposia in Pure Math.* **38**, Providence, RI, 1981, 175–716.
- [4] E. G. Effros, D. E. Handelman, and C. L. Shen, *Dimension groups and their affine representations*. *Amer. J. Math.* **102**(1980), 385–407. <http://dx.doi.org/10.2307/2374244>
- [5] G. A. Elliott, *Automorphisms determined by multipliers on ideals of a  $C^*$ -algebra*. *J. Funct. Anal.* **23**(1976), 1–10. [http://dx.doi.org/10.1016/0022-1236\(76\)90054-9](http://dx.doi.org/10.1016/0022-1236(76)90054-9)
- [6] G. A. Elliott and D. E. Evans, *The structure of the irrational rotation  $C^*$ -algebra*. *Ann. of Math.* (2) **138**(1993), 477–501. <http://dx.doi.org/10.2307/2946553>
- [7] G. A. Elliott and Z. Niu, *A canonical Pimsner-Voiculescu embedding*. In: *Oberwolfach Reports*, Mathematisches Forschungsinstitut Oberwolfach, 43, European Mathematical Society Publishing House, Zürich, 2013, pp. 2438–2440. <http://dx.doi.org/10.4171/OWR/2013/43>
- [8] G. A. Elliott and Z. Niu, *Extended rotation algebras: adjoining spectral projections to rotation algebras*. *J. Reine Angew. Math.* **665**(2012), 1–71. <http://dx.doi.org/10.1515/CRELLE.2011.112>
- [9] ———, *Remarks on the Pimsner-Voiculescu embedding*. In: *Operator Algebra and Dynamics (Nordforsk Network Closing Conference, Faroe Islands, May 2012)*, Springer Proceedings in Mathematics & Statistics **58**, 2013, pp. 131–140.
- [10] G. Gong, X. Jiang, and H. Su, *Obstructions to  $\mathcal{Z}$ -stability for unital simple  $C^*$ -algebras*. *Canad. Math. Bull.* **43**(2000), 418–426. <http://dx.doi.org/10.4153/CMB-2000-050-1>

- [11] H. Lin, *Localizing the Elliott conjecture at strongly self-absorbing  $C^*$ -algebras, II*. J. Reine Angew. Math. **2014**(2014) no. 692, 233–243. <http://dx.doi.org/10.1515/crelle-2012-0182>
- [12] H. Lin and Z. Niu, *Lifting KK-elements, asymptotic unitary equivalence and classification of simple  $C^*$ -algebras*. Adv. Math. **219**(2008), 1729–1769. <http://dx.doi.org/10.1016/j.aim.2008.07.011>
- [13] H. Matui and Y. Sato, *Strict comparison and  $\mathcal{Z}$ -absorption of nuclear  $C^*$ -algebras*. Acta Math. **209**(2012), 179–196. <http://dx.doi.org/10.1007/s11511-012-0084-4>
- [14] H. Matui and Y. Sato, *Decomposition rank of UHF-absorbing  $C^*$ -algebras*. Duke Math. J., to appear. [arxiv:1303.4371](https://arxiv.org/abs/1303.4371).
- [15] N. C. Phillips, *Large subalgebras*. Preprint, 2014.
- [16] M. Pimsner and D. Voiculescu, *Imbedding the irrational rotation  $C^*$ -algebra into an AF-algebra*. J. Operator Theory **4**(1980), 201–210.
- [17] I. F. Putnam, *The  $C^*$ -algebras associated with minimal homeomorphisms of the Cantor set*. Pacific J. Math. **136**(1989):329–353. <http://dx.doi.org/10.2140/pjm.1989.136.329>
- [18] Y. Sato, *Trace spaces of simple nuclear  $C^*$ -algebras with finite-dimensional extreme boundary*. Preprint, 2012. [arxiv:1209.3000](https://arxiv.org/abs/1209.3000).
- [19] K. Thomsen, *On the KK-theory and the E-theory of amalgamated free products of  $C^*$ -algebras*. J. Funct. Anal. **201**(2003), 30–56. [http://dx.doi.org/10.1016/S0022-1236\(03\)00084-3](http://dx.doi.org/10.1016/S0022-1236(03)00084-3)
- [20] W. Winter, *Localizing the Elliott conjecture at strongly self-absorbing  $C^*$ -algebras*. J. Reine Angew. Math. **2014**(2014) no. 692, 193–231. <http://dx.doi.org/10.1515/crelle-2012-0082>

*Department of Mathematics, University of Toronto, Toronto, ON M5S 2E4*  
e-mail: [elliott@math.toronto.edu](mailto:elliott@math.toronto.edu)

*Department of Mathematics, University of Wyoming, Laramie, WY 82071, USA*  
e-mail: [zniu@uwyo.edu](mailto:zniu@uwyo.edu)