

## O191

**Sensory experience in obsessive compulsive disorder - sensiocd: Do they think or feel differently?**

I. Soares Da Costa\*, V. Covelo and R. Moreira

Psychiatry And Mental Health Department, Centro Hospitalar e Universitário de São João, Porto, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.377

**Introduction:** Obsessive-compulsive disorder (OCD) is associated to a wide range of symptomatic expression and treatment response variability [1]. Sensory perception has been identified as an emerging factor in this process [2]. Sensory vulnerability and atypical sensory experience were identified as risk factors for the development of OCD [3] and a sensory subtype of the disease was proposed in which there is a positive correlation with early onset sensory symptoms, male gender and family background [4]. Adding to the atypical sensory profile, obsessions are frequently experienced as partially perceptual.

**Objectives:** Our main goals are to characterize the sensory perception in OCD patients; assess the prevalence and intensity of the sensory properties of the obsessive thoughts and explore the how sensory perception, obsessive thoughts and obsessive dimensions/clusters are interrelated.

**Methods:** Patients with OCD diagnosis, aged 18 to 65 years and no comorbid mental disorder (except depression) will be recruited. The study battery will include participant form with demographical and clinical features, assessment of depressive and anxiety symptoms (HAM-A and HAM-D) evaluation of clinical outcome measures and obsessive dimensions/clusters (Yale-Brown Obsessive-Compulsive Scale (Y-BOQS) and Obsessive Beliefs Questionnaire-44 (OBQ-44)), assessment of sensory perception and sensory properties of obsessive thoughts (Sensory Perception Quotient (SPQ 21) and Sensory Properties of Obsessive Thoughts Questionnaire (SPOQ)).

**Results:** The results will help us understand the interaction between perceptual and cognitive processes in OCD.

**Conclusions:** Better definition of OCD psychopathology and the establishment of a sensory subtype may indicate the need of specific therapeutic indications or a different escalation of treatment measures.

**Disclosure:** No significant relationships.

**Keywords:** ocd; sensory experience; obsessive spectrum; obsessive thoughts

## O193

**Can we reduce the stigmatisation experience with psychosocial interventions? An investigation of the meeting centre support programme impact on people with cognitive impairments**K. Lion<sup>1\*</sup>, D. Szcześniak<sup>2</sup>, S. Evans<sup>3</sup>, S. Evans<sup>3</sup>, E. Farina<sup>4</sup>, D. Brooker<sup>3</sup>, R. Chattat<sup>5</sup>, F. Meiland<sup>6</sup>, R.-M. Droes<sup>7</sup> and J. Rymaszewska<sup>2</sup>

<sup>1</sup>Menzies Health Institute Queensland, Griffith University, Nathan, Australia; <sup>2</sup>Department Of Psychiatry, Wroclaw Medical University, Wroclaw, Poland; <sup>3</sup>Association For Dementia Studies, University of Worcester, Worcester, United Kingdom; <sup>4</sup>Fondazione Don Carlo Gnocchi, IRCCS, Milan, Italy; <sup>5</sup>Department Of Psychology, University

of Bologna, Bologna, Italy; <sup>6</sup>Department Of Medicine For Older People, Amsterdam UMC University Medical Centres, Amsterdam, Netherlands and <sup>7</sup>Amsterdam Public Health Research Institute Department Of Psychiatry, Amsterdam UMC, Amsterdam, Netherlands

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.378

**Introduction:** People living with dementia or mild cognitive impairment (MCI) experience stigmatisation and there are not many specific psychosocial interventions dedicated to help them coping with this issue, reducing its impact on their lives.

**Objectives:** This study aimed to a) investigate the stigmatisation level among people with dementia and MCI in Poland, Italy and the United Kingdom and b) assess the role of the Meeting Centre Support Programme (MCSP) in decreasing stigmatisation.

**Methods:** We investigated outcomes for 114 people with dementia and MCI living in Italy, Poland and the UK who participated 6 months in MCSP or usual care (UC) using a pre/post-test control group study design. Level of stigmatisation was assessed with the Stigma Impact Scale: neurological impairment (SIS).

**Results:** Stigmatisation level (SIS) among participants varied from 2 to 65 (median=33.5; Q1=27; Q3=41) with people from the UK experiencing a statistically significantly higher level of stigmatisation than people in Italy and Poland. In Italy, stigmatisation was lower (p=0.02) in the MCSP group following the intervention. In Poland, the social isolation level did not significantly change in MCSP, but increased (p=0.05) in UC. In the UK, the social rejection level raised (p=0.03) in MCSP. Overall, the combined data of the three countries did not show statistically significant differences in SIS between MCSP and UC.

**Conclusions:** Stigmatisation among people with dementia and MCI is complex and seems culturally dependent. There is a great opportunity in psychosocial interventions to reduce the burden of stigma among people with dementia which requires further investigation.

**Disclosure:** No significant relationships.

**Keywords:** dementia; Stigma; attitude; social isolation

## O194

**Rates of 1-year cognitive impairment in older adults who developed delirium due to a systemic infection**A.R. Silva\*, A.L. Cardoso, I. Baldeiras, I. Santana and J. Cerejeira  
University Of Coimbra, Center for Neuroscience and Cell Biology, Coimbra, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.379

**Introduction:** Delirium affects a significant proportion of hospitalized older patients with acute infections. There is growing evidence that delirium accelerates the cognitive decline at long term.

**Objectives:** We aimed to determine if delirium during hospitalization was independently associated with cognitive deterioration at one-year.

**Methods:** From a total of 22 patients (12 C, 4 Dem, 2 D, and 4 DD) delirium (D and DD groups) was associated with a worse score in MOCA of 3-points (p<.02) and 2.5-points (p<.03), respectively, at one year, follow up. Dementia patients without delirium had a