Vol. LXXVII No. 1

# The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR

## Contents

J. C. MCALPINE, PRIMARY AMYLOIDOSIS OF THE UPPER AIR PASSAGES . A. RADCLIFFE and I. FRIEDMANN MYCOTIC INFECTION OF THE FRONTAL SINUS D. A. OSBORN JAMES LISTER TREATMENT OF MALIGNANT DISEASE OF THE NASAL SINUSES BY SUPERVOLTAGE RADIOTHERAPY THOMAS J. DEELEY and ROBERT MORRISON THE EARLY MANAGEMENT OF BELL'S PALSY BY CERVICAL SYMPATHETIC BLOCK . . . . F. BOYES KORKIS CLINICAL RECORDS-WAARDENBURG'S SYNDROME . J. CALINIKOS CHRONIC MIDDLE-EAR INFECTION-MASTOID SURGERY WITHOUT MEATAL FLAPS OR SKIN GRAFTS . . . HECTOR A. THOMAS SEPTAL DERMOPLASTY: A NEW OPERATIVE PROCEDURE FOR CONTROL OF NOSEBLERDS IN PATIENTS WITH HEREDITARY HÆMOR-WILLIAM H. SAUNDERS SOCIETIES' PROCEEDINGS-ROYAL SOCIETY OF MEDICINE-SECTION OF LARYNGOLOGY

London

# Headley Brothers

109 Kingsway WC2

GENERAL NOTES

# The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

EDITED BY

### G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C. and Brown, D. (1951) J. Laryng., 65, 55. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World

Health Organization, 1952.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of fro per article; beyond this authors are expected to pay half the cost. Coloured illustrations will

be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2.

5. Orders for reprints must be sent when returning galley proofs, and for this

purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the Journal of Laryngology. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of

Laryngology, c/o Headiey Brothers, 109 Kingsway, London, WCz.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free, and is payable in advance.

9. Single copies will be on sale at 10s, oil, each; copies of parts up to Vol.

LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.


United States of America

Orders for this Journal may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-35 East 10th Street, New York, or direct to the publishers, Headley Brothers, 109 Kingsway, London, WC2, England. D Journal of Laryngology and Otology, 1963

#### CONTENTS

	PAGE
Primary Amyloidosis of the Upper Air Passages. J. C. McAlpine, A. Radcliffe and I. Friedmann (London)	I
Mycotic Infection of the Frontal Sinus, D. A. Osborn (London)	29
Nasal Glioma. James Lister (London)	34
TREATMENT OF MALIGNANT DISEASE OF THE NASAL SINUSES BY SUPERVOLTAGE RADIOTHERAPY. Thomas J. Deeley and Robert Morrison (London)	43
THE EARLY MANAGEMENT OF BELL'S PALSY BY CERVICAL SYMPATHETIC BLOCK. F. Boyes Korkis (London)	50
Clinical Records—	
Waardenburg's Syndrome. J. Calinikos (Johannesburg) .	<b>5</b> 9
CHRONIC MIDDLE-EAR INFECTION—MASTOID SURGERY WITHOUT MEATAL FLAPS OR SKIN GRAFTS. Hector A. Thomas (Cardiff)	63
Septal Dermoplasty: A New Operative Procedure for Control of Nosebleeds in Patients with Hereditary Hæmorrhagic Telangiectasia. William H. Saunders	
(Ohio)	69
Societies' Proceedings—	
ROYAL SOCIETY OF MEDICINE—SECTION OF LARVINGOLOGY .	77
General Notes	95

For Advertisement space in this Journal apply to: HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2



And the reply was  $\mu\alpha\rho\nu\delta\iota\alpha$  which is hardly surprising when you consider that in that helmet he could hardly have heard him anyway.

Fortunately, few, if any, patients wear Grecian helmets these days and most are delighted to have the name of an independent hearing aid consultant of impeccable reputation murmured tactfully into their ears.

Today more people than ever before can benefit from a carefully selected aid. But careful selection is important. INGRAMS, the Independent Suppliers, have twenty-six different makes—the largest selection in the country—for patients to choose from. Couple this with expert impartial advice and a high reputation and you have the reason why so many E.N.T. Specialists refer their patients to Ingrams.



Domiciliary visits in any part of the country



# ORBENIN (BRL) SODIUM GLOXACILLIN

# ORAL PENICILLIN ACTIVE AGAINST RESISTANT STAPHYLOCOCCI

**ORBENIN** is indicated for immediate use in known or suspected resistant staphylococcal infections, and in mixed Gram-positive infections (e.g. infected burns) where resistant staphylococci may be present.

- Orbenin is stable to staphylococcal penicillinase and highly effective against 'resistant' staphylococci. Its activity is four times that of methicillin.
- Orbenin is also highly effective against other Gram-positive organisms, including streptococci, pneumococci and penicillin-sensitive staphylococci.
- Orbenin is bactericidal; it is acid-stable and well absorbed by either the oral or intramuscular routes.

Dosage by mouth is 500 mg. 6 hourly. Dosage by intramuscular injection is 250 mg. four to six hourly.



#### ORBENIN

is a product of British research at the BEECHAM RESEARCH LABORATORIES LTD BRENTFORD · ENGLAND · TELEPHONE: ISLeworth 4111

## 'PETERS' **NEW** CLINIC AUDIOMETER

FOR PURE-TONE AND SPEECH



#### **CONSIDER THESE FEATURES:**

- A continuous frequency range with continuous automatic threshold compensation.
- Twin continuously variable attenuators accurate to 1 db.
- Narrow band masking of pure tones plus insert receiver.
- Manual or automatic completely silent interruption.
- Loudness Balance and Difference Limen Recruitment Tests.
- High Fidelity Speech Circuit with Accessories for Free Field Speech Audiometry.
- Full Spectrum of White Noise for Masking Speech.
- Completely Transistorised for reliability.

The Westrex Company Limited, who are leading suppliers of Deaf Education Equipment, have a nationwide servicing organisation. In the immediate future their engineers in the field will be co-operating in the speedy servicing of all "Peters" audiometers in Great Britain.

Write for a descriptive brochure giving full details of this magnificent instrument and ask for a demonstration from either:

## WESTREX COMPANY Ltd. 152 COLES GREEN ROAD, LONDON, N.W.2

#### ALFRED PETERS & SONS Ltd.

VICTORIA WORKS, GELL STREET, SHEFFIELD 3

### ROYAL EAR HOSPITAL PATTERN ULTRASONIC EQUIPMENT

for the conservative surgery of Ménière's Disease. (Arslan Technique.) U.K. and U.S.A. patents applied for. Now available with the following special advantages:

TRANSDUCER and cable will withstand sterilization in autoclave.

Low operating voltage of transducer.

APPLICATORS can be interchanged to suit anatomy of middle ear.

Applicator tip normally  $2\frac{1}{2}$  mm. in diameter.

Intensities up to 100 Watts per sq. cm., giving ample power for pituitary irradiation in carcinoma.

GENERATOR operates on 90-250 volts 50 or 60 cycles.

Weight 50 lb. Size 21 by 14 by 12 in.

Tuning and intensity can be checked continuously.

#### REFERENCES:

Arslan. Proc. Fifth Int. Congress Oto-Rhino-Laryngol., Amsterdam, 1953.

Proc. roy. Soc. Med., 1958, 51, 617.

J. Laryng., 1960, LXXIV, 628.

Ibid., 1960, LXXIV, 730.

Ibid., 1961, LXXV, 345.

Ibid., 1962, LXXVI, 299.

Made in England by

FRISTON ELECTRONICS LTD., 25 CHURCH ROAD, RICHMOND, SURREY.

# Whatever the Season...

'Actifed' relieves nasal and respiratory congestion from the inside. It is taken by mouth and reaches areas inaccessible to sprays and nasal drops.

Rapidity of action is an outstanding feature of 'Actifed' products—relief of symptoms is felt within 10 to 30 minutes.

'Actifed' combines the most potent antihistamine, triprolidine, with pseudoephedrine, a powerful vaso-constrictor and bronchodilator. Unlike its analogue ephedrine, pseudoephedrine causes no marked central stimulation.

With 'Actifed' there is no risk of secondary engorgement or inhibitions of ciliary activity. 'Actifed' Tablets are recommended for adults, while the Syrup is particularly suitable for children. Also available—'Actifed' Compound Linctus, which contains codeine in addition to the other two ingredients, for relief of coughs or coughs and congestion.

# 'Actifed' TABLETS & S

clear Respiratory Congestion



Burroughs Wellcome & Co. (The Wellcome Foundation Ltd.) London