

SPARSITY OF ITERATIONS OF RATIONAL FUNCTIONS ON THE UNIT CIRCLE

IGOR E. SHPARLINSKI 

(Received 23 January 2026; accepted 11 February 2026)

Abstract

We use a recent result of Orevkov and Pakovich [‘On intersection of lemniscates of rational functions’, *Arnold J. Math.* **11** (2025), 7–26] to obtain a bound on the frequency with which elements in an orbit of polynomial iterations can fall on the unit circle.

2020 *Mathematics subject classification*: primary 37F10.

Keywords and phrases: iterations of rational functions, unit circle.

1. Introduction

Given a rational function $f \in \mathbb{C}(X)$ and an initial point $\alpha \in \mathbb{C}$, we consider the orbit

$$\mathcal{O}_f(\alpha) = (f^n(\alpha) : n = 0, 1, \dots),$$

where we set $f^0(X) = X$ and then define recursively

$$f^n(X) = f(f^{n-1}(X)), \quad n = 1, 2, \dots$$

We also assume that the orbit $\mathcal{O}_f(\alpha)$ terminates if α is a pole of $f^n(X)$ for some n .

We emphasise that we treat orbits as naturally ordered sequences, rather than as sets. If this sequence becomes eventually periodic, we call α *preperiodic*.

There is a vast amount of work studying the distribution of orbits $\mathcal{O}_f(\alpha)$ in various subsets $\mathcal{S} \subseteq \mathbb{C}$, under various algebraic and arithmetic constraints on the function f , the initial point α , and the set \mathcal{S} (see [1, 2, 4, 5, 7–10] and the references therein).

Here, we use a recent result of Orevkov and Pakovich [6] to study an apparently new question and estimate the frequency of the elements in an orbit $\mathcal{O}_f(\alpha)$, which fall on the unit circle

$$\mathbb{U} = \{z \in \mathbb{C} : |z| = 1\}.$$

During the preparation of this work, the author was supported in part by the Australian Research Council Grant DP230100530.

© The Author(s), 2026. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Some arithmetic analogues of this question have been considered by Ostafe [7], where a rational function $f \in \mathbb{K}(X)$ is defined over a number field \mathbb{K} and the unit circle is replaced by the set $\mathcal{U}_{\mathbb{K}}$ of roots of unity of \mathbb{K} . In particular, in this setting, using [7, Theorem 1.2], one can derive the finiteness of the sets $\mathcal{O}_f(\alpha) \cap \mathcal{U}_{\mathbb{K}}$ (under some natural conditions on α and f).

2. Main result

Given an integer N , we define

$$T_f(\alpha, N) = \#\{n : f^n(\alpha) \in \mathbb{U}, 1 \leq n \leq N\}.$$

We say that $f(z) \in \mathbb{C}(z)$ is *special* if $f^n(\mathbb{U}) \subseteq \mathbb{U}$ for some n , that is, f^n maps \mathbb{U} to \mathbb{U} .

For example, if $f(z)$ is a polynomial, then as follows from the discussion in [6, Section 1], in particular, [6, (3)], this is possible only if $f^n(z)$ is a monomial, that is, $f^n(z) = z^k$ for some integer $k \geq 0$ and thus $f(z)$ is also a monomial.

THEOREM 2.1. *For any nonspecial rational function $f(z) \in \mathbb{C}(z)$ and an initial point $\alpha \in \mathbb{C}$, which is not preperiodic, we have*

$$T_f(\alpha, N) = O(N/\log N),$$

where the implied constant depends only on $\deg f$.

As we have mentioned, nonmonomial polynomials are always nonspecial. However, it is not immediately clear how to characterise nonspecial rational functions, which we pose as an open question.

3. Preliminaries

The following result is a special case of [6, Theorem 1.1], which improves a similar bound from [11] on the number of intersections of lemniscates.

LEMMA 3.1. *Let $F(z) \in \mathbb{C}(z)$ be a rational function of degree D such that $F(\mathbb{U}) \not\subseteq \mathbb{U}$. Then, $\#\{z \in \mathbb{U} : F(z) \in \mathbb{U}\} \leq 2D$.*

We also need the following simple combinatorial statement given by [3, Lemma 5.7].

LEMMA 3.2. *Let $2 \leq M < N/2$. For any sequence $0 \leq n_1 < \dots < n_M \leq N$, there exists $r \leq 2N/(M-1)$ such that $n_{i+1} - n_i = r$ for at least $(M-1)^2/4N$ values of $i \in \{1, \dots, M-1\}$.*

4. Proof of Theorem 2.1

We fix some $\alpha \in \mathbb{C}$, which is not preperiodic, and define $z_n = f^n(\alpha)$, $n = 0, 1, \dots$, which are obviously pairwise distinct.

If $T_f(\alpha, N) = 0$, there is nothing to prove. Thus, we now assume that $T_f(\alpha, N) \geq 1$.

For $t = T_f(\alpha, N) - 1$, let $1 \leq n_1 < \dots < n_{t+1} \leq N$ be such that $z_{n_i} \in \mathbb{U}$. Consider the gaps $d_i = n_{i+1} - n_i$, $i = 1, \dots, t$.

By Lemma 3.2, there is some $r \leq 2N/t$ such that $d_i = r$ for at least $I \geq t^2/(4N)$ values of $i \in \{1, \dots, t\}$. Hence, there are at least I solutions to the equation

$$|f^r(z)| = |z| = 1$$

given by $z_i = f^{n_i}(\alpha)$ for each i such that $d_i = r$. Note that since $\alpha \in \mathbb{C}$ is not preperiodic, all these points are distinct.

Since $\deg f^r \leq (\deg f)^r$ and f is nonspecial, by Lemma 3.1,

$$\frac{t^2}{4N} \leq I \leq 2(\deg f)^r \leq 2(\deg f)^{2N/t}. \quad (4.1)$$

If

$$2(\deg f)^{2N/t} \leq N^{1/2}, \quad (4.2)$$

then from (4.1), we have $t \leq 2N^{3/4}$, which is much stronger than the desired bound. Otherwise, that is, if (4.2) is false,

$$\frac{2N}{t} \log(\deg f) + \log 2 > \frac{1}{2} \log N,$$

which implies the result.

Acknowledgement

The author is grateful to the referee for the careful reading of the manuscript.

References

- [1] J. P. Bell, S. Chen and E. Hossain, ‘Rational dynamical systems, S -units, and D -finite power series’, *Algebra Number Theory* **15** (2021), 1699–1728.
- [2] E. Chen, ‘Avoiding algebraic integers of bounded house in orbits of rational functions over cyclotomic closures’, *Proc. Amer. Math. Soc.* **146** (2018), 4189–4198.
- [3] C. D’Andrea, A. Ostafe, I. Shparlinski and M. Sombra, ‘Reductions modulo primes of systems of polynomial equations and algebraic dynamical systems’, *Trans. Amer. Math. Soc.* **371** (2019), 1169–1198.
- [4] A. Ferraguti, A. Ostafe and U. Zannier, ‘Cyclotomic and abelian points in backward orbits of rational functions’, *Adv. Math.* **438** (2024), 109463.
- [5] J. Mello, ‘Cyclotomic preperiodic points for morphisms in affine spaces and preperiodic points with bounded house and height’, *Funct. Approx. Comment. Math.* **67** (2022), 259–279.
- [6] S. Orevkov and F. Pakovich, ‘On intersection of lemniscates of rational functions’, *Arnold J. Math.* **11** (2025), 7–26.
- [7] A. Ostafe, ‘On roots of unity in orbits of rational functions’, *Proc. Amer. Math. Soc.* **145** (2017), 1927–1936.
- [8] A. Ostafe, L. Pottmeyer and I. E. Shparlinski, ‘Perfect powers in value sets and orbits of polynomials’, *New York J. Math.* **27** (2021), 903–917.
- [9] A. Ostafe, M. Sha, I. E. Shparlinski and U. Zannier, ‘On multiplicative dependence of values of rational functions and a generalization of the Northcott theorem’, *Michigan Math. J.* **68** (2019), 385–407.

- [10] A. Ostafe and M. Young, 'On algebraic integers of bounded house and preperiodicity in polynomial semigroup dynamics', *Trans. Amer. Math. Soc.* **373** (2020), 2191–2206.
- [11] F. Pakovich and I. E. Shparlinski, 'Level curves of rational functions and unimodular points on rational curves', *Proc. Amer. Math. Soc.* **148** (2020), 1829–1833.

IGOR E. SHPARLINSKI, School of Mathematics and Statistics,
University of New South Wales, Sydney, New South Wales 2052, Australia
e-mail: igor.shparlinski@unsw.edu.au