

SPIN CORRELATION IN BINARY SYSTEMS

N. Farbiash¹ and R. Steinitz¹

RESUMEN

Examinamos la correlación entre las velocidades de rotación proyectadas en sistemas binarios. Este estudio es una extensión de un trabajo previo (Steinitz y Pyper, 1970; Levato, 1974). Una base de datos más amplia y nuevas pruebas nos permiten concluir que, en efecto, existe una correlación entre las velocidades de rotación proyectadas de las binarias. Sugerimos, de hecho, que los espines están correlacionados.

ABSTRACT

We examine the correlation of projected rotational velocities in binary systems. It is an extension of previous work (Steinitz and Pyper, 1970; Levato, 1974). An enlarged data basis and new tests enable us to conclude that there is indeed correlation between the projected rotational velocities of components of binaries. In fact we suggest that spins are already correlated.

Key Words: **BINARIES: GENERAL — STARS: ROTATION**

1. INTRODUCTION

Evolution of binary systems could be a result of two main processes: three body collision, or evolving of binaries in one disk. The probability for three body collision is extremely small; thus it is more likely that binary systems evolve in one disk. In that case we expect to find correlation between the measured $V_e \sin(i)$ values of the members of such a system. Therefore, we study the degree of projected rotational velocity correlation between members of binary systems.

Slettebak (1963) did not find any significant difference between mean rotational velocity distribution of members of binary systems and those of single stars. Further, Abt (2001) concluded that spin axes are probably randomly oriented. On the other hand, Steinitz and Pyper (1970) concluded that some correlation of projected rotational velocities is present for the components of visual binaries. Levato (1974) also discussed this issue for visual binaries and close binary systems and found that there is indeed correlation of projected rotational velocities in binary systems. We now extend the original study (Steinitz and Pyper, 1970), which included only 50 systems. The significance of our results stems from the use of 1010 binary systems.

Actually, we will examine three samples of binary systems, as defined in the next section.

2. DATA

The Catalogue of Stellar Projected Rotational Velocities (Glebocki, Gnacinski & Stawikowski 2000)

is our source for the sample of binaries, chosen by imposing the following criteria:

1. The spectral type of both components is earlier than A9 (slow rotation of stars later than A9 would automatically simulate correlation).
2. Giants and supergiants may have lost their original rotational velocities. So we select only those binaries both of whose components are on the main sequence.
3. Multiple systems including more than two stars are excluded.

We are now left with a sample of 1010 real binary (RB) systems. Since the mean rotational velocity along the main sequence changes, it could happen that the choice of a sample restricted in spectral type will automatically exhibit the correlation we are looking for. To eliminate this possibility, and show that projected rotational velocity correlation is present in real binary systems only, we use two artificial samples of binaries. Define sample AB (Artificial Binaries) through shuffling the components of the real systems, and exclude the real ones (containing 2038180 systems). The second sample ABR (Artificial Binaries, Restricted), is obtained from sample AB by eliminating of all pairs having spectral type difference larger than two spectral subclasses (containing 263344 systems). This restriction is more stringent than the one we admit for the basic, real systems.

¹Ben-Gurion University, Beer-Sheva, 84105, Israel.

3. ILLUSTRATION

In Fig.1. we plot the projected rotational velocities of one component against the other one, for all the three samples previously defined. From Fig.1. we see that samples AB and ABR do not exhibit any correlation whatsoever. In contrast, the correlation in sample RB is clearly evident. To quantify this result we apply more tests in the next section.




Fig. 1. Illustration of the projected rotational velocities of one component against the other for the samples AB (top), ABR (middle), and RB (bottom).

4. ANALYSIS

4.1. Bivariate distribution, marginal distribution, and linear regression

We prepare a table for each of the three samples, AB, ABR and RB: it gives the discrete bivariate distribution of the samples. We give also the regression of the mean velocity of one component on the other one, as defined by equation (1).

$$\bar{u}_1(u_2(j)) = \frac{\sum_{i=1}^7 u_1(i) F(u_1(i), u_2(j))}{\sum_{i=1}^7 F(u_1(i), u_2(j))} \quad (1)$$

Here $u_1 = v_1 \sin i_1$ is the projected rotational velocity of the component, and $F(u_1, u_2)$ is the bivariate distribution.

For better illustration we show the regression of projected rotational velocity distribution of one component on the other one for each of the three samples. The $F(u_1, u_2)$ distributions for the samples AB, ABR, and RB are given in Table 1. The marginal distribution and the regression lines are also shown. The latter are plotted in Fig.2. For the ease of comparing the current results with previous ones, the velocity range has been divided into subintervals of 50 km s^{-1} .

No significant differences between the main diagonal and other components of the table are evident for the samples AB and ABR. But the set RB shows a distinct difference between the main diagonal and other elements of the table, indicating the presence of correlation of projected spins in real binaries.




Fig. 2. Regression of the mean rotational velocities of one component (ordinate) as a function of the rotational velocity interval of the second component (abscissa) for samples AB (top), ABR (middle), and RB (bottom).

TABLE 1

DISCRETE DISTRIBUTION FUNCTION $F(Y_1, Y_2)$ OF THE PROJECTED ROTATIONAL VELOCITIES OF THE COMPONENTS OF SAMPLE AB (TOP), ABR(MIDDLE), AND RB(BOTTOM).

u_2	u_1	0-50	50-100	100-150	150-200	200-250	250-300	300-350	$\phi_2(u_2)$	$\bar{u}_1(u_2)$
0-50		563	525	406	351	200	90	51	0.22	115
50-100		617	575	446	388	221	99	58	0.24	116
100-150		451	420	324	282	161	73	42	0.18	115
150-200		433	404	313	271	155	70	41	0.17	116
200-250		276	260	200	174	99	45	26	0.11	116
250-300		122	115	88	77	44	20	12	0.05	116
300-350		104	99	76	67	39	17	10	0.04	117
$\phi_1(u_1)$		0.26	0.24	0.19	0.16	0.09	0.04	0.02		
$\bar{u}_2(u_1)$		125	126	126	126	126	126	127		
0-50		576	496	385	356	205	105	0	0.21	112
50-100		657	627	466	412	215	94	0	0.25	108
100-150		452	447	331	277	141	56	22	0.17	110
150-200		465	454	335	276	143	57	22	0.18	109
200-250		290	264	201	172	96	45	18	0.11	112
250-300		115	99	80	65	37	19	7	0.04	112
300-350		115	88	74	66	43	25	10	0.04	119
$\phi_1(u_1)$		0.27	0.25	0.19	0.16	0.09	0.04	0.01		
$\bar{u}_2(u_1)$		126	125	126	124	125	125	201		
0-50		1901	198	20	0	0	0	0	0.21	31
50-100		614	1564	158	40	10	0	0	0.24	68
100-150		40	515	1000	158	0	0	0	0.17	112
150-200		69	79	574	752	158	59	10	0.17	156
200-250		10	50	99	535	347	50	40	0.11	190
250-300		0	0	30	59	228	119	30	0.05	231
300-350		0	10	10	50	129	198	89	0.05	254
$\phi_1(u_1)$		0.26	0.24	0.19	0.16	0.09	0.04	0.02		
$\bar{u}_2(u_1)$		43	89	144	193	242	278	284		

^a u_1 and u_2 are given in units of km s^{-1} .

4.2. Modified convolution test

Further illustration of the correlation between the rotational velocity distribution in real binary systems, and not in artificial ones, is to evaluate the modified convolution of the distribution functions. To validate the result, we first look at the single star velocity distribution, as shown in Fig. 3. The distribution is rather flat and does not indicate a strong maximum at any specific speed. Our modified convolution is essentially $g(u) = \int P(v - u|v)dv$, which in the discrete case is:

$$g(u) = \sum_v F(v - u, v). \tag{2}$$

As expected from the flatness of the distribution in Fig. 3., the convolution of artificial binary systems do not indicate a concentrated, sharp peak. However, for real binaries, there is a rather sharp peak present. It means that probability for the projected rotational velocity difference to be smaller than 50 km s^{-1} is almost 60%. The probability for the difference to be smaller than 100 km s^{-1} is more than 90%(!) in real binaries.




Fig. 3. Projected rotational velocity distribution of all components grouped into intervals of 50 km s^{-1} .




Fig. 4. Modified convolution of rotational velocities.

5. CONCLUSIONS

We examined three samples of binaries. For each of the samples we applied identical tests, to render significant comparisons between artificial binaries and real ones.

As expected, correlation in artificial systems (Set AB and ABR) is insignificant. This is indicated by the minuscule slope of the regression lines. However, correlation for the binary set RB (Real Binaries) is clearly evident ($u_1 = .856u_2 + 32$). One can describe our results roughly as follows:

$$v_1 \sin i_1 \cong v_2 \sin i_2. \quad (3)$$

This relation can be understood either as:

1. $v_1 \ll v_2$ while $\sin i_1 \gg \sin i_2$,
- or
2. $v_1 \simeq v_2$ as well as $\sin i_1 \simeq \sin i_2$.

Since we have used a sample containing 1010 systems, the probability of the first case being real is extremely small. We rather accept the second explanation. We interpret this as meaning that:

1. Spin axes of members in binary systems are roughly parallel.
2. Rotational speeds are correlated.

There have been several investigations related to binary systems (Giuricin, Mardirossian & Mezzetti, 1984a,b; Levato, 1976; Zahn, 1977; and Pan, 1997). These papers point out the importance of tidal interaction in close binary systems (especially the theory given by Zahn, 1966a,b,c, 1970, 1975, 1977).

We suppose that only a small fraction of the sample RB contains close binary systems. Thus we lack a general theory which can account for the empirical results demonstrated here.

REFERENCES

- Abt, H. A. 2001, *AJ*, 122, 2008
 Giuricin, G., Mardirossian, F., & Mezzetti, M. 1984a, *A&A*, 131, 152
 Giuricin, G., Mardirossian, F., & Mezzetti, M. 1984b, *A&A*, 135, 393
 Glebocki, R., Gnacinski, P., & Stawikowski, A. 2000, *AcA*, 50, 509
 Levato, H. 1974, *A&A*, 35, 259
 Levato, H. 1976, *ApJ*, 203, 680
 Pan, K. 1997, *A&A*, 321, 202
 Slettebak, A. 1963, *ApJ*, 138, 118
 Steinitz, R., & Pyper, D. M., 1970, in *Stellar Rotation*, ed. A. Slettebak (IAU Coll. 4) (New York: Gordon & Breach), 165
 Zahn, J.-P. 1966a, *AnAp*, 29, 313
 Zahn, J.-P. 1966b, *AnAp*, 29, 489
 Zahn, J.-P. 1966c, *AnAp*, 29, 685
 Zahn, J.-P. 1970, *A&A*, 4, 452
 Zahn, J.-P. 1975, *A&A*, 41, 329
 Zahn, J.-P. 1977, *A&A*, 57, 383

DISCUSSION

Sterzik – Could the spins be preferentially anti-parallel oriented to reach the same conclusion?


Farbiash – We have no practical way to say if spins are parallel or anti-parallel.

Mathieu – How does the correlation that you find depend on binary separation? Can you distinguish between axis alignment due to tidal effects as compared to other effects, such as formation?

Farbiash – We find very small differences between the correlation for spectroscopic binaries (~ 100 systems), visual ones (~ 80 systems), and other binaries we used in this study. The tidal effect is very important for close binaries, and is the main reason for synchronization among these systems. The reason we have made the Omega test is to see how many systems are already synchronized. We find that very few systems in the sample tested are synchronized. Data on the periods of the systems we tested are required in order to get a clear answer to the dependence between spin correlation and the binary separation.

Hanawa – Can you separate the correlation in $\sin i_j - \sin i_k$ and that in V_1 and V_2) by assuming, for example, that $\sin i$ has a random distribution?

Farbiash – The assumption that $\sin(i)$ is distributed randomly is probably true (Abt 2001). With the assumption that the spin axes are perfectly parallel (which is a very strong assumption!) we get a theoretical correlation value of ≈ 1 for the rotation velocity. I have to be very careful saying that, since this value is meaningless if the spins are not perfectly parallel.



Cathie Clarke and Bo Reipurth.